

Conservation of Ramsey's theorem for pairs and well-foundedness

Quentin Le Hou  rou Ludovic Levy Patey
Keita Yokoyama

February 18, 2024

Abstract

In this article, we prove that Ramsey's theorem for pairs and two colors is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$ and over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \bigcup_n \text{WF}(\omega_n^\omega)$. These results improve theorems from Chong, Slaman and Yang [6] and Ko  dziejczyk and Yokoyama [20] and belong to a long line of research towards the characterization of the first-order part of Ramsey's theorem for pairs.

1 Introduction

We investigate the proof-theoretic strength of Ramsey's theorem for pairs and some of its consequences, by proving that they are Π_1^1 -conservative over the base theory of Reverse Mathematics (RCA_0), augmented with the collection principle for Σ_2^0 formulas ($\text{B}\Sigma_2^0$) and the assumption that some appropriate ordinals are well-founded ($\text{WF}(\alpha)$). This article belongs to a long line of research towards the characterization of the first-order part of Ramsey's theorem for pairs, and is part of a larger foundational program known as Reverse Mathematics, and whose goal is to find optimal axioms for proving ordinary theorems. See Simpson [30] or Dzhafarov and Mummert [7] for a reference book on Reverse Mathematics, and Hirschfeldt [8] to a gentle introduction to the reverse mathematics of combinatorial principles.

Ramsey's theorem is the founding theorem of Ramsey's theory, a field of research which studies the conditions under which structure appears among a sufficiently large amount of data. We identify a set k with its ordinal $\{0, 1, \dots, k-1\}$. Given a set $X \subseteq \mathbb{N}$ and some integer $n \in \mathbb{N}$, we let $[X]^n$ denote the set of all unordered n -tuples over X . Given a coloring $f : [\mathbb{N}]^n \rightarrow k$, a set $H \subseteq \mathbb{N}$ is *f-homogeneous* (for color $i < k$) if $f(\sigma) = i$ for every $\sigma \in [H]^n$.

Statement 1.1 (Ramsey's theorem). *Given $n, k \in \mathbb{N}$, RT_k^n is the statement "Every coloring $f : [\mathbb{N}]^n \rightarrow k$ admits an infinite f-homogeneous set".*

The proof-theoretic strength of Ramsey's theorem for triples and larger size of tuples is well-understood: RT_k^n is equivalent to the Arithmetic Comprehension

Axiom (ACA_0) over RCA_0 whenever $n \geq 3$ and $k \geq 2$ (see Jockusch [18] for a computability-theoretic equivalence, and Simpson [30, Theorem III.7.6] for a proof-theoretic version of it). The first-order part of ACA_0 is precisely Peano Arithmetic.

Ramsey's theorem for pairs and two colors (RT_2^2) can be phrased as stating, for every infinite graph, the existence of an infinite subset of vertices such that the induced subgraph is either a clique or an anticlique. RT_2^2 is known to be strictly weaker than ACA_0 over RCA_0 (see Seetapun and Slaman [29]), and incomparable with weak König's lemma (WKL_0), the restriction of König's lemma to binary trees (see Liu [24]). The characterization of the first-order part of Ramsey's theorem for pairs and two colors is a long-standing open question.

Definition 1.2 (Collection principle). *Given a family Γ of formulas, $\text{B}\Gamma$ is the statement “For every formula $\varphi \in \Gamma$, and every $a \in \mathbb{N}$, if $\forall x < a \exists y \varphi(x, y)$, then there is some $b \in \mathbb{N}$ such that $\forall x < a \exists y < b \varphi(x, y)$ ”.*

Ramsey's theorem for pairs implies $\text{B}\Sigma_2^0$ over RCA_0 (see Hirst [16]). The following question is a major open problem in Reverse Mathematics:

Question 1.3. *Is RT_2^2 a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0$?*

1.1 Well-foundedness and bounded monotone enumerations

In Computability Theory, the ability to compute fast-growing functions can be seen as a unified framework to measure the strength of some Turing degrees, e.g., hyperimmune degrees, high degrees, and degrees admitting a modulus. In Proof Theory, the strength of a theory is better measured by the ability to prove the well-foundedness of some ordinals. The Π_1^1 *proof-theoretic ordinal* of a theory T is the least ordinal α such that T does not prove its well-foundedness ($\text{WF}(\alpha)$). For instance, the Π_1^1 proof-theoretic ordinals of RCA_0 and ACA_0 are ω^ω and ϵ_0 , respectively (see Simpson [30]).

The well-foundedness of ordinals naturally occurred in the proof-theoretic analysis of Ramsey's theorem for pairs. Chong, Slaman and Yang [6] proved that RT_2^2 does not imply the induction scheme for Σ_2^0 formulas ($\text{I}\Sigma_2^0$) over RCA_0 using a model of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \neg \text{I}\Sigma_2^0$ which furthermore satisfied a principle called Bounded Monotone Enumeration (BME).

Definition 1.4 ([6]). *Let E be a procedure to recursively enumerate a finite branching enumerable tree. We say that E is a monotone enumeration if and only if the following conditions apply to its stage-by-stage behavior.*

- (1) *The empty sequence is enumerated by E during stage 0.*
- (2) *Only finitely many sequences are enumerated by E during any stage.*
- (3) *Suppose $T[s]$ is the tree enumerated by E at the end of stage s . Then any sequence enumerated by E during stage $s + 1$ must extend some terminal node of $T[s]$.*

A monotone enumeration E is b -bounded if every node in E has length at most b .

Statement 1.5. BME_\star is the statement “Every bounded monotone enumeration is finite”.

The statement BME is an iterated version of BME_\star . Its formal definition is rather technical and will not be used within this article. Kreuzer and Yokoyama [22] proved that BME_\star is equivalent to $\text{WF}(\omega^\omega)$ over IS_1^0 , and that BME is equivalent to $\bigcup_k \text{WF}(\omega_k^\omega)$, where $\omega_0^\alpha = \alpha$ and $\omega_{k+1}^\alpha = \omega_k^{\omega_k^\alpha}$.

The principles BME and BME_\star naturally occur in the proofs of conservation of Ramsey’s theorem for pairs. On the other hand, Patey and Yokoyama [28] proved that RT_2^2 is $\forall \Pi_3^0$ -conservative over $\text{RCA}_0 + \text{BS}_2^0$, which implies in particular that the Π_1^1 proof-theoretic ordinal of RT_2^2 is ω^ω , hence that RT_2^2 does not imply BME_\star over RCA_0 .

In this article, we show that the separation of Chong, Slaman and Yang [6] of RT_2^2 from IS_2^0 only depends on the assumption that the original model satisfies $\text{BS}_2^0 + \text{BME}$, by proving the following conservation theorems.

Main Theorem 1.6. Fix $n \geq 2$. Let T_n be either BS_n^0 or IS_n^0 . $\text{RCA}_0 + T_n + \text{RT}_2^2 + \text{WKL} + \text{WF}(\epsilon_0)$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + T_n + \text{WF}(\epsilon_0)$.

Main Theorem 1.7. Fix $n \geq 2$. Let T_n be either BS_n^0 or IS_n^0 . $\text{RCA}_0 + T_n + \text{RT}_2^2 + \text{WKL} + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega)$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega)$.

This strengthens a result from Kołodziejczyk and Yokoyama [20] who showed that $\text{RCA}_0 + \text{RT}_2^2 + \text{BME}$ is a Π_4^0 -conservative extension of $\text{RCA}_0 + \text{BS}_2^0 + \text{BME}$. By Cholak, Jockusch and Slaman [3], the result holds when $\text{BS}_2^0 + \text{WF}(\epsilon_0)$ is replaced by IS_2^0 . Thus, conservations over $\text{RCA}_0 + (\text{WF}(\epsilon_0) \vee \text{IS}_2^0)$ and over $\text{RCA}_0 + (\bigcup_{n \in \omega} \text{WF}(\omega_n^\omega) \vee \text{IS}_2^0)$ are the best known results.

In order to better understand the computational and proof-theoretic content of Ramsey’s theorem for pairs, two main decompositions of RT_2^2 into combinatorially weaker statements have been studied. These decompositions will be used to prove separate conservation theorems for simpler statements, and then combined to obtain our main theorems. We now detail those statements.

1.2 Tournaments and linear orders

Ramsey’s theorem for pairs and two colors can be decomposed into the Erdős-Moser theorem about infinite tournaments, and the Ascending Descending Sequence principle.

Definition 1.8. A tournament on a domain D is an irreflexive binary relation $R \subseteq D^2$ such that for every $a, b \in D$ with $a \neq b$, exactly one of $R(a, b)$ and $R(b, a)$ holds. A tournament R is transitive if for every $a, b, c \in D$, if $R(a, b)$ and $R(b, c)$ then $R(a, c)$.

A subtournament of $R \subseteq \mathbb{N}^2$ is fully characterized by its domain and the parent tournament R . We therefore identify subtournaments with infinite sets $H \subseteq \mathbb{N}$. The following statement is known as the Erdős-Moser theorem, and is an infinitary version of some theorem by Erdős and Moser [9].

Statement 1.9 (Erdős-Moser). *EM is the statement “Every infinite tournament admits an infinite transitive subtournament”.*

There exists a one-to-one correspondence between a tournament R and a coloring $f : [\mathbb{N}]^2 \rightarrow 2$, by letting $f(\{x, y\}) = 1$ if $(R(x, y) \leftrightarrow x <_{\mathbb{N}} y)$. EM can be restated as “for every coloring $f : [\mathbb{N}]^2 \rightarrow 2$, there exists an infinite f -transitive subset H , that is, for every $x, y, z \in H$ such that $x < y < z$ and every $i < 2$, then if $f(\{x, y\}) = f(\{y, z\}) = i$, we have $f(\{x, z\}) = i$ ”.

Statement 1.10 (Ascending Descending Sequence). *ADS is the statement “Every infinite linear order admits an infinite ascending or descending subsequence”.*

Here again, there is a one-to-one correspondence between linear orders $\mathcal{L} = (\omega, <_{\mathcal{L}})$, and transitive colorings $f : [\mathbb{N}]^2 \rightarrow 2$, by letting $f(\{x, y\}) = 1$ if $(x <_{\mathbb{N}} y \leftrightarrow x <_{\mathcal{L}} y)$. Then ADS can be restated as “for every transitive coloring $f : [\mathbb{N}]^2 \rightarrow 2$, there exists an infinite f -homogeneous set”. Using the reformulations of EM and ADS, the following proposition is immediate:

Proposition 1.11 (Bovyking and Weiermann [2]). $\text{RCA}_0 \vdash \text{RT}_2^2 \leftrightarrow \text{EM} \wedge \text{ADS}$

Proof. EM and ADS are both particular cases of RT_2^2 , hence $\text{RCA}_0 \vdash \text{RT}_2^2 \rightarrow \text{EM} \wedge \text{ADS}$. On the other hand, given a coloring $f : [\mathbb{N}]^2 \rightarrow 2$, first apply EM to obtain an infinite f -transitive subset $X = \{x_0 < x_1 < \dots\}$. Let $g : [\mathbb{N}]^2 \rightarrow 2$ be the transitive coloring defined by $g(a, b) = f(x_a, x_b)$. By ADS, there exists an infinite g -homogeneous subset $H \subseteq \mathbb{N}$. The set $\{x_a : a \in H\}$ is infinite and f -homogeneous. \square

From a combinatorial viewpoint, the decomposition of RT_2^2 into EM and ADS divides the difficulty of Ramsey’s theorem for pairs as follows: the combinatorics of EM are very close to those of RT_2^2 , but without the disjunctive nature of RT_2^2 . Thus, there is no need to work with pairing arguments. On the other hand, ADS has very simple combinatorics, and does not imply any form of compactness, but is a disjunctive statement.

From a reverse mathematical viewpoint, the decomposition is non-trivial, in that EM and ADS are both strictly weaker than RT_2^2 (see Lerman, Solomon and Towsner [23]).

From a proof-theoretic viewpoint, both statements imply BS_2^0 over RCA_0 (see Hirschfeldt and Shore [15, Proposition 4.5] and Kreuzer [21, Proposition 16]), and $\text{RCA}_0 + \text{ADS}$ is known to be Π_1^1 -conservative extension of $\text{RCA}_0 + \text{BS}_2^0$ (see Chong, Slaman and Yang [5]). By an amalgamation theorem of Yokoyama [33], it follows that $\text{RCA}_0 + \text{RT}_2^2$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{BS}_2^0$ iff so is $\text{RCA}_0 + \text{EM}$.

1.3 Cohesiveness and stability

The second (and historically the first) decomposition of RT_2^2 was done by Cholak, Jockusch and Slaman [3] in their seminal article on Ramsey's theorem for pairs. It consists in reducing RT_2^2 to a stable version of it (SRT_2^2), using the Cohesiveness principle (COH). Then, it is easy to see that SRT_2^2 is equivalent to the infinite pigeonhole principle for Δ_2^0 instances, which is a combinatorially simple statement.

Definition 1.12. A coloring $f : [\mathbb{N}]^2 \rightarrow k$ is stable if for every $x \in \mathbb{N}$, $\lim_y f(x, y)$ exists. Given an infinite sequence of sets $\vec{R} = R_0, R_1, \dots$, an infinite set C is \vec{R} -cohesive if for every $s \in \mathbb{N}$, $C \subseteq^* R_s$ or $C \subseteq^* \overline{R}_s$, where \subseteq^* denotes inclusion up to finite changes.

Statement 1.13 (Stable Ramsey's theorem for pairs). SRT_k^2 denotes the restriction of RT_k^2 to stable colorings.

Statement 1.14 (Cohesiveness). COH is the statement "Every infinite sequence of sets has a cohesive set".

The following equivalence is essentially due to Cholak, Jockusch and Slaman [3], with the proof $\text{RCA}_0 \vdash \text{RT}_2^2 \rightarrow \text{COH}$ fixed by Mileti [25, Appendix A].

Proposition 1.15. $\text{RCA}_0 \vdash \text{RT}_2^2 \leftrightarrow \text{SRT}_2^2 \wedge \text{COH}$.

Proof. The proof of the forward direction will be omitted, as it will be of no use in the article. Let us prove that $\text{RCA}_0 \vdash \text{SRT}_2^2 \wedge \text{COH} \rightarrow \text{RT}_2^2$. Given a coloring $f : [\mathbb{N}]^2 \rightarrow 2$, define the sequence of sets $\vec{R} = R_0, R_1, \dots$ by $R_x = \{y : f(x, y) = 1\}$. Given an infinite \vec{R} -cohesive set $C = \{x_0 < x_1 < \dots\}$, let $g : [\mathbb{N}]^2 \rightarrow 2$ be defined by $g(\{a, b\}) = f(\{x_a, x_b\})$. By cohesiveness of C , the coloring g is stable, so by SRT_2^2 , there is an infinite g -homogeneous set H . The set $\{x_a : a \in H\}$ is infinite and f -homogeneous. \square

From a reverse mathematical viewpoint, the decomposition of RT_2^2 into SRT_2^2 and COH is also non-trivial (see Hirschfeldt et al. [14] and Monin and Patey [26]).

The computability-theoretic content of the cohesiveness principle is well-understood: it admits a maximally complex instance, which is the sequence of all primitive recursive sets. A cohesive set for this sequence is called *p-cohesive*. For every uniformly computable sequence of sets \vec{R} , every *p*-cohesive set computes an \vec{R} -cohesive set. Moreover, a Turing degree bounds a *p*-cohesive set iff its jump is PA over $0'$. See Jockusch and Stephan [17].

The idea of using cohesiveness to transform a coloring into a stable one can be applied to EM and ADS. We call SEM and SADS the restriction of EM and ADS to stable tournaments and linear orders, respectively. Stable linear orders can be of three types: $\omega + \omega^*$, $\omega + k$ and $k + \omega^*$, for some $k \in \omega$. The two former cases are computably true, so the only interesting case is whenever it is of order type $\omega + \omega^*$. This is why SADS is often defined as the statement "Every linear order of type $\omega + \omega^*$ admits an infinite ascending or descending sequence".

Combining the two kind of decompositions, we shall use the following equivalence:

Proposition 1.16. $\text{RCA}_0 \vdash \text{RT}_2^2 \leftrightarrow \text{SEM} \wedge \text{SADS} \wedge \text{COH}$

We will therefore prove conservation results about WKL , COH , SEM and SADS independently, and use the previous decomposition to obtain our main theorems.

1.4 Definitions and notation

Since $[X]^n$ is in one-to-one correspondence with the set of all n -tuples in increasing order, given a coloring $f : [\mathbb{N}]^n \rightarrow k$, we might omit the brackets, and simply write $f(x_0, x_1, \dots, x_{n-1})$ instead of $f(\{x_0, x_1, \dots, x_{n-1}\})$, assuming that $x_0 < \dots < x_{n-1}$.

A Π_2^1 *problem* P is a formula $\forall X[\varphi(X) \rightarrow \psi(X, Y)]$ where φ and ψ are arithmetic formulas. A set X such that $\varphi(X)$ holds is a *P-instance*, and a Y such that $\psi(X, Y)$ holds is a *P-solution* to X .

Theories. Let IS_n^0 be the following induction scheme for every Σ_n^0 formula ϕ :

$$\phi(0) \wedge \forall x[\phi(x) \rightarrow \phi(x+1)] \rightarrow \forall x\phi(x)$$

Let $\Delta_n^0\text{-CA}$ be the following comprehension scheme for every Σ_n^0 formula ϕ and every Π_n^0 formula ψ :

$$\forall x[\phi(x) \leftrightarrow \psi(x)] \rightarrow \exists A \forall x[x \in A \leftrightarrow \phi(x)]$$

RCA_0^* denotes the theory of Robinson arithmetic (Q) augmented with the comprehension axiom for Δ_1^0 predicates ($\Delta_1^0\text{-CA}$), the collection scheme for Σ_1^0 formulas (BS_1^0) and the totality of the exponentiation function (exp). RCA_0 is the theory of $Q + \Delta_1^0\text{-CA} + \text{IS}_1^0$. Note that RCA_0 entails RCA_0^* . The theory RCA_0 is the base theory of Reverse Mathematics, but RCA_0^* is sometimes more convenient to use, as shown by Belanger [1].

Models. A *model* is a second-order structure $\mathcal{M} = (M, S, +_M, \times_M, <_M)$, where M is the first-order part, that is, the set of integers and $S \subseteq \mathcal{P}(M)$ is the second-order part. We will usually omit the operations, and simply write $\mathcal{M} = (M, S)$. A model $\mathcal{M} = (M, S)$ is *topped* if there is some set $X \in S$ such that every $Y \in S$ is $\Delta_1^0(X)$. A model $\mathcal{N} = (N, T)$ is an ω -*submodel* of $\mathcal{M} = (M, S)$ if $N = M$ and $T \subseteq S$. Note that the notion of ω -submodel must not be confused with the notion of ω -model, in which $M = \omega$, that is, is the set of all standard integers.

Given a model $\mathcal{M} = (M, S)$ and a set $G \subseteq M$, we write $\mathcal{M}[G]$ for the model whose first-order part is M , and whose second-order part consists of all $\Delta_1^0(X, G)$ sets for $X \in S$. Therefore, if $\mathcal{M} \models \text{RCA}_0$, and $\mathcal{M} \cup \{G\} \models \text{IS}_1^0$, then $\mathcal{M}[G] \models \text{RCA}_0$.

A set X is M -bounded if there is some $x \in M$ such that $\forall y \in X, y \leq x$. A set X is M -finite or M -coded if there exists some $x \in M$ such that $x = \sum_{n \in X} 2^n$.

Given two sets X and Y , we write $X \gg Y$ to say that X is of PA degree over Y . More formally, it means that $\Delta_1^0(Y)$ infinite binary tree admits a $\Delta_1^0(X)$ path.

Ordinals. In this article, we shall work with ordinals smaller than ϵ_0 in models of weak arithmetic. By Cantor's normal form theorem, every ordinal α below ϵ_0 can be written uniquely as $\alpha = \omega^{\gamma_1} n_1 + \dots + \omega^{\gamma_k} n_k$ for some ordinals $\gamma_k < \dots < \gamma_1 < \alpha$ and positive integers n_1, \dots, n_k . So, every such ordinal can be represented as a finite rooted tree and therefore be coded by an integer which will be called its "index". For a suitable encoding of the finite rooted trees, the cantor normal form of any ordinal below ϵ_0 and the standard operations on ordinals (sums, product, comparison) are computable from their indexes. See Hájek and Pudlák [13, Chapter II.3] for a formal study of ordinals in the context of weak arithmetic.

1.5 Structure of the paper

In Section 2, we develop some basic techniques about Π_1^1 -conservation over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$. We prove in particular that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha) + \Gamma$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ for $\Gamma \in \{\text{COH}, \text{WKL}\}$ and α any primitive recursive ordinal.

In Section 3, we handle the case of **SADS** and **SEM** which does not admit such a direct preservation of well-foundedness. In these cases, one needs to assume $\text{WF}(\omega^{\alpha \times \omega})$ to obtain an ω -extension satisfying $\text{WF}(\alpha)$, where $\alpha \leq \epsilon_0$. We prove in particular that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0) + \Gamma$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$ for $\Gamma \in \{\text{SADS}, \text{SEM}\}$.

In Section 4, we combine the previous conservation theorems to obtain our main theorems. Last, in Section 5, we state some remaining open questions.

2 Conservation over $\text{B}\Sigma_2^0 + \text{WF}(\alpha)$

In this section, we introduce the framework for preserving $\text{WF}(\alpha)$ for an ordinal $\alpha \leq \epsilon_0$ over $\text{RCA}_0 + \text{B}\Sigma_2^0$. We then prove that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha) + \text{WKL}$ and $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha) + \text{COH}$ are both Π_1^1 -conservative extensions of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

The proofs of conservation will all follow the same pattern: Given a countable topped model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$, and an instance $I \in S$ of our problem **P**, we will show the existence of a **P**-solution $G \subseteq M$ of I such that $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$. Then, the conservation result follows from a standard argument that we give for the sake of completeness:

Definition 2.1. A Π_2^1 -problem **P** is a statement of the form $\forall X \Phi(X) \rightarrow \exists Y \Psi(X, Y)$ for some arithmetical formulas Φ and Ψ . A **P**-instance is a set X

such that $\Phi(X)$ holds. A P -solution of a P -instance X is a set Y such that $\Psi(X, Y)$ holds.

In particular, RT_2^2 , EM and ADS are Π_2^1 -problems. For example, an RT_2^2 -instance is a coloring $f : [\mathbb{N}]^2 \rightarrow 2$ and an RT_2^2 -solution to f is an infinite f -homogeneous set.

Theorem 2.2. *Let P be a Π_2^1 problem and $\alpha \leq \epsilon_0$ be an ordinal such that for every countable topped model $\mathcal{M} = (M, S) \models RCA_0 + B\Sigma_2^0 + WF(\alpha)$ and for every P -instance $I \in \mathcal{M}$, there exists a P -solution $G \subset M$ of I such that $\mathcal{M}[G] \models RCA_0 + B\Sigma_2^0 + WF(\alpha)$. Then $RCA_0 + B\Sigma_2^0 + WF(\alpha) + P$ is Π_1^1 -conservative over $RCA_0 + B\Sigma_2^0 + WF(\alpha)$.*

Proof. Assume $RCA_0 + B\Sigma_2^0 + WF(\alpha) \not\models \forall X \phi(X)$ for ϕ an arithmetic formula. Then by the completeness theorem and the downward Löwenheim–Skolem theorem, there is a countable model $\mathcal{M} = (M, S) \models RCA_0 + B\Sigma_2^0 + WF(\alpha) + \neg\phi(A)$ with $A \in S$. We can furthermore assume that \mathcal{M} is topped by A .

From our assumption, we can build a sequence $\mathcal{M} = \mathcal{M}_0 \subseteq \mathcal{M}_1 = (M, S_1) \subseteq \mathcal{M}_2 = (M, S_2) \subseteq \dots$ of countable topped model of $RCA_0 + B\Sigma_2^0 + WF(\alpha)$ such that every P -instance appearing in one of the \mathcal{M}_i will eventually have a solution in one \mathcal{M}_j . Then $\bigcup_{n < \omega} \mathcal{M}_n$ is a model of $RCA_0 + B\Sigma_2^0 + WF(\alpha) + P + \neg\phi(A)$. So $RCA_0 + B\Sigma_2^0 + WF(\alpha) + P \not\models \forall X \phi(X)$. \square

2.1 Conservation over $B\Sigma_2^0$

There exist two main methods for extending a model \mathcal{M} into a model $\mathcal{M}[G]$: by external forcing, and by an internal construction. In the case of conservation theorems over $B\Sigma_2^0 + \neg I\Sigma_2^0$, one arguably needs to use an internal construction. Indeed, using an isomorphism theorem for WKL_0^* , Fiori-Carones et al [11] proved that Π_1^1 -conservation of any Π_2^1 problem over $B\Sigma_2^0 + \neg I\Sigma_2^0$ is equivalent to a formalized version of the first-jump control.

Let $I\Delta_n^0$ be the following induction scheme for every Σ_n^0 formula ϕ and every Π_n^0 formula ψ :

$$\forall x[\phi(x) \leftrightarrow \psi(x)] \wedge \phi(0) \wedge \forall x[\phi(x) \rightarrow \phi(x+1)] \rightarrow \forall x\phi(x)$$

A set X is M -regular (or *piecewise-coded*, or even *amenable*) if all its initial segments are M -finite. Over this article, we will use the following characterizations of $B\Sigma_2^0$:

Proposition 2.3 (Chong and Mourad [4] and Slaman [32]). *The following are equivalent over RCA_0 :*

- (1) $B\Sigma_2^0$
- (2) The induction principle for Δ_2^0 predicates ($I\Delta_2^0$)
- (3) Every Δ_2^0 set is regular

Given a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ for some $\alpha \leq \epsilon_0$ topped by a set $Y \in S$, and a \mathbf{P} -instance $I \in S$ for some Π_2^1 problem \mathbf{P} , we will create a decreasing sequence of conditions $c_0 \geq c_1 \geq \dots$ (for the appropriate notion of forcing depending on \mathbf{P}) such that

- (a) The sequence is $\Delta_1^0(P)$ for some set $P \subseteq M$ such that $\mathcal{M}[P] \models \text{I}\Delta_1^0$
- (b) The set of $s \in M$ such that c_s is defined is a cut
- (c) For every $k \in M$, there is some $s \in M$ such that c_s decides $(G \oplus Y)' \upharpoonright_k$
- (d) For every $k \in M$, there is some $s \in M$ such that c_s forces $\Phi_e^{G \oplus Y}$ not to be an infinite α -decreasing sequence for every $e < k$

The item (a) will ensure that there is a $\Delta_1^0(P)$ formula $\psi(s)$ stating that the construction can be pursued up to stage s . By item (b), $\{s : \psi(s)\}$ is a cut, so by $\text{I}\Delta_1^0(P)$, the construction can be pursued at every stage. By item (c), $(G \oplus Y)'$ will be $\Delta_1^0(P)$, hence $\mathcal{M}[G] \models \text{B}\Sigma_2^0$. Last, by item (d), $\mathcal{M}[G] \models \text{WF}(\alpha)$.

The use of an internal construction prevents from using a bijection between M and ω to satisfy the requirements one by one. Therefore, one must use *Shore blocking* arguments, that is, force the requirements simultaneously for larger and larger initial segments of M , as mentioned in items (c) and (d).

2.2 Well-foundedness and Shore blocking

Before working on Π_1^1 -conservation of WKL and COH, we prove two combinatorial lemma which say that Shore blocking comes for free for preserving $\text{WF}(\alpha)$ for any ordinal $\alpha \leq \epsilon_0$. Instead of using the standard sums and products over ordinals, the natural (Hessenberg) sums and products will be more convenient:

Definition 2.4 (Natural sum and product). *Let α and β be two ordinals less than ϵ_0 . Let $\alpha = \omega^{\gamma_1}n_1 + \dots + \omega^{\gamma_k}n_k$ and $\beta = \omega^{\gamma_1}m_1 + \dots + \omega^{\gamma_k}m_k$ (We allow the n_i and m_i to be equal to 0 in order to write α and β using the same exponents γ_i). We define the natural sum of α and β (denoted by $\alpha \dot{+} \beta$) to be the following ordinal (It does not depend on the choice of the family (γ_i)) :*

$$\omega^{\gamma_1}(n_1 + m_1) + \dots + \omega^{\gamma_k}(n_k + m_k)$$

We also define $\alpha \dot{\times} k$ to be equal to be the natural sum of α with itself k times and $\alpha \dot{\times} \omega = \omega^{\gamma_1+1}n_1 + \dots + \omega^{\gamma_k+1}n_k$ (whereas $\alpha \times \omega = \omega^{\gamma_1+1}n_1$)

Note that $\alpha \dot{\times} k \leq \alpha \times \omega$, where $\alpha \times \omega = \sup_{k < \omega} \alpha \times k = \omega^{\gamma_1+1}n_1$ is the usual ordinal product.

Proposition 2.5. *For all $\beta < \epsilon_0$, and $\alpha' < \alpha < \epsilon_0$: $\alpha' \dot{+} \beta < \alpha \dot{+} \beta$ and $\beta \dot{+} \alpha' < \beta \dot{+} \alpha$.*

Proof. Clear from the definition of $\dot{+}$. □

Lemma 2.6. *Let $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{WF}(\alpha)$ for α an ordinal less than ϵ_0 . Then $\mathcal{M} \models \text{WF}(\alpha \dot{\times} k)$ for every $k \in M$.*

Proof. Since $\alpha \dot{\times} k \leq \alpha \times \omega$ for every $k \in M$ and RCA_0 proves that the product of two well-orders is a well-order, hence $\text{RCA}_0 \vdash \text{WF}(\alpha) \rightarrow \text{WF}(\alpha \times \omega)$, we get the desired result. \square

Lemma 2.7. *Consider a model $\mathcal{M} = (M, S)$ and $\alpha \in M$ an ordinal less than ϵ_0 . For every $k \in M$, there is a Turing functional Γ_k such that, letting $\beta \in \alpha$ be the largest ordinal less than α of index at most k , for every $X \in 2^M$ such that $\mathcal{M}[X] \models \text{RCA}_0$, if there is some $e < k$ such that Φ_e^X is an M -infinite decreasing sequence of elements smaller than β , then Γ_k^X is an M -infinite decreasing sequence of elements smaller than $\beta \dot{\times} k$.*

Moreover, an index of Γ_k can be found computably in k .

Proof. By twisting the Turing functionals, we can assume that for every $e, n \in M$, if $\Phi_e^\sigma(n) \downarrow$, then

- (1) $n < |\sigma|$;
- (2) $\Phi_e^\sigma(m) \downarrow$ for every $m < n$;
- (3) $\Phi_e^\sigma(0), \Phi_e^\sigma(1), \dots, \Phi_e^\sigma(m)$ is a strictly decreasing sequence of elements smaller than β .

Given $\sigma \in 2^{<M}$ and $e < k$, let $\alpha(\sigma, e) = \Phi_e^\sigma(s)$ for the largest $s < |\sigma|$ such that $\Phi_e^\sigma(s) \downarrow$. If there is no such s , then $\alpha(\sigma, e) = \beta$. Note that if $\sigma' \geq \sigma$, then $\alpha(\sigma', e) \leq \alpha(\sigma, e)$.

Let $\sigma_{-1} = \epsilon$. Let Γ_k^X be the Turing functional which, on input n , searches for some $x > |\sigma_{n-1}|$ and some $\sigma_n < X$ such that $\Phi_e^{\sigma_n}(x) \downarrow$ for some $e < k$. If found, it outputs $\alpha(\sigma, 0) + \dots + \alpha(\sigma, k-1)$. Note that if $\Gamma_k^X(n) \downarrow$, then by (3), $\Gamma_k^X(n)$ is an ordinal smaller than $\beta \dot{\times} k$.

Suppose that X is such that $\mathcal{M}[X] \models \text{RCA}_0$ and there is an $e < k$ such that Φ_e^X is total. Since $\mathcal{M}[X] \models \text{IS}_1^0$, then Γ_k^X is total.

Moreover, since $x > |\sigma_{n-1}|$, then letting $e < k$ be such that $\Phi_e^{\sigma_n}(x) \downarrow$, by (1) we have $\Phi_e^{\sigma_{n-1}}(x) \uparrow$, so by (2) and (3), $\alpha(\sigma_{n+1}, e) < \alpha(\sigma_n, e)$, hence $\Gamma_k^X(n+1) < \Gamma_k^X(n)$. It follows that Γ_k^X is an M -infinite decreasing sequence of ordinals smaller than $\beta \dot{\times} k$. \square

2.3 Π_1^1 -conservation of COH

We now turn to the proof that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha) + \text{COH}$ is Π_1^1 -conservative of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ for every ordinal $\alpha \leq \epsilon_0$. In particular, letting $\alpha = \omega$, we reprove the theorem of Chong, Slaman and Yang [5] stating that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{COH}$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$. Note that the original proof by Chong, Slaman and Yang uses a case analysis, depending on whether the model satisfies IS_2^0 or not. Belanger [1] gave a more direct proof using characterization of p-cohesive sets are those whose jump is PA over 0' (see Jockusch and Stephan [17]). Our proof follows the ideas from Belanger, but with an even

more direct construction based on an instance-wise correspondence between instances of COH and $\Pi_1^0(\emptyset')$ classes. As in Belanger's proof, the proof relies on the following theorem:

Theorem 2.8 (Simpson–Smith [31]). *Every countable model of RCA_0^* can be ω -extended into a model of $\text{RCA}_0^* + \text{WKL}$.*

Proposition 2.9. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ topped by a set $Y \in S$, where $\alpha \leq \epsilon_0$ is an ordinal. Then for every $\vec{R} \in S$ and every $P \gg Y'$ such that $\mathcal{M}[P] \models \text{RCA}_0^*$, there exists $G \subseteq M$ such that*

1. G is \vec{R} -cohesive ;
2. $P \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Proof. Fix a $\Delta_2^0(Y)$ tree T such that $[T] = \mathcal{C}(\vec{R})$. In particular, T is $\Delta_1^0(Y')$, so since $P \gg Y'$, there is a $\Delta_1^0(P)$ set $Q \in [T]$.

We will define the set G using an infinite $\Delta_1^0(P)$ decreasing sequence of Mathias-like conditions.

Definition 2.10. *A condition is a pair of M -finite binary strings (σ, ρ) such that $\rho < Q$. We have $(\sigma_2, \rho_2) \leq (\sigma_1, \rho_1)$ if $\sigma_2 > \sigma_1$, $\rho_2 > \rho_1$ and $\sigma_2 - \sigma_1 \subseteq R_{\rho_1}$ (where $\sigma_2 - \sigma_1$ is the difference between the sets corresponding to the two sequences).*

One can think of a condition (σ, ρ) as a Mathias condition (σ, \vec{R}_ρ) . Requiring that $\rho < Q$ ensures that \vec{R}_ρ is M -infinite. When increasing the length of ρ , we make progress in cohesiveness. Note that being a condition is a $\Delta_1^0(Q)$ predicate, as it simply requires checking that $\sigma < Q$, while the extension relation is $\Delta_1^0(Y)$. In particular, both are $\Delta_1^0(Q \oplus Y)$, so $\Delta_1^0(P)$.

A sequence τ is said to be *compatible* with a condition (σ, ρ) if $\sigma < \tau$ and $\tau - \sigma \subseteq R_\rho$.

Definition 2.11. *Let (σ, ρ) be a condition, and Φ_e be a Turing functional.*

1. $(\sigma, \rho) \Vdash \exists x \Phi_e^{G \oplus Y}(x) \downarrow$ if there is some $x, t < |\sigma|$ such that $\Phi_e^{\sigma \oplus Y}(x)[t] \downarrow$
2. $(\sigma, \rho) \Vdash \forall x \Phi_e^{G \oplus Y}(x) \uparrow$ if for every τ compatible with (σ, ρ) and every $x \in M$, $\Phi_e^{\tau \oplus Y}(x) \uparrow$.

Note that the relations $(\sigma, \rho) \Vdash \exists x \Phi_e^{G \oplus Y}(x) \downarrow$ and $(\sigma, \rho) \Vdash \forall x \Phi_e^{G \oplus Y}(x) \uparrow$ are $\Delta_0^0(Y)$ and $\Pi_1^0(Y)$, respectively.

To ensure that $P \geq_T (G \oplus Y)'$, we will also construct $(G \oplus Y)'$ in parallel.

For this, we will need to satisfy three kind of requirements for every $k \in M$:

- \mathcal{R}_k : $(G \oplus Y)' \upharpoonright_k$ is decided
- \mathcal{S}_k : for every $e < k$, Φ_e^G is not an M -infinite α -decreasing sequence

- \mathcal{T}_k : for every $e < k$, $G \subseteq^* R_e$ or $G \subseteq^* \overline{R}_e$

We will need the following two technical lemmas, which are Shore blocking arguments for satisfying the two kind of requirements, namely, deciding the jump and preserving $\text{WF}(\alpha)$. In what follows, $(\tau, \rho) \Vdash (G \oplus Y)' \upharpoonright_k = \sigma'$ means that for every $e < |\sigma'|$, if $\sigma'(e) = 1$ then $(\tau, \rho) \Vdash \Phi_e^{G \oplus Y}(e) \downarrow$ and if $\sigma'(e) = 0$, then $(\tau, \rho) \Vdash \Phi_e^{G \oplus Y}(e) \uparrow$. Thus, this relation is $\Pi_1^0(Y)$.

Lemma 2.12. *Let (σ, ρ) be a condition. For every $k \in M$, there exists some M -finite $\sigma' \in 2^k$ and an extension $(\tau, \rho) \leq (\sigma, \rho)$ such that $(\tau, \rho) \Vdash (G \oplus Y)' \upharpoonright_k = \sigma'$.*

Proof. Let W be the set of all $\sigma' \in 2^k$ such that $(\exists \tau > \sigma)(\exists t)(\forall e < s)(\tau - \sigma \in R_\rho \wedge |\tau - \sigma| \geq 1 \wedge \sigma'(e) = 1 \rightarrow \Phi_e^{\sigma \oplus Y}(e)[t] \downarrow)$. Note that W is $\Sigma_1^0(Y)$, hence is M -finite. Moreover, W is non-empty, as it contains the string $000\dots$. Let $\sigma' \in W$ be the lexicographically maximal element, and let $\tau \geq \sigma$ witness that $\sigma' \in W$.

We claim that (τ, ρ) forces $(G \oplus Y)' \upharpoonright_k = \sigma'$. Fix some $e < k$.

Case (a): $\sigma'(e) = 1$. Then $\Phi_e^{\tau \oplus Y}(e) \downarrow$, hence for every G compatible with (τ, ρ) , $e \in (G \oplus Y)'$.

Case (b): $\sigma'(e) = 0$. The maximality of σ' ensures that for every μ compatible with (τ, ρ) , $\Phi_e^{\mu \oplus Y}(e) \uparrow$. It follows that $\Phi_e^{G \oplus Y}(e) \uparrow$, hence $e \notin (G \oplus Y)'$. \square

Lemma 2.13. *Let (σ, ρ) be a condition. For every $k \in M$, there exists some $n \in M$ an extension $(\tau, \rho) \leq (\sigma, \rho)$ forcing $\Phi_e^{G \oplus Y}$ not to be an α -decreasing sequence of length more than n for every $e < k$.*

Proof. By twisting the Turing functionals, we can assume that for every $e, n \in M$, if $\Phi_e^\sigma(n) \downarrow$, then

- $\Phi_e^\sigma(m) \downarrow$ for every $m < n$
- $\Phi_e^\sigma(0), \Phi_e^\sigma(1), \dots, \Phi_e^\sigma(m)$ is a strictly α -decreasing sequence

Suppose the lemma does not hold. Then in particular, for every M -finite string τ compatible with (σ, ρ) and every $n \in M$, there is some $\tau_1 \geq \tau$ compatible with (σ, ρ) and some $e < k$ such that $\Phi_e^{\tau_1 \oplus Y}(n) \downarrow$.

Build Y -recursively an infinite sequence $\tau_0 \leq \tau_1 \leq \tau_2 \leq \dots$ such that for every n ,

- τ_n is compatible with (σ, ρ)
- $\Phi_e^{\tau_n \oplus Y}(n) \downarrow$ for some $e < k$ (which depends on n)

Since $\mathcal{M} \models \text{BS}_2^0$, then $\mathcal{M} \models \forall k \text{RT}_k^1$, so there is an M -infinite set $Z \in \mathcal{M}$ and some $e < k$ such that for every $n \in Z$, $\Phi_e^{\tau_n \oplus Y}(n) \downarrow$.

Then, letting $f(n) = \Phi_e^{\tau_n \oplus Y}(n)$, we have an M -infinite α -decreasing sequence, contradicting $\mathcal{M} \models \text{WF}(\alpha)$. \square

Note that the statement " (τ, ρ) forces $\Phi_e^{G \oplus Y}$ not to be an α -decreasing sequence of length more than n for every $e < k$ " is $\Pi_1^0(Y)$. Indeed, it means that for every μ compatible with (τ, ρ) and every $e < k$, $\Phi_e^{\mu \oplus Y}(n) \uparrow$."

Construction. We will build a decreasing sequence (σ_s, ρ_s) of conditions and then take for G the union of the σ_s . We will also build an increasing sequence (σ'_s) such that $(G \oplus Y)'$ will be the union of the σ'_s . Initially, we take $\sigma_0 = \sigma'_0 = \rho_0 = \epsilon$, and during the construction, we will ensure that we have $|\sigma_s|, |\sigma'_s|, |\rho_s| \leq s$ at every stage. Each stage will be either of type \mathcal{R} , of type \mathcal{S} or of type \mathcal{T} . The stage 0 is of type \mathcal{R} .

Assume that (σ_s, ρ_s) and σ'_s are already defined. Let $s_0 < s$ be the latest stage at which we switched the stage type. We have three cases.

Case 1: s is of type \mathcal{R} . If there exists some $\tau \in 2^{\leq s}$ and some $\sigma' \in 2^{s_0}$ such that $(\tau, \rho_s) \leq (\sigma_s, \rho_s)$, and (τ, ρ_s) forces $(G \oplus Y)' \upharpoonright_{s_0} = \sigma'$, then let $\sigma_{s+1} = \tau$, $\rho_{s+1} = \rho_s$, $\sigma'_{s+1} = \sigma'$ and let $s+1$ be of type \mathcal{S} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 2: s is of type \mathcal{S} . If there exists some $\tau \in 2^{\leq s}$ such that $(\tau, \rho_s) \leq (\sigma_s, \rho_s)$, and (τ, ρ_s) forces \mathcal{S}_{s_0} , then let $\sigma_{s+1} = \tau$, $\rho_{s+1} = \rho_s$, $\sigma'_{s+1} = \sigma'_s$ and let $s+1$ be of type \mathcal{T} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 3: s is of type \mathcal{T} . If there exists some $\tau \in 2^{\leq s}$ such that $(\tau, \rho_s) \leq (\sigma_s, \rho_s)$ and $\tau - \sigma_s \neq \emptyset$. Then let $\sigma_{s+1} = \tau$, $\rho_{s+1} = P \upharpoonright_{s_0}$, $\sigma'_{s+1} = \sigma'_s$, and let $s+1$ be of type \mathcal{R} . Otherwise, the elements are left unchanged and we go to the next stage.

This completes the construction.

Verification. Since the size of σ_s, σ'_s and ρ_s are bounded by s , there is a $\Delta_1^0(Y' \oplus Q)$ -formula $\phi(s)$ stating that the construction can be pursued up to stage s . Our construction implies that the set $\{s | \phi(s)\}$ is a cut, so by $\text{I}\Delta_1^0(P)$, the construction can be pursued at every stage.

Let $G = \bigcup_{s \in M} \sigma_s$. By Lemma 2.12 and Lemma 2.13 and the fact that each R_{ρ_s} is M -infinite, each type of stage changes M -infinitely often. Thus, G is M -infinite and $\{|\sigma'_s| : s \in M\}$ is M -infinite, so $P \geq_T (G \oplus Y)'$. In particular, $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0$. Moreover, for every $e \in M$, there is a stage $s_0 > e$ at which the stage type switches from \mathcal{R} to \mathcal{S} , and a stage $s > s_0$ at which the stage switches from \mathcal{S} to \mathcal{T} . Then, $(\sigma_{s+1}, \rho_{s+1})$ forces \mathcal{S}_{s_0} , so $\Phi_e^{G \oplus Y}$ is not an M -infinite α -decreasing sequence. Thus, $\mathcal{M}[G] \models \text{WF}(\alpha)$. Similarly, for every $e \in M$, there is a stage $s > e$ such that the stage type switches from \mathcal{S} to \mathcal{T} . Then $(\sigma_{s+1}, \rho_{s+1})$ is such that $|\rho_{s+1}| > e$, so $G \subseteq^* R_e$ or $G \subseteq^* \bar{R}_e$. Thus G is \vec{R} -cohesive. This completes the proof of Proposition 2.9. \square

Corollary 2.14. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ topped by a set $Y \in S$, where $\alpha \leq \epsilon_0$ is an ordinal. Then for every $\vec{R} \in S$, there exists $G \subseteq M$ such that*

1. G is \vec{R} -cohesive ;

2. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Proof. By Theorem 2.8, there exists a set $P \subseteq M$ such that $P \gg Y'$ and $\mathcal{M}[P] \models \text{RCA}_0^*$. The conclusion follows from Proposition 2.9. \square

We therefore obtain our first conservation theorem:

Theorem 2.15. *Fix an ordinal $\alpha \leq \epsilon_0$. $\text{RCA}_0 + \text{COH} + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.*

Proof. Immediate by Proposition 2.9 and Theorem 2.2. \square

2.4 Π_1^1 -conservation of WKL

We now turn to the proof of Π_1^1 -conservation of WKL_0 . As for COH , in the case $\alpha = \omega$, we recover the well-known conservation theorem of Hájek [12] stating that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WKL}$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0$.

Proposition 2.16. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ topped by a set $Y \in S$, where $\alpha \in M$ is an ordinal $\leq \epsilon_0$. Then for every M -infinite primitive Y -recursive tree $T \in S$, there exists $G \subseteq M$ such that*

1. G is a path in T ;
2. $Y' \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$

Proof. We will build the path G using forcing on trees. A condition will be a pair (σ, T') with $T' \in S$ an M -infinite primitive Y -recursive subtree of T with root σ and we will have $(\sigma, T_1) \leq (\tau, T_2)$ if $\tau < \sigma$ and $T_1 \subseteq T_2$.

Definition 2.17. *Let (σ, U) be a condition, and Φ_e be a Turing functional.*

1. $(\sigma, U) \Vdash \exists x \Phi_e^{G \oplus Y}(x) \downarrow$ if there is some $x, t < |\sigma|$ such that $\Phi_e^{\sigma \oplus Y}(x)[t] \downarrow$
2. $(\sigma, U) \Vdash \forall x \Phi_e^{G \oplus Y}(x) \uparrow$ if for every $\tau \in U$ and every $x \in M$, $\Phi_e^{\tau \oplus Y}(x) \uparrow$.

Note that the relations $(\sigma, U) \Vdash \exists x \Phi_e^{G \oplus Y}(x) \downarrow$ and $(\sigma, U) \Vdash \forall x \Phi_e^{G \oplus Y}(x) \uparrow$ are $\Delta_0^0(Y)$ and $\Pi_1^1(Y)$, respectively.

For our construction, we want to satisfy the following requirements :

- \mathcal{R}_k : $(G \oplus Y)' \upharpoonright_k$ is decided.
- \mathcal{S}_k : Let $\beta \in \alpha$ be the $<_{\epsilon_0}$ -biggest element of index less than k , for every $e < k$, $\Phi_e^{G \oplus Y}$ is not an M -infinite $<_{\epsilon_0}$ -decreasing sequence of elements smaller than β .
- \mathcal{T}_k : $|\sigma| \geq k$.

We first prove a simple technical lemma saying that there exists arbitrarily large extendible nodes. This lemma even holds over RCA_0^* and can be found in various places in the literature.

Lemma 2.18 (Fernandes et al [10]). *Let $\mathcal{M} = (M, S) \models \text{RCA}_0^*$ be a model and $T \subseteq 2^{<M}$ be an M -infinite tree in \mathcal{M} . Then for every $\ell \in M$, there is some $\sigma \in T$ of length ℓ such that $\{\tau \in T : \tau \text{ is comparable with } \sigma\}$ is M -infinite.*

Proof. Assume by contradiction that for every $\sigma \in 2^\ell$ the tree

$$\{\tau \in T : \tau \text{ is comparable with } \sigma\}$$

is M -bounded, this means that $\mathcal{M} \models \forall \sigma \in 2^\ell \exists b_\sigma \forall \tau \in 2^{b_\sigma}, \sigma < \tau \Rightarrow \tau \notin T$.

The formula $\forall \tau \in 2^{b_\sigma}, \sigma < \tau \Rightarrow \tau \notin T$ is $\Delta_1^0(T)$ and therefore Δ_1^0 in \mathcal{M} . Therefore, we can use BS_1^0 (which holds in RCA_0^*) to get :

$$\mathcal{M} \models \exists b \forall \sigma \in 2^\ell \exists b_\sigma < b \forall \tau \in 2^{b_\sigma}, \sigma < \tau \Rightarrow \tau \notin T$$

Which yields that T is bounded by such a b , contradicting our assumption that T is M -infinite. So there is some $\sigma \in T$ of length ℓ such that $\{\tau \in T : \tau \text{ is comparable with } \sigma\}$ is M -infinite. \square

We will need the following two technical lemmas, which are Shore blocking arguments for satisfying the two kind of requirements, namely, deciding the jump and preserving $\text{WF}(\alpha)$. In what follows, $(\tau, U) \Vdash (G \oplus Y)' \upharpoonright_k = \sigma'$ means that for every $e < |\sigma'|$, if $\sigma'(e) = 1$ then $(\tau, U) \Vdash \Phi_e^{G \oplus Y}(e) \downarrow$ and if $\sigma'(e) = 0$, then $(\tau, U) \Vdash \Phi_e^{G \oplus Y}(e) \uparrow$. Thus, this relation is $\Pi_1^0(Y)$.

Lemma 2.19. *Let (σ, T_1) a condition. For every $k \in M$, there exists some M -finite σ' and an extension $(\tau, T_2) \leq (\sigma, T_1)$ forcing $(G \oplus Y)' \upharpoonright_k = \sigma'$.*

Proof. Let W be the set of all $\sigma' \in 2^k$ such that the tree $\{\rho \in T_1 \mid \sigma'(e) = 0 \rightarrow \Phi_e^{\rho \oplus Y}(e) \uparrow\}$ is M -infinite. W is $\Pi_1^0(Y)$ and hence M -finite, and it is non-empty as it contains the string 1111... .

We can consider $\sigma' \in W$ its lexicographically minimal element, and consider the corresponding tree T_3 . For every $e < k$ such that $\sigma'(e) = 1$, the minimality of σ' gives us that the set of $\tau \in T_3$ such that $\Phi_e^{\tau \oplus Y}(e) \uparrow$ is M -finite, so we have a level h_e such that for every $\tau \in T_3 \cap 2^{h_e}$, $\Phi_e^{\tau \oplus Y}(e) \downarrow$. The set $\{e < k \mid \sigma'(e) = 1\}$ is M -finite, so we can consider an upperbound h of all the h_e . Finally, by Lemma 2.18, there is a $\tau \in T_3 \cap 2^h$ such that $T_2 := \{\rho \in T_3 : \rho \text{ is comparable with } \tau\}$ is M -infinite (since T_3 is M -infinite).

We claim that (τ, T_2) forces $(G \oplus Y)' \upharpoonright_k = \sigma'$. Fix some $e < k$.

Case (a): $\sigma'(e) = 0$. Then $\Phi_e^{\tau \oplus Y}(e) \downarrow$ for every $\tau \in T_2$ (since $T_2 \subseteq T_3$), hence for every G compatible with (τ, T_2) , so $e \notin (G \oplus Y)'$.

Case (b): $\sigma'(e) = 1$. The definition of τ ensure that $\Phi_e^{\tau \oplus Y}(e) \downarrow$, so $e \in (G \oplus Y)'$. \square

Lemma 2.20. *Let (σ, T_1) be a condition. For every $k \in M$, letting Γ_k be the functional of Lemma 2.7, there is an extension $(\sigma, T_2) \leq (\sigma, T_1)$ forcing $\Gamma_k^{G \oplus Y}$ to be partial.*

Proof. By twisting Γ_k , we can assume that for every $e, n \in M$, if $\Gamma_k^{\sigma \oplus Y}(n) \downarrow$, then

- $\Gamma_k^{\sigma \oplus Y}(m) \downarrow$ for every $m < n$;
- $\Gamma_k^{\sigma \oplus Y}(0), \Gamma_k^{\sigma \oplus Y}(1), \dots, \Gamma_k^{\sigma \oplus Y}(m)$ is a strictly $<_{\epsilon_0}$ -decreasing sequence of ordinals smaller than $\beta \times k$.

We have two cases.

Case 1: there exists some $n \in M$ such that for every $\ell \in M$, there is some $\tau \in T_1$ of length ℓ such that $\Gamma_k^{\sigma \oplus Y}(n) \uparrow$. Let $T_2 = \{\tau \in T_1 : \Gamma_k^{\sigma \oplus Y}(n) \uparrow\}$. Note that the set T_2 is a Y -primitive recursive, as the set T_1 and the predicate $\Gamma_k^{\sigma \oplus Y}(n) \uparrow$ are Y -primitive recursive. The set T_2 is closed-downward, and by assumption, is M -infinite. Then $(\sigma, T_2) \leq (\sigma, T_1)$. Moreover, by construction, $(\sigma, T_2) \Vdash \Gamma_k^{G \oplus Y}(n) \uparrow$.

Case 2: for every $n \in M$, there is some $\ell_n \in M$ such that for every $\sigma \in T$ of length ℓ_n , $\Gamma_k^\sigma(n) \downarrow$.

Then, for every n , let $\alpha_n = \max \{\Gamma_k^\sigma(n) : \sigma \in T_1 \wedge |\sigma| = \ell_n\}$. We claim that for every $n \in M$, $\alpha_{n+1} <_{\epsilon_0} \alpha_n$. Indeed, for every $\sigma \in T_1$ such that $|\sigma| = \ell_{n+1}$, $\Gamma_k^\sigma(n+1) <_{\epsilon_0} \Gamma_k^\sigma \upharpoonright^{\ell_n}$, so

$$\max \{\Gamma_k^\sigma(n+1) : \sigma \in T_1 \wedge |\sigma| = \ell_{n+1}\} <_{\epsilon_0} \max \{\Gamma_k^\sigma(n) : \sigma \in T_1 \wedge |\sigma| = \ell_n\}$$

So $\mathcal{M} \Vdash \text{WF}(\alpha \times k)$. However, by Lemma 2.6, since $\mathcal{M} \models \text{B}\Sigma_2^0 + \text{WF}(\alpha)$, then $\mathcal{M} \models \text{WF}(\alpha \times k)$. Contradiction. \square

Note that the statement “ (σ, U) forces $\Gamma_k^{G \oplus Y}(n) \uparrow$ ” is $\Pi_1^0(Y)$.

Construction We will build a decreasing sequence (σ_s, T_s) of conditions and then take G for the union of the σ_s . We will also build an increasing sequence (σ'_s) such that G' will be the union of the σ'_s . Initially we take $\sigma_0 = \sigma'_0 = \epsilon$ and $T_0 = T$, and during the construction we will ensure that we have $|\sigma_s|, |\sigma'_s| \leq s$ and that the index describing T_s is also less than s at every stage. Each stage will be either of type \mathcal{R} , of type \mathcal{S} or of type \mathcal{T} . The stage 0 is of type \mathcal{R} .

Assume that (σ_s, T_s) and σ'_s are already defined. Let $s_0 < s$ be the latest stage at which we switched the stage type. We have three cases.

Case 1: s is of type \mathcal{R} . If there exists some index $e \leq s$ such that the e -th primitive recursive tree T' is M -infinite and satisfy $(\sigma_s, T') \leq (\sigma_s, T_s)$ and is such that (σ_s, T') forces $(G \oplus Y)' \upharpoonright_k = \sigma'$ for a $\sigma' \in 2^{s_0}$. Then let $\sigma_{s+1} = \sigma_s$, $T_{s+1} = T'$, $\sigma'_{s+1} = \sigma'$ and let $s+1$ be of type \mathcal{S} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 2: s is of type \mathcal{S} . If there exists some $e, n \leq s$ such that the e -th primitive recursive tree T' is M -infinite and (σ_s, T') forces $\Gamma_k^{G \oplus Y}(n) \uparrow$, then let $\sigma_{s+1} = \sigma_s$, $T_{s+1} = T'$, $\sigma'_{s+1} = \sigma'_s$ and let $s+1$ be of type \mathcal{T} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 3: s is of type \mathcal{T} . Then we can find a $\sigma \in 2^s$ such that $\{\rho \in T_s \mid \rho \text{ compatible with } \sigma\}$ is M -infinite (with $\text{B}\Sigma_2^0$). We then take this tree for T_{s+1} and let $\sigma_{s+1} = \sigma$, $\sigma'_{s+1} = \sigma'_s$ and let $s+1$ be of type \mathcal{R} .

This completes the construction.

Verification. Since the size of σ_s, σ'_s and the index of T_s are bounded by s , there is a $\Delta_1^0(Y')$ -formula $\phi(s)$ stating that the construction can be pursued up to stage s . Our construction implies that the set $\{s | \phi(s)\}$ is $\Delta_1^0(Y')$ and forms a cut, so by $\text{I}\Delta_1^0(Y')$, the construction can be pursued at every stage.

Let $G = \bigcup_{s \in M} \sigma_s$. By Lemma 2.19 and Lemma 2.20, each type of stage changes M -infinitely often. Thus, $\{|\sigma'_s| : s \in M\}$ is M -infinite, so $Y' \geq_T (G \oplus Y)'$. In particular, $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0$.

Moreover, the stages s_0 where we force \mathcal{S}_{s_0} are infinite, so for every $\beta \in \alpha$ and $e \in M$, there is a stage that will force $\Phi_e^{G \oplus Y}$ not to be an M -infinite α -decreasing sequence of elements less than β . And thus $\Phi_e^{G \oplus Y}$ will not be an M -infinite α -decreasing sequence for any e , so $\mathcal{M}[G] \models \text{WF}(\alpha)$.

For every $e \in M$, there is a stage $s > e$ such that the stage type switches from \mathcal{S} to \mathcal{T} . Then (σ_{s+1}, T_{s+1}) is such that $|\sigma_{s+1}| = s$, so G is a path in T . This completes the proof of Proposition 2.16. \square

Theorem 2.21. *Fix an ordinal $\alpha \leq \epsilon_0$. $\text{RCA}_0 + \text{WKL} + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.*

Proof. Immediate by Proposition 2.16 and Theorem 2.2. \square

3 Conservation over $\text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$

The cases of SEM and SADS is more complicated than WKL and COH, as $\text{WF}(\omega^{\alpha \times \omega})$ seems to be necessary to preserve $\text{WF}(\alpha)$ where $\alpha \leq \epsilon_0$. We therefore get a fixed-point in the preservation of well-foundedness by letting $\alpha = \epsilon_0$.

3.1 Π_1^1 -conservation of SEM

As mentioned in the introduction, the bounded monotone enumeration principle naturally shows up when formalizing the first-jump control of Ramsey's theorem for pairs. Within the decomposition of RT_2^2 into $\text{COH} + \text{SADS} + \text{SEM}$ of Proposition 1.16, SEM is the only statement not known to be Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$. It is therefore natural to expect to involve BME_* in the proof below. Since BME_* is equivalent to $\text{WF}(\omega^\omega)$ and we suppose $\mathcal{M} \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega^{\alpha \times \omega})$, we will be able to use BME_* .

Proposition 3.1. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega^{\alpha \times \omega})$ topped by a set $Y \in S$, where $\alpha \in M$ is a non-zero ordinal $\leq \epsilon_0$. Then for every infinite stable tournament R in S and every set $P \gg Y'$ such that $\mathcal{M}[P] \models \text{RCA}_0^*$, there exists $G \subseteq M$ such that*

1. G is an M -infinite transitive subtournament ;
2. $P \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Proof. Fix a uniform enumeration of all Y -primitive recursive non-empty tree functionals T_0, T_1, \dots , that is, for every $n \in M$ and every X , T_n^X is an M -infinite tree. Let

$$\mathcal{C} = \left\{ \bigoplus_a X_a : \forall a \forall n \ X_{\langle a, n \rangle} \in [T_n^{Y \oplus X_0 \oplus \dots \oplus X_a}] \right\}$$

The class \mathcal{C} is $\Pi_1^0(Y)$ and non-empty. There exists a primitive Y -recursive tree whose infinite paths are the elements of \mathcal{C} , so Proposition 2.16 gives us a set $\vec{X} = \bigoplus_a X_a \in \mathcal{C}$ such that $Y' \geq_T (\vec{X} \oplus Y)'$ and $\mathcal{M}[\vec{X}] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega^{\alpha \times \omega})$.

Furthermore, Proposition 2.9 applied on $\mathcal{M}[\vec{X}]$ gives us a set $C \subseteq M$ satisfying :

- C is \vec{X} -cohesive ;
- $P \geq_T (C \oplus \vec{X} \oplus Y)'$;
- $\mathcal{M}[\vec{X} \oplus C] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega^{\alpha \times \omega})$

Let $A_0(R) := \{x : \forall^\infty y R(x, y)\}$ and $A_1(R) := \{x : \forall^\infty y R(y, x)\}$. The stability of R gives us $A_0(R) \sqcup A_1(R) = M$, so $A_0(R)$ and $A_1(R)$ are $\Delta_2^0(Y)$.

Definition 3.2. For two sets F and G , the notation $F \rightarrow_R G$ denotes the formula $(\forall x \in F)(\forall y \in G)R(x, y)$.

Definition 3.3 (Minimal interval). Let $a, b \in \mathcal{M}$, the interval (a, b) is the set of all x such that $R(a, x)$ and $R(x, b)$.

Let F be an M -finite transitive subtournament of R , for $a, b \in F \cup \{-\infty, +\infty\}$ we say that (a, b) is a minimal interval of F if $F \cap (a, b) = \emptyset$, where we consider that $R(-\infty, x)$ and $R(x, +\infty)$ holds for every $x \in \mathcal{M}$.

Definition 3.4. A condition is a pair (σ, a) where σ is a M -finite binary string corresponding to an R -transitive subtournament, $a \in \mathcal{M}$ is such that $C \subseteq^* X_a$ and X_a is included in a minimal interval of σ .

Being a condition is a $\Delta_1^0((C \oplus X_a \oplus Y)')$ -formula. Indeed, being a finite R -transitive subtournament is Y computable, checking that X_a is included in a minimal interval of σ is $(X_a \oplus Y)'$ -computable, and checking if $C \subseteq^* X_a$ can be decided using $(C \oplus X_a)'$ (since C is \vec{X} -cohesive, we know that $C \subseteq^* X_a$ or $C \subseteq^* \vec{X}_a$ and these events are $\Sigma_2^0(C \oplus X_a)$). Since $P \geq_T (C \oplus \vec{X} \oplus Y)'$, it is a $\Delta_1^0(P)$ -formula.

Definition 3.5. A condition (τ, b) extends a condition (σ, a) (and we write $(\tau, b) \leq (\sigma, a)$) if $\sigma < \tau$, $X_b \subseteq X_a$ and $\tau - \sigma \subseteq X_a$.

For our construction, we want to satisfy the following requirements :

- \mathcal{R}_k : $(G \oplus Y)' \upharpoonright_k$ is decided.
- \mathcal{S}_k : Let $\beta < \alpha$ be the $<_{\epsilon_0}$ -biggest ordinal of index less than k , for every $e < k$, $\Phi_e^{G \oplus Y}$ is not an M -infinite $<_{\epsilon_0}$ -decreasing sequence of elements smaller than β .

- \mathcal{T}_k : $\text{card } G \geq k$.

We will need two technical lemmas, which are Shore blocking arguments for satisfying the two kinds of requirements, namely, deciding the jump and preserving $\text{WF}(\alpha)$. Before this, we introduce some basic combinatorial concepts which will be useful for both lemmas. A Σ_1^0 formula $\varphi(G)$ is in *normal form* if $\varphi(G) \equiv \exists i \psi(G \upharpoonright i)$ for a Δ_0^0 formula ψ .

Definition 3.6. Fix a condition (σ, a) .

- (1) A (σ, a) -node is an M -finite sequence of the form

$$\nu = \langle (i_0, \rho_0), (i_1, \rho_1), \dots, (i_{q-1}, \rho_{q-1}) \rangle$$

such that for every $s < q$, $i_s < 2$, $\rho_s \subseteq X_a$ is a non-empty M -finite R -transitive set with $\min \rho_s > \max \rho_{s-1}$ (letting $\rho_{-1} = \sigma$), and for every $s < t < q$, if $i_s = 0$ then $\rho_s \rightarrow_R \rho_t$, and if $i_s = 1$ then $\rho_t \rightarrow_R \rho_s$. We then write

$$\pi(\nu) = \sigma \cup \rho_0 \cup \dots \cup \rho_{q-1}$$

- (2) This (σ, a) -node is *valid* if for every $s < q$, $\rho_s \subseteq A_{i_s}(R)$
- (3) A (σ, a) -node ν satisfies a $\Sigma_1^0(\mathcal{M})$ -formula $\varphi(G, x) \equiv \exists i \psi(G \upharpoonright i, x)$ if $\psi(\pi(\nu), |\nu| - 1)$ holds.

Lemma 3.7. Fix a (σ, a) -node ν . Then $\pi(\nu)$ is R -transitive.

Proof. Say $\nu = \langle (i_0, \rho_0), (i_1, \rho_1), \dots, (i_{q-1}, \rho_{q-1}) \rangle$. We prove by Σ_0^0 -induction that for every $\ell < t$, ρ_ℓ is an R -transitive tournament included in a minimal interval of $\sigma \cup \rho_0 \cup \dots \cup \rho_{\ell-1}$ and that $\sigma \cup \rho_0 \cup \dots \cup \rho_\ell$ is R -transitive.

This is true for $\ell = 0$ since (σ, a) is a condition. And assuming the property to be true at a rank $\ell < t - 1$, the construction of $\rho_{\ell+1}$ ensures that it is included in the minimal interval of (σ, a) and that $\rho_{\ell+1} \rightarrow_R \rho_j$ if $i_j = 1$ and $\rho_j \rightarrow_R \rho_{\ell+1}$ if $i_j = 0$, so no elements of $\sigma \cup \rho_0 \cup \dots \cup \rho_\ell$ is between two elements of $\rho_{\ell+1}$. Since $\sigma \cup \rho_0 \cup \dots \cup \rho_\ell$ is R -transitive (by the induction hypothesis), this means that $\rho_{\ell+1}$ is included in one of its minimal intervals, and since it is itself R -transitive, $\sigma \cup \rho_0 \cup \dots \cup \rho_\ell \cup \rho_{\ell+1}$ is R -transitive. \square

Lemma 3.8. Fix a valid (σ, a) -node ν . Then $(\pi(\nu), b)$ is an extension of (σ, a) for some $b \in M$.

Proof. Say $\nu = \langle (i_0, \rho_0), (i_1, \rho_1), \dots, (i_{q-1}, \rho_{q-1}) \rangle$. By definition of validity and stability of R , for every $s < q$, if $i_s = 0$ then $\forall x \in \rho_s \forall^\infty y \in X_a$, $R(x, y)$, and if $i_s = 1$ then $\forall x \in \rho_s \forall^\infty y \in X_a$, $R(y, x)$. By $\text{B}\Sigma_2^0$, there exists some $r \in M$ such that for every $s < q$, if $i_s = 0$ then $\rho_s \rightarrow_R X_a \setminus \{0, \dots, r\}$, and if $i_\ell = 1$ then $X_a \setminus \{0, \dots, r\} \rightarrow_R \rho_\ell$. Let $b \in M$ be such that $X_b = X_a \setminus \{0, \dots, r\}$ and let $\tau = \sigma \cup \rho_0 \cup \dots \cup \rho_{q-1}$. By Lemma 3.7, τ is R -transitive and by construction, $X_b \subseteq X_a$ is included in a minimal interval of τ . Thus (τ, b) is an extension of (σ, a) . \square

Definition 3.9. Fix a condition (σ, a) .

- (1) A (σ, a) -tree is a set of (σ, a) -nodes closed downward under the prefix relation.
- (2) A (σ, a) -tree E is universal if for every (σ, a) -node ν which is not a leaf, there is some $r \in M$ such that for every M -finite 2-partition $A_0 \sqcup A_1 = \{0, \dots, r-1\}$, there is a side $i < 2$ and an M -finite R -transitive $\rho \subseteq A_i \cap X_a$ such that $\nu \cdot \langle i, \rho \rangle \in E$.
- (3) A (σ, a) -tree E satisfies a $\Sigma_1^0(\mathcal{M})$ formula if all its nodes satisfies it.

Lemma 3.10. For every $\Sigma_1^0(\mathcal{M})$ formula $\varphi(G, x)$ and every condition (σ, a) , there is a monotone enumeration of a maximal (for inclusion) universal (σ, a) -tree E satisfying $\varphi(G, x)$.

Proof. At stage 0 enumerate the empty sequence $\langle \rangle$. At stage s , for every leaf $\nu \in E$, if for every 2-partition $A_0 \sqcup A_1 = \{0, \dots, s\}$, there is some $i < 2$ and some $\rho \subseteq A_i$ for which

- (i) $\varphi(\pi(\nu) \cup \rho, |\nu|)$ holds
- (ii) $\nu \cdot \langle (i, \rho) \rangle$ is a (σ, a) -node.

then enumerate $\nu \cdot \langle (i, \rho) \rangle$ for all such partitions, and go to the next stage. If nothing is enumerated, simply go to the next stage. \square

A maximal universal (σ, a) -tree is not necessarily M -finite.

Lemma 3.11. Every M -finite universal (σ, a) -tree E has a valid leaf.

Proof. We prove it by $\text{I}\Delta_2^0$ induction. The root $\langle \rangle$ is vacuously valid. Assume $\nu \in E$ is a valid non-leaf. By universality of E , there is some $r \in M$ such that for every M -finite 2-partition $A_0 \sqcup A_1 = \{0, \dots, r-1\}$, there is a side $i < 2$ and an M -finite R -transitive $\rho \subseteq A_i \cap X_a$ such that $\nu \cdot \langle i, \rho \rangle \in E$. By $\text{BC}\Delta_2^0(Y)$ (which is equivalent to $\text{B}\Sigma_2^0(Y)$), the following partition is M -finite:

$$A_0(R) \upharpoonright_r \sqcup A_1(R) \upharpoonright_r = \{0, \dots, r-1\}$$

Thus, there exists some $i < 2$ and some $\rho \subseteq A_i(R)$ such that $\nu \cdot \langle i, \rho \rangle \in E$. By choice of i and ρ , the node $\nu \cdot \langle i, \rho \rangle$ is valid. \square

Lemma 3.12. Let (σ, a) be a condition, $\varphi(G, x)$ be a $\Sigma_1^0(\mathcal{M})$ formula, and E be an M -finite maximal universal (σ, a) -tree satisfying $\varphi(G, x)$. Then there is an extension $(\tau, c) \leq (\sigma, a)$ and some $q \in M$ such that $(\tau, c) \Vdash \varphi(G, q-1)$ and $(\tau, c) \Vdash \neg \varphi(G, q)$.

Proof. By Lemma 3.11, there is a valid leaf $\nu \in E$. Since E satisfies $\varphi(G, x)$, then $\varphi(\pi(\nu), |\nu| - 1)$ holds. By Lemma 3.8, there is some $b \in M$ such that $(\pi(\nu), b) \leq (\sigma, a)$.

By maximality of the tree, for every $s \in M$, there exists a 2-partition $A_0 \sqcup A_1 = \{0, \dots, s\}$, such that for every side $i < 2$ and every M -finite $\rho \subseteq A_i \cap X_a$ such that $\nu \cdot \langle i, \rho \rangle$ is a (σ, a) -node, then $\varphi(\pi(\nu) \cup \rho, |\nu|)$ does not hold.

Note that by choice of X_b , for every M -finite R -transitive $\rho \subseteq A_i \cap X_b$, $\nu \cdot \langle i, \rho \rangle$ is a (σ, a) -node. Thus, for every $s \in M$, there exists a 2-partition $A_0 \sqcup A_1 = \{0, \dots, s\}$, such that (P) for every side $i < 2$ and every M -finite R -transitive $\rho \subseteq A_i \cap X_b$, $\varphi(\pi(\nu) \cup \rho, |\nu|)$ does not hold.

By compactness, there exists a 2-partition $A_0 \sqcup A_1 = X_b$ such that (P) holds. Moreover, the class \mathcal{C} of all such partitions is $\Pi_1^0(Y \oplus X_b)$ and non-empty. By definition of \vec{X} , there exists some $c, d \in M$ such that X_c, X_d is a 2-partition of X_b in \mathcal{C} . Moreover, since $G \subseteq^* X_b$, then either $G \subseteq^* X_c$ or $G \subseteq^* X_d$. Since they play a symmetric role, say $G \subseteq^* X_c$. Then (τ, c) is an extension of (τ, b) forcing $\varphi(G, |\nu| - 1)$ and $\neg\varphi(G, |\nu|)$. \square

We are now ready to prove our two Shore blocking lemmas.

Lemma 3.13. *Let (σ, a) be a condition. For every $k \in M$, there exists some M -finite $\sigma' \in 2^k$ and an extension $(\tau, b) \leq (\sigma, a)$ such that $(\tau, b) \Vdash (G \oplus Y)' \upharpoonright k = \sigma'$*

Proof. Let $\varphi(G, x)$ be the $\Sigma_1^0(Y)$ formula which holds if there exists some $e_0 < \dots < e_{x-1} < k$ such that for every $s < x$, $\Phi_{e_s}^{G \oplus Y}(e_s) \downarrow$.

By Lemma 3.10, there exists a monotone enumeration of a maximal universal (σ, a) -tree E satisfying $\varphi(G, x)$. Moreover, since $\varphi(G, k+1)$ never holds, then E is k -bounded. Since $\alpha > 0$, then $\mathcal{M} \models \text{WF}(\omega^\omega)$, hence $\mathcal{M} \models \text{BME}_*$, so E is M -finite. By Lemma 3.12, there is an extension $(\tau, c) \leq (\sigma, a)$ and some $q \in M$ such that $(\tau, c) \Vdash \varphi(G, q-1)$ and $(\tau, c) \Vdash \neg\varphi(G, q)$. Let $\sigma' = \{e < k : \Phi_e^{\tau \oplus Y}(e) \downarrow\}$. Then $(\tau, b) \Vdash (G \oplus Y)' \upharpoonright k = \sigma'$. \square

Lemma 3.14. *Let (σ, a) be a condition. For every $k \in M$, letting Γ_k be the functional of Lemma 2.7 for the order α , there is an extension $(\tau, b) \leq (\sigma, a)$ forcing $\Gamma_k^{G \oplus Y}$ to be partial.*

Proof. By twisting Γ_k , we can assume that for every $e, n \in M$, if $\Gamma_k^{\sigma \oplus Y}(n) \downarrow$, then

- $\Gamma_k^{\sigma \oplus Y}(m) \downarrow$ for every $m < n$;
- $\Gamma_k^{\sigma \oplus Y}(0), \Gamma_k^{\sigma \oplus Y}(1), \dots, \Gamma_k^{\sigma \oplus Y}(m)$ is a strictly $<_{e_0}$ -decreasing sequence of elements smaller than $\beta \times k$.

Let $\varphi(G, x)$ be the $\Sigma_1^0(Y)$ formula which holds if $\Gamma_k^{G \oplus Y}(x) \downarrow$. By Lemma 3.10, there exists a monotone enumeration of a maximal universal (σ, a) -tree E satisfying $\varphi(G, x)$. We have two cases.

Case 1: E is M -finite. Then by Lemma 3.12, there is an extension $(\tau, c) \leq (\sigma, a)$ and some $q \in M$ such that $(\tau, c) \Vdash \varphi(G, q-1)$ and $(\tau, c) \Vdash \neg\varphi(G, q)$. In particular, $(\tau, c) \Vdash \Gamma_k^{G \oplus Y}(x) \uparrow$.

Case 2: E is not M -finite. Then there exists M -infinitely many stages $s \in M$ such that $E[s] \neq E[s+1]$. By speeding up the enumeration, we can assume without loss of generality that $E[s+1] \neq E[s] \neq \{\langle \rangle\}$ for every $s \in M$.

For every $s \in M$ and $\nu \in E[s]$, let

$$\zeta_s(\nu) = \begin{cases} \omega^{\Gamma_k^{\pi(\nu) \oplus Y}(|\nu|-1)} & \text{if } \nu \text{ is a leaf of } E[s] \\ \sum_{\mu \sqsupset_{E[s]} \nu} \zeta_s(\mu) & \text{otherwise} \end{cases}$$

Finally, let $\zeta_s = \zeta_s(\langle \rangle)$.

Claim 1: $\zeta_{s+1} <_{\epsilon_0} \zeta_s$ for all $s \in M$. Since $E[s+1] \neq E[s]$, there is some leaf $\nu \in E[s]$ which is not a leaf in $E[s+1]$. By assumption on Γ_k , for every $\mu \sqsupset_{E[s+1]} \nu$,

$$\Gamma_k^{\pi(\mu) \oplus Y}(|\mu|-1) <_{\epsilon_0} \Gamma_k^{\pi(\nu) \oplus Y}(|\nu|-1)$$

Hence

$$\sum_{\mu \sqsupset_{E[s+1]} \nu} \omega^{\Gamma_k^{\pi(\mu) \oplus Y}(|\mu|-1)} <_{\epsilon_0} \omega^{\Gamma_k^{\pi(\nu) \oplus Y}(|\nu|-1)}$$

So $\zeta_{s+1}(\nu) <_{\epsilon_0} \zeta_s(\nu)$. By Δ_0^0 induction along the ancestors of ν up to the root, we obtain $\zeta_{s+1}(\langle \rangle) <_{\epsilon_0} \zeta_s(\langle \rangle)$. This concludes our proof of Claim 1.

Claim 2: $\zeta_s <_{\epsilon_0} \omega^{\alpha \dot{\times} \omega}$ for all $s \in M$. For all $s \in M$, $\zeta_s \leq_{\epsilon_0} \zeta_0 \leq_{\epsilon_0} \omega^{\beta \dot{\times} k} <_{\epsilon_0} \omega^{\alpha \dot{\times} \omega}$.

Then Claim 1 and Claim 2 together contradict the fact that $\mathcal{M} \models \text{WF}(\omega^{\alpha \dot{\times} k})$. \square

Construction We will build a decreasing sequence (σ_s, a_s) of conditions and G to be the union of the σ_s . We will also build an increasing sequence (σ'_s) such that G' will be the union of the σ'_s . Initially we take $\sigma_0 = \sigma'_0 = \epsilon$ and a_0 an index corresponding to M , and during the construction we will ensure that we have $|\sigma_s|, |\sigma'_s| \leq s$ and that the index a_s is also less than s at every stage. Each stage will be either of type \mathcal{R} , of type \mathcal{S} or of type \mathcal{T} . The stage 0 is of type \mathcal{R} .

Assume that (σ_s, a_s) and σ'_s are already defined. Let $s_0 < s$ be the latest stage at which we switched the stage type. We have three cases.

Case 1: s is of type \mathcal{R} . If there exists some condition $(\sigma, a) \leq (\sigma_s, a_s)$ with $|\sigma| \leq s$ and $a \leq s$ such that $(\sigma, a) \Vdash (G \oplus Y)' \upharpoonright_{s_0} = \sigma'$ for a $\sigma' \in 2^{s_0}$. Then let $\sigma_{s+1} = \sigma$, $a_{s+1} = a$, $\sigma'_{s+1} = \sigma'$ and let $s+1$ be of type \mathcal{S} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 2: s is of type \mathcal{S} . If there exists some condition $(\sigma, a) \leq (\sigma_s, a_s)$ with $|\sigma| \leq s$ and $a \leq s$ such that (σ, a) forces $\Gamma_{s_0}^{G \oplus Y}(m) \uparrow$ for some $m \leq s$, then let $\sigma_{s+1} = \sigma$, $a_{s+1} = a$, $\sigma'_{s+1} = \sigma'_s$ and let $s+1$ be of type \mathcal{T} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 3: s is of type \mathcal{T} . If there is some condition $(\sigma, a) \leq (\sigma_s, a_s)$ with $\sigma - \sigma_s$ non-empty, $|\sigma| \leq s$ and $a \leq s$, then we let $\sigma_{s+1} = \sigma$, $a_{s+1} = a$ and $\sigma'_{s+1} = \sigma'_s$. Otherwise, the elements are left unchanged and we go to the next stage.

This completes the construction.

Verification. Since the size of σ_s, σ'_s and the index a_s are bounded by s , there is a $\Delta_1^0(P \oplus Y')$ -formula $\phi(s)$ stating that the construction can be pursued up to stage s . Our construction implies that the set $\{s | \phi(s)\}$ is $\Delta_1^0(P)$ and a cut, so by $\text{I}\Delta_1^0(P)$, the construction can be pursued at every stage.

Let $G = \bigcup_{s \in M} \sigma_s$. By Lemma 3.13, Lemma 3.14, every time we enter a stage of type \mathcal{R} or \mathcal{S} we eventually leave it. Also, for every condition (σ_s, a_s) , we can add an element of X_{a_s} to σ_s and then extract an infinite minimal interval from X_{a_s} using P , so every time we enter a stage of type \mathcal{T} , we eventually leave it. So, each type of stage changes M -infinitely often. Thus, $\{|\sigma'_s| : s \in M\}$ is M -infinite, so $P \geq_T (G \oplus Y)'$. In particular, $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0$.

Moreover, the stages s_0 where we force \mathcal{S}_{s_0} are infinite, so for every $\beta \in \alpha$ and $e \in M$, there is a stage that will force $\Phi_e^{G \oplus Y}$ not to be an M -infinite ϵ_0 -decreasing sequence of elements less than β . And thus $\Phi_e^{G \oplus Y}$ will not be an M -infinite α -decreasing sequence for any e , so $\mathcal{M}[G] \models \text{WF}(\alpha)$.

The stages of type \mathcal{T} happens M -infinitely often, so the final set G will be M -infinite. This completes the proof of Proposition 3.1. \square

Corollary 3.15. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega^{\alpha \times \omega})$ topped by a set $Y \in S$, where $\alpha \in M$ is an ordinal $\leq \epsilon_0$. Then for every infinite stable tournament R in S , there exists $G \subseteq M$ such that*

1. G is an M -infinite transitive subtournament ;
2. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Proof. By Theorem 2.8, there exists a set $P \subseteq M$ such that $P \gg Y'$ and $\mathcal{M}[P] \models \text{RCA}_0^*$. The conclusion follows from Proposition 3.1. \square

Theorem 3.16. $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0) + \text{SEM}$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$.

Proof. Immediate by Corollary 3.15 and Theorem 2.2. \square

3.2 Π_1^1 -conservation of SADS

We now turn to the proof that $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0) + \text{SADS}$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$. There exists two main constructions of solutions to instances of SADS: an asymmetric one, which assumes that there is no infinite computable descending sequence, and constructs an ascending one, as in Hirschfeldt and Shore [15, Theorem 2.11], and a symmetric one using split pairs, as in Lerman, Solomon and Towsner [23] or Patey [27, Theorem 26]. Our proof follows the second construction.

Proposition 3.17. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \forall k \text{WF}(\omega^{\alpha \times k})$ topped by a set $Y \in S$, where $\alpha \in M$ is an ordinal $\leq \epsilon_0$. Then for every linear order $L = (M, \leq_L) \in S$ of order type $\omega + \omega^*$, there exists $G \subseteq M$ such that*

1. G is an M -infinite ascending or descending sequence for \leq_L ;
2. $Y' \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Proof. Let $L = (M, \leq_L)$ be such an order, let $U \subseteq M$ and $V \subseteq M$ be the sets of elements that have a finite amount of predecessors and successors, respectively. By assumption, $U \sqcup V = M$, and U and V are both $\Delta_2^0(Y)$.

A condition is a pair (σ, τ) with σ an M -finite L -ascending sequence included in U and τ an M -finite L -descending sequence included in V . We have $(\sigma_2, \tau_2) \leq (\sigma_1, \tau_1)$ if $\sigma_2 \geq \sigma_1$ and $\tau_2 \geq \tau_1$.

Being a condition is a $\Delta_2^0(Y)$ -predicate, while the extension relation is Δ_1^0 .

Definition 3.18. Let (σ_0, σ_1) be a condition, $e, x \in M$ and $i < 2$, we write:

- $(\sigma_0, \sigma_1) \Vdash_i \Phi_e^{G \oplus Y}(x) \downarrow$ if $\Phi_e^{\sigma_i \oplus Y}(x) \downarrow$.
- $(\sigma_0, \sigma_1) \Vdash_i \Phi_e^{G \oplus Y}(x) \uparrow$ if $\forall \sigma'_i \geq \sigma_i$ such that $(\sigma'_i, \sigma_{1-i})$ is a split pair, $\Phi_e^{\sigma'_i \oplus Y}(x) \uparrow$.

We want our construction to satisfy for some $i < 2$ the following requirements for every $k \in M$:

- \mathcal{R}_k^i : $(G_i \oplus Y)' \upharpoonright_k$ is decided.
- \mathcal{S}_k^i : Let $\beta < \alpha$ be the $<_{\epsilon_0}$ -biggest ordinal of index less than k , for every $e < k$, $\Phi_e^{G_i \oplus Y}$ is not an M -infinite $<_{\epsilon_0}$ -decreasing sequence of elements smaller than β .
- \mathcal{T}_k^i : $\text{card } G_i \geq k$.

For this, we will use a pairing argument, and ensure that for every $k \in M$, the following requirements are met: $\mathcal{R}_k^0 \vee \mathcal{R}_k^1$, $\mathcal{R}_k^0 \vee \mathcal{S}_k^1$, $\mathcal{S}_k^0 \vee \mathcal{R}_k^1$, $\mathcal{S}_k^0 \vee \mathcal{S}_k^1$, and $\mathcal{T}_k^0 \wedge \mathcal{T}_k^1$.

Definition 3.19. Fix a condition (σ, τ) .

A (σ, τ) -tree is a binary tree T labelled by a family of split pairs $(\sigma_\rho, \tau_\rho)_{\rho \in T}$ such that for all $\rho \in T$:

- $\sigma \leq \sigma_\rho$ and $\tau \leq \tau_\rho$;
- For every $\mu \in T$, if $\rho \cdot 0 \leq \mu$ then $\sigma_\rho \leq \sigma_\mu$ and if $\rho \cdot 1 \leq \mu$ then $\tau_\rho \leq \tau_\mu$;

A node ρ is said to be valid if the biggest index i such that $\rho(i) = 0$ satisfies $\sigma_\rho \upharpoonright_i \subseteq U$ (if no such index exists this condition is verified) and if the biggest index j such that $\rho(i) = 1$ satisfies $\tau_\rho \upharpoonright_j \subseteq V$ (if no such index exists this condition is verified). Being valid is a $\Delta_2^0(Y)$ condition.

Definition 3.20. Let T a (σ, τ) -tree

- A node (σ_ρ, τ_ρ) in T satisfies a pair of $\Sigma_1^0(Y)$ -formulas $(\varphi_0(G_0, x), \varphi_1(G_1, x))$ if $\varphi_0(\sigma, |\rho| - \text{card } \rho - 1)$ and $\varphi_1(\tau, \text{card } \rho - 1)$ holds (where $\text{card } \rho$ is the cardinal of ρ seen as a set).
- T satisfies $(\varphi_0(G_0, x), \varphi_1(G_1, x))$ if every node of T satisfies it.

Lemma 3.21. For every pair $(\varphi_0(G_0, x), \varphi_1(G_1, x))$ of $\Sigma_1^0(Y)$ formulas and every condition (σ, τ) , there is a maximal Σ_1^0 (σ, τ) -tree T satisfying it.

Proof. T will be the limit of the following increasing sequence of computable (σ, τ) -trees $(T_s)_{s \in M}$ that all satisfy $(\varphi_0(G_0, x), \varphi_1(G_1, x))$ (and therefore T will also be a (σ, τ) -tree satisfying $(\varphi_0(G_0, x), \varphi_1(G_1, x))$).

At stage 0, let T_0 be the empty tree. It vacuously satisfies $(\varphi_0(G_0, x), \varphi_1(G_1, x))$.

At stage $s + 1$, if there exists a split pair (σ', τ') with $|\sigma'|, |\tau'| \leq s$, a $\rho \in T_s$ and an $i < 2$ with $\rho \cdot i \notin T_s$ such that adding $\rho \cdot i$ to T_s with the index (σ', τ') still yields a (σ, τ) -tree satisfying $(\varphi_0(G_0, x), \varphi_1(G_1, x))$, then let T_{s+1} be that tree (choose the smallest split pair possible to enforce the maximality of T). If no such element is found, simply go to the next stage. \square

Lemma 3.22. Let (σ, τ) be a condition, $(\varphi_0(G_0, x), \varphi_1(G_1, x))$ be a pair of $\Sigma_1^0(Y)$ formulas, and T be an M -finite maximal (σ, τ) -tree satisfying $(\varphi_0(G_0, x), \varphi_1(G_1, x))$. Then there is an extension $(\hat{\sigma}, \hat{\tau}) \leq (\sigma, \tau)$ and some $p, q \in M$ such that the following holds:

- $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \varphi_0(G_0, p - 1)$ and $(\hat{\sigma}, \hat{\tau}) \Vdash_1 \varphi_1(G_1, q - 1)$
- $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \neg \varphi_0(G_1, p)$ or $(\hat{\sigma}, \hat{\tau}) \Vdash_1 \neg \varphi_1(G_1, q)$.

Proof. Consider a valid $\rho \in T$ having no valid descendant in T , this is possible since the set of valid element of T is non empty because $\pi(\epsilon) = (\sigma, \tau)$ is a condition, so by can just take a maximal valid element of T using $\mathbf{L}\Delta_2^0$ (which we have by $\mathbf{B}\Sigma_2^0$).

We have two cases:

Case 1: $\sigma_\rho \subseteq U$. In that case $\rho \cdot 0 \notin T$ by maximality of ρ . Then take $p = |\rho| - \text{card } \rho$, $q = \text{card } \rho$ and $(\hat{\sigma}, \hat{\tau}) = (\sigma_\rho, \tau_{\rho \upharpoonright i_0})$ with i_0 the biggest index $i < |\rho|$ such that $\rho(i) = 1$ (if no such index exists we take τ instead of $\tau_{\rho \upharpoonright i_0}$), the pair $(\hat{\sigma}, \hat{\tau})$ is a condition by definition of validity and the hypothesis $\sigma_\rho \subseteq U$. $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \varphi_0(G_0, p - 1)$ and $(\hat{\sigma}, \hat{\tau}) \Vdash_1 \varphi_1(G_1, q - 1)$ because T satisfies $(\varphi_0(G_0, x), \varphi_1(G_1, x))$. And $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \neg \varphi_0(G_0, p)$ by maximality of T (Otherwise we could add $\rho \cdot 0$ to T indexed by a split pair contradicting $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \neg \varphi_0(G_0, p)$).

Case 2: The case $\tau_\rho \subseteq V$ can be treated similarly. \square

Lemma 3.23. Let $k \in M$, let $\Phi_k(G, x)$ be the formula: $\exists e_0 < e_1 < \dots < e_{x-1} < k$ such that $\Phi_{e_i}^{G \oplus Y}(e_i) \downarrow$ for all $i < x$, let $\Psi_k(G, x)$ be the formula: $\Gamma_k^{G \oplus Y}(x) \downarrow$ (with Γ_k the functional of Lemma 2.7 for the order α).

For any combination $\phi_0(G, x), \phi_1(G, x) \in \{\Phi_k(G, x), \Psi_k(G, x)\}$ and every condition (σ, τ) , there exists an extension $(\hat{\sigma}, \hat{\tau}) \leq (\sigma, \tau)$ and some $p, q \in M$ such that the following holds:

- $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \varphi_0(G_0, p - 1)$ and $(\hat{\sigma}, \hat{\tau}) \Vdash_1 \varphi_1(G_1, q - 1)$
- $(\hat{\sigma}, \hat{\tau}) \Vdash_0 \neg \varphi_0(G_1, p)$ or $(\hat{\sigma}, \hat{\tau}) \Vdash_1 \neg \varphi_1(G_1, q)$.

Proof. By Lemma 3.21 there exists a maximal $\Sigma_1^0(\sigma, \tau)$ -tree T satisfying

$$(\varphi_0(G, x), \varphi_1(G, x))$$

. To apply Lemma 3.22 and conclude, we just need to prove the M -finiteness of T .

By twisting Γ_k , we can assume that for every $e, n \in M$, if $\Gamma_k^{\sigma \oplus Y}(n) \downarrow$, then

- $\Gamma_k^{\sigma \oplus Y}(m) \downarrow$ for every $m < n$;
- $\Gamma_k^{\sigma \oplus Y}(0), \Gamma_k^{\sigma \oplus Y}(1), \dots, \Gamma_k^{\sigma \oplus Y}(m)$ is a strictly $<_{\epsilon_0}$ -decreasing sequence of elements smaller than $\beta \times k$.

Assume by contradiction that T is M -infinite and let $(T_s)_{s \in M}$ be an increasing sequence of computable tree approaching T . Without loss of generality we can assume that $T_s \neq T_{s+1}$ for all $s \in M$.

For every $s \in M$ and ρ node of T_s indexed by (σ_ρ, τ_ρ) with set of immediate successors S_ρ ($\text{card } S_\rho \leq 2$ since T_s is a binary tree), define inductively :

$$\zeta_s(\rho) = \omega^{f_0(\rho) + f_1(\rho)} (2 - \text{card } S_\rho) \dot{+} \sum_{\mu \in S_\rho} \zeta_s(\mu)$$

Where $f_0(\rho) = \Gamma_k^{\sigma_\rho \oplus Y}(|\rho| - \text{card } \rho)$ if $\varphi_0(G, x) = \Psi_k(G, x)$ and $f_0(\rho) = k - (|\rho| - \text{card } \rho)$ if $\varphi_0(G, x) = \Phi_k(G, x)$. And $f_1(\rho) = \Gamma_k^{\tau_\rho \oplus Y}(\text{card } \rho)$ if $\varphi_1(G, x) = \Psi_k(G, x)$ and $f_1(\rho) = k - \text{card } \rho$ if $\varphi_1(G, x) = \Phi_k(G, x)$.

And let $\zeta_s = \zeta_s(\epsilon)$.

Claim. The sequence ζ_s is an M -infinitely $<_{\epsilon_0}$ -decreasing sequence of ordinals less than ω^α

Let $s \in M$, since $T_{s+1} \neq T_s$, there is a $\rho \in T_s$ an $i < 2$ such that $T_{s+1} = T_s \cup \rho \cdot i$. Therefore:

$$\zeta_{s+1}(\rho) = \zeta_s(\rho) \dot{-} \omega^{f_0(\rho) + f_1(\rho)} \dot{+} \omega^{f_0(\rho \cdot i) + f_1(\rho \cdot i)} \times 2$$

But for every possible values of φ_0 and φ_1 , we have $f_0(\rho) > f_0(\rho \cdot 0)$, $f_0(\rho) = f_0(\rho \cdot 1)$, $f_1(\rho) = f_1(\rho \cdot 0)$ and $f_1(\rho) > f_1(\rho \cdot 1)$. So for every possible value of i , we have

$$\omega^{f_0(\rho) + f_1(\rho)} > \omega^{f_0(\rho \cdot i) + f_1(\rho \cdot i)} \times 2$$

and therefore $\zeta_{s+1}(\rho) < \zeta_s(\rho)$ which gives us $\zeta_{s+1} < \zeta_s$ (as $\rho \cdot i$ is the only difference between T_s and T_{s+1}).

Therefore, for all $s \in M$, $\zeta_s \leq_{\epsilon_0} \zeta_0 \leq_{\epsilon_0} \omega^{\beta \dot{\times} k} < \omega^{\alpha \dot{\times} \omega}$.

This contradicts $\text{WF}(\omega^{\alpha \dot{\times} \omega})$, so T is M -finite and the result follows. \square

Remark 3.24. If $(\sigma, \tau) \Vdash_i \Phi_k(G_i, p-1)$ and $(\sigma, \tau) \Vdash_i \neg \Phi_k(G_i, p)$, then there exists a $\rho' \in 2^k$ such that $(\sigma, \tau) \Vdash_i (G_i \oplus Y)' \upharpoonright k = \rho'$. Such a ρ' can be found computably in Y .

Moreover, if $(\sigma, \tau) \Vdash_i \Psi_k(G_i, p-1)$ and $(\sigma, \tau) \Vdash_i \neg \Psi_k(G_i, p)$ then $(\sigma, \tau) \Vdash_i \Gamma_{s_0}^{G_i \oplus Y}(p) \uparrow$, so (σ, τ) forces $\Gamma_{s_0}^{G_i \oplus Y}$ to be partial.

Remark 3.25. Because of the disjunctive requirements, preservation of $\text{WF}(\epsilon_0)$ and first jump control must be combined. Thus, at first sight, $\text{WF}(\epsilon_0)$ is used to control the first jump. However, if one wants to control only the first jump, no well-foundedness assumption is necessary. Indeed, let U be the set of pairs $(p, q) \in k^2$ such that there is a pair (σ_1, τ_1) with $\sigma_1 \geq \sigma$, $\tau_1 \geq \tau$ and $\max_{\mathcal{L}} \sigma_1 < \min_{\mathcal{L}} \tau_1$ satisfying $\Psi_k(\sigma_1, p)$ and $\Psi_k(\tau_1, q)$. Then U exists by bounded Σ_1^0 comprehension. Using this set U , one can find an extension with the desired property. Therefore, one can prove that $\text{RCA}_0 + \text{BS}_2^0 + \text{SADS}$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{BS}_2^0$, as already proved by Chong, Slaman and Yang [5] using the first construction.

Construction We will build a decreasing sequence (σ_s, τ_s) of conditions and let G_0 and G_1 to be the union of the σ_s and τ_s . We will also build two increasing sequence (σ'_s) and (τ'_s) such that G'_0 and G'_1 will be the union of the σ'_s and τ'_s . Initially we take $\sigma_0 = \sigma'_0 = \tau_s = \tau'_s = \epsilon$, and during the construction we will ensure that we have $|\sigma_s|, |\sigma'_s|, |\tau_s|, |\tau'_s| \leq s$.

Each stage of the construction will be of one of the following 5 types corresponding to the requirements: $\mathcal{R} \vee \mathcal{R}$, $\mathcal{S} \vee \mathcal{R}$, $\mathcal{R} \vee \mathcal{S}$, $\mathcal{S} \vee \mathcal{S}$ or of type $\mathcal{T} \wedge \mathcal{T}$. The stage 0 is of type $\mathcal{R} \vee \mathcal{R}$.

Assume that (σ_s, τ_s) , σ'_s and τ'_s are already defined. Let $s_0 < s$ be the latest stage at which we switched the stage type. We have 5 cases.

Case 1: s is of type $\mathcal{R} \vee \mathcal{R}$. If there exists some condition $(\sigma, \tau) \leq (\sigma_s, \tau_s)$ with $|\sigma|, |\tau| \leq s$ such that $(\sigma, \tau) \Vdash_0 (G_0 \oplus Y)' \upharpoonright_{s_0} = \sigma'$ for a $\sigma' \in 2^{s_0}$ or $(\sigma, \tau) \Vdash_1 (G_1 \oplus Y)' \upharpoonright_{s_0} = \tau'$ for a $\tau' \in 2^{s_0}$. Then let $\sigma_{s+1} = \sigma$, $\tau_{s+1} = \tau_s$, and if we are in the first case, let $\sigma'_{s+1} = \sigma'$ and $\tau'_{s+1} = \tau'_s$ and in the second case let $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'$. Then let $s+1$ be of type $\mathcal{R} \vee \mathcal{S}$. Otherwise, the elements are left unchanged and we go to the next stage.

Case 2: s is of type $\mathcal{R} \vee \mathcal{S}$. If there exists some condition $(\sigma, \tau) \leq (\sigma_s, \tau_s)$ with $|\sigma|, |\tau| \leq s$ such that $(\sigma, \tau) \Vdash_0 (G_0 \oplus Y)' \upharpoonright_{s_0} = \sigma'$ for a $\sigma' \in 2^{s_0}$ or $(\sigma, \tau) \Vdash_1 (G_1 \oplus Y)' \upharpoonright_{s_0} = \tau'$ for a $\tau' \in 2^{s_0}$. Then let $\sigma_{s+1} = \sigma$, $\tau_{s+1} = \tau_s$, and if we are in the first case, let $\sigma'_{s+1} = \sigma'$ and $\tau'_{s+1} = \tau'_s$ and in the second case let $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'_s$. Then let $s+1$ be of type $\mathcal{S} \vee \mathcal{R}$. Otherwise, the elements are left unchanged and we go to the next stage.

Case 3: s is of type $\mathcal{S} \vee \mathcal{R}$. If there exists some condition $(\sigma, \tau) \leq (\sigma_s, \tau_s)$ with $|\sigma|, |\tau| \leq s$ such that $(\sigma, \tau) \Vdash_0 \Gamma_{s_0}^{G_0 \oplus Y} \text{ partial}$ or $(\sigma, \tau) \Vdash_1 (G_1 \oplus Y)' \upharpoonright_{s_0} = \tau'$ for a $\tau' \in 2^{s_0}$. Then let $\sigma_{s+1} = \sigma$, $\tau_{s+1} = \tau_s$, and if we are in the first case, let $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'_s$ and in the second case let $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'$.

Then let $s + 1$ be of type $\mathcal{S} \vee \mathcal{S}$. Otherwise, the elements are left unchanged and we go to the next stage.

Case 4: s is of type $\mathcal{S} \vee \mathcal{S}$. If there exists some condition $(\sigma, \tau) \leq (\sigma_s, \tau_s)$ with $|\sigma|, |\tau| \leq s$ such that $(\sigma, \tau) \Vdash_0 \Gamma_{s_0}^{G_0 \oplus Y} \text{partial}$ or $(\sigma, \tau) \Vdash_1 \Gamma_{s_0}^{G_1 \oplus Y} \text{partial}$. Then let $\sigma_{s+1} = \sigma$, $\tau_{s+1} = \tau_s$, and if we are in the first case, let $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'_s$ and in the second case let $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'_s$. Then let $s + 1$ be of type \mathcal{T} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 5: s is of type \mathcal{T} . If there exists some condition $(\sigma, \tau) \leq (\sigma_s, \tau_s)$ with $\#\sigma > \#\sigma_s$ and $\#\tau > \#\tau_s$, then let $\sigma_{s+1} = \sigma$, $\tau_{s+1} = \tau$, $\sigma'_{s+1} = \sigma'_s$ and $\tau'_{s+1} = \tau'_s$. Then let $s + 1$ be of type $\mathcal{R} \vee \mathcal{R}$. Otherwise, the elements are left unchanged and we go to the next stage.

This completes the construction.

Verification. Since the size of $\sigma_s, \tau_s, \sigma'_s$ and τ'_s are bounded by s , there is a $\Delta_2^0(Y)$ -formula $\phi(s)$ stating that the construction can be pursued up to stage s . Our construction implies that the set $\{s \mid \phi(s)\}$ is a cut, so by $\text{I}\Delta_2^0(Y)$, the construction can be pursued at every stage.

Let $G_0 = \bigcup_{s \in M} \sigma_s$ and $G_1 = \bigcup_{s \in M} \tau_s$. By Lemma 3.23, and the fact that U and V are M -infinite, each type of stage changes M -infinitely often. Thus, for M -infinitely many k 's, the four requirements $\mathcal{R}_k^0 \vee \mathcal{R}_k^1$, $\mathcal{S}_k^0 \vee \mathcal{R}_k^1$, $\mathcal{R}_k^0 \vee \mathcal{S}_k^1$ and $\mathcal{S}_k^0 \vee \mathcal{S}_k^1$ are satisfied and G_0 and G_1 are M -infinite.

So, for M -infinitely many k 's, $(\mathcal{R}_k^0 \wedge \mathcal{S}_k^0) \vee (\mathcal{R}_k^1 \wedge \mathcal{S}_k^1)$ is satisfied which gives us an $i < 2$ such that G_i is an ascending or descending sequence, $Y' \geq_T (G_i \oplus Y)'$ and therefore $\mathcal{M}[G_i] \models \text{B}\Sigma_2^0$ and $\mathcal{M}[G_i] \models \text{WF}(\alpha)$. This completes the proof of Proposition 3.17. \square

Theorem 3.26. $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0) + \text{SADS}$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$.

Proof. Immediate by Proposition 3.17 and Theorem 2.2. \square

4 Π_1^1 -conservation of RT_2^2

Thanks to the decomposition of Proposition 1.16, we can combine the conservation theorems developed in Section 2 and Section 3 to prove our main theorems. Yokoyama [33] proved the following amalgamation theorem:

Theorem 4.1 ([33]). *Let T_0, T_1, T_2 be Π_2^1 theories such that $T_0 \supseteq \text{RCA}_0$ and T_1 and T_2 are both Π_1^1 -conservative extensions of T_0 . Then $T_1 + T_2$ is a Π_1^1 -conservative extension of T_0 .*

In particular, letting T_0 be $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$, one can combine Theorem 2.15, Theorem 2.21, Theorem 3.16 and Theorem 3.26 to deduce that $\text{RCA}_0 + \text{RT}_2^2 + \text{WKL} + \text{WF}(\epsilon_0)$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\epsilon_0)$.

We shall however provide a direct proof using ω -extensions with the following proposition.

Proposition 4.2. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega_4^\alpha)$ topped by a set $Y \in S$, where $\alpha \in M$ is an ordinal $\leq \epsilon_0$. Then, for every coloring $f : [M]^2 \rightarrow 2$ in S and every set $P \gg Y'$ such that $\mathcal{M}[P] \models \text{RCA}_0^*$, there exists $G \subseteq M$ such that*

1. G is an M -infinite f -homogeneous set ;
2. $P \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Proof. By Fiori-Carones et al [11, Lemma 3.2], there exists a P -computable countable coded model of WKL_0^* containing Y' . In particular, there exists a set $P_0 \subseteq M$ such that $P \gg P_0 \gg Y'$. Let $\vec{R} = R_0, R_1, \dots$ be defined by $R_x = \{y : f(x, y) = 1\}$. By Proposition 2.9 applied to \mathcal{M} and P_0 , there exists some M -infinite \vec{R} -cohesive set $G_0 \subseteq M$ such that $P_0 \geq_T (G_0 \oplus Y)'$ and $\mathcal{M}[G_0] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega_3^\alpha)$. In particular, $f : [G_0]^2 \rightarrow 2$ is stable. One can see $f : [G_0]^2 \rightarrow 2$ as an instance of SEM. By Proposition 3.1 applied to $\mathcal{M}[G_0]$ and P , there exists some M -infinite f -transitive set $G_1 \subseteq G_0$ such that $P \geq_T (G_1 \oplus G_0 \oplus Y)'$ and $\mathcal{M}[G_1 \oplus G_0] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\omega_2^\alpha)$. One can see $f : [G_1]^2 \rightarrow 2$ as an instance of SADS. By Proposition 3.17 applied to $\mathcal{M}[G_0 \oplus G_1]$ and P , there exists some M -infinite f -homogeneous set $G \subseteq G_1$ such that $P \geq_T (G \oplus G_1 \oplus G_0 \oplus Y)'$ and $\mathcal{M}[G \oplus G_1 \oplus G_0] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$. \square

Corollary 4.3. *Fix $n \geq 2$. Let T_n be either $\text{B}\Sigma_n^0$ or $\text{I}\Sigma_n^0$. Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + T_n + \text{WF}(\omega_4^\alpha)$ topped by a set $Y \in S$, where $\alpha \in M$ is an ordinal $\leq \epsilon_0$. Then, for every coloring $f : [M]^2 \rightarrow 2$ in S , there exists $G \subseteq M$ such that*

1. G is an M -infinite f -homogeneous set ;
2. $\mathcal{M}[G] \models \text{RCA}_0 + T_n + \text{WF}(\alpha)$.

Proof. By Harrington; Simpson-Smith; Hajek (see Belanger [1]), every countable topped model of $\text{RCA}_0^* + T_{n-1}$ can be ω -extended into a model of $\text{RCA}_0^* + T_{n-1} + \text{WKL}$. Since $\mathcal{M}[Y'] \models \text{RCA}_0^* + T_{n-1}$, there is some $P \subseteq M$ such that $P \gg Y'$ and $\mathcal{M}[P] \models \text{RCA}_0^* + T_{n-1}$. By Proposition 4.2, there exists $G \subseteq M$ such that

1. G is an M -infinite f -homogeneous set ;
2. $P \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{WF}(\alpha)$.

Since every Σ_n^0 formula in $\mathcal{M}[G]$ is Σ_{n-1}^0 in $\mathcal{M}[P]$ and $\mathcal{M}[P] \models T_{n-1}$, then $\mathcal{M}[G] \models T_n$. \square

Main Theorem 1.6. Fix $n \geq 2$. Let T_n be either BS_n^0 or IS_n^0 . $\text{RCA}_0 + T_n + \text{RT}_2^2 + \text{WKL} + \text{WF}(\epsilon_0)$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + T_n + \text{WF}(\epsilon_0)$.

Proof. From Corollary 4.3, Proposition 2.16 and Theorem 2.2, we get the Π_1^1 -conservativity of $\text{WKL} + \text{RT}_2^2$ over $\text{BS}_2^0 + \text{WF}(\epsilon_0)$. \square

4.1 Preserving $\bigcup_n \text{WF}(\omega_n)$

In their proof that RT_2^2 does not imply IS_2^0 over RCA_0 , Chong, Slaman and Yang [6] started with a model satisfying a slightly weaker well-foundedness axiom than $\text{WF}(\epsilon_0)$, namely, $\bigcup_n \text{WF}(\omega_n)$. We show that our assumption can also be weakened to $\bigcup_k \text{WF}(\omega_k^\omega)$ with a little bit of extra work.

Theorem 4.4. Fix $n \geq 2$. Let T_n be either BS_n^0 or IS_n^0 . Let P be a Π_2^1 problem such that for every countable topped model $\mathcal{M} = (M, S) \models \text{RCA}_0 + T_n + \text{WF}(\omega_k^\alpha)$ with $\alpha \in M$ an ordinal less than ϵ_0 and k a standard integer and every instance of P in \mathcal{M} there exists a solution $G \subset M$ of that instance such that $\mathcal{M}[G] \models \text{RCA}_0 + T_n + \text{WF}(\alpha)$. Then P is Π_1^1 -conservative over $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega)$.

Proof. Assume $\text{RCA}_0 + T_n + \bigcup_{n \in \omega} \text{WF}(\omega_n) \not\models \forall X \phi(X)$ for ϕ arithmetical.

Extend the language of second order arithmetics with a first-order constant symbol c . The theory $\text{RCA}_0 + T_n + \text{WF}(\omega_c) + \bigcup_{k \in \omega} c > k + \exists X \neg \phi(X)$ is finitely satisfiable by our assumption, so by compactness we have a model $\mathcal{M} \models \text{RCA}_0 + T_n + \text{WF}(\omega_c) + \bigcup_{k \in \omega} c > k + \exists X \neg \phi(X)$. Such a c will therefore be non-standard.

From our assumption, for any countable topped model $\mathcal{N} = (N, S) \models \text{RCA}_0 + T_n + \text{WF}(\omega_{d+k})$ and instance of P in this model, we can find a countable topped ω -extension that will be a model of $\text{RCA}_0 + T_n + \text{WF}(\omega_d)$ and will contain a solution to that instance.

So by iterating that reasoning, we can build a sequence $\mathcal{M}_0 = \mathcal{M} \subseteq \mathcal{M}_1 \subseteq \dots$ of countable topped model such that $\mathcal{M}_s \models \text{RCA}_0 + T_n + \text{WF}(\omega_{c-ks})$ and such that every instance of P appearing inside one \mathcal{M}_i will eventually have a solution in one \mathcal{M}_j . Since c is non-standard and every such s is standard, every \mathcal{M}_i is a model of $\bigcup_{k \in \omega} \text{WF}(\omega_k^\omega)$, and the model $\bigcup_{s \in \omega} \mathcal{M}_s$ will therefore be a model of $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega) + \exists X \neg \phi(X) + P$.

So $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega) + P \not\models \forall X \phi(X)$. \square

Main Theorem 1.7. Fix $n \geq 2$. Let T_n be either BS_n^0 or IS_n^0 . $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega) + \text{RT}_2^2 + \text{WKL}$ is Π_1^1 -conservative over $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega)$.

Proof. From Corollary 4.3, Proposition 2.16 and Theorem 4.4, we get the Π_1^1 -conservativity of $\text{RCA}_0 + T_n + \text{RT}_2^2 + \text{WKL}$ over $\text{RCA}_0 + T_n + \bigcup_{k \in \omega} \text{WF}(\omega_k^\omega)$. \square

Remark 4.5. The combination of Proposition 2.9, Proposition 3.1 and Proposition 3.17 provides the \ll_2 -basis theorem for RT_2^2 while preserving $\text{WF}(\epsilon_0)$ (see [11, Definition 4.9]). This implies that there exists a polynomial proof transformation for the Π_1^1 -conservation for Main Theorem 1.6 (it will appear in the forthcoming thesis of H. Ikari). On the other hand, it is not clear whether there exists a polynomial proof transformation for Main Theorem 1.7.

4.2 Preserving $\text{I}\Sigma_n^0$ and $\text{B}\Sigma_{n+1}^0$ for $n \geq 2$

One can use the previous theorems to reprove a known conservation result over $\text{I}\Sigma_2^0$.

Proposition 4.6. *Fix $n \geq 2$ and let T_n be either $\text{I}\Sigma_n^0$, or $\text{B}\Sigma_{n+1}^0$. Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + T_{n-1}$ topped by a set $Y \in S$. Then, for every coloring $f : [M]^2 \rightarrow 2$ in S and every set $P \gg Y'$ such that $\mathcal{M}[P] \models \text{RCA}_0$, there exists $G \subseteq M$ such that*

1. G is an M -infinite f -homogeneous set ;
2. $P \geq_T (G \oplus Y)'$;
3. $\mathcal{M}[G] \models \text{RCA}_0 + T_n$.

Proof. By Fiori-Carones et al [11, Lemma 3.2], there exists a P -computable countable coded model of WKL_0^* containing Y' . In particular, there exists a set $P_0 \subseteq M$ such that $P \gg P_0 \gg Y'$. Moreover, $\mathcal{M}[P]$ and $\mathcal{M}[P_0]$ both satisfy T_{n-1} .

Let $\vec{R} = R_0, R_1, \dots$ be defined by $R_x = \{y : f(x, y) = 1\}$. By Proposition 2.9 applied to \mathcal{M} , P_0 and $\alpha = 1$, there exists some M -infinite \vec{R} -cohesive set $G_0 \subseteq M$ such that $P_0 \geq_T (G_0 \oplus Y)'$ and $\mathcal{M}[G_0] \models \text{RCA}_0 + \text{B}\Sigma_2^0$. In particular, $f : [G_0]^2 \rightarrow 2$ is stable. Since any $\Sigma_n^0(G_0 \oplus Y)$ formula is $\Sigma_{n-1}^0(P_0)$, and $\mathcal{M}[P_0] \models T_{n-1}$, then $\mathcal{M}[G_0] \models T_n$. In particular, since $\text{RCA}_0 + T_n \vdash \text{WF}(\omega^\omega)$, $\mathcal{M}[G_0] \models \text{WF}(\omega^\omega)$.

One can see $f : [G_0]^2 \rightarrow 2$ as an instance of SEM. By Proposition 3.1 applied to $\mathcal{M}[G_0]$, P and $\alpha = 1$, there exists some M -infinite f -transitive set $G_1 \subseteq G_0$ such that $P \geq_T (G_1 \oplus G_0 \oplus Y)'$. Here again, since any $\Sigma_n^0(G_1 \oplus G_0 \oplus Y)$ formula is $\Sigma_{n-1}^0(P)$ and $\mathcal{M}[P] \models T_{n-1}$, then $\mathcal{M}[G_1 \oplus G_0] \models T_n$.

One can see $f : [G_1]^2 \rightarrow 2$ as an instance of SADS. By Proposition 3.17 applied to $\mathcal{M}[G_0 \oplus G_1]$, P and $\alpha = 0$, there exists some M -infinite f -homogeneous set $G \subseteq G_1$ such that $P \geq_T (G \oplus G_1 \oplus G_0 \oplus Y)'$ and $\mathcal{M}[G \oplus G_1 \oplus G_0] \models \text{RCA}_0 + \text{B}\Sigma_2^0$. Still by the same argument, $\mathcal{M}[G \oplus G_1 \oplus G_0] \models T_n$. \square

Remark 4.7. *Contrary to Proposition 2.9 and Proposition 3.17 in which the well-foundedness hypothesis on the ground model is only used to preserve well-foundedness, in the proof of Proposition 3.1, one exploits $\text{WF}(\omega^\omega)$ to preserve $\text{B}\Sigma_2^0$, and one uses $\text{WF}(\omega^{\omega \times \omega})$ to preserve $\text{WF}(\omega^\omega)$. If one starts with a model of $\text{I}\Sigma_2^0$, the ground model is therefore also a model of $\text{WF}(\omega^\omega)$, in which case $\text{B}\Sigma_2^0$ can be preserved. Thanks to the first-jump control, $\text{I}\Sigma_2^0$ is actually preserved, and therefore so is $\text{WF}(\omega^\omega)$.*

Since $\text{I}\Sigma_2^0$ and $\text{WF}(\omega^{\omega \times \omega})$ are incomparable over RCA_0 , it would be interesting to investigate the common factor between these two statements which is sufficient to preserve $\text{WF}(\omega^\omega)$.

Corollary 4.8. *Fix $n \geq 2$ and let T_n be either $\text{I}\Sigma_n^0$, or $\text{B}\Sigma_{n+1}^0$. Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + T_n$ topped by a set $Y \in S$. Then, for every coloring $f : [M]^2 \rightarrow 2$ in S , there exists $G \subseteq M$ such that*

1. G is an M -infinite f -homogeneous set ;
2. $\mathcal{M}[G] \models \text{RCA}_0 + T_n$.

Proof. By Harrington; Simpson–Smith; Hajek (see Belanger [1]), every countable topped model of $\text{RCA}_0^* + T_{n-1}$ can be ω -extended into a model of $\text{RCA}_0^* + T_{n-1} + \text{WKL}$. Since $\mathcal{M}[Y'] \models \text{RCA}_0^* + T_{n-1}$, there is some $P \subseteq M$ such that $P \gg Y'$ and $\mathcal{M}[P] \models \text{RCA}_0^* + T_{n-1}$. The conclusion follows from Proposition 4.6. \square

Corollary 4.9 (Cholak, Jockusch and Slaman [3]). *Fix $n \geq 2$ and let T_n be either IS_n^0 , or BS_{n+1}^0 . $\text{RCA}_0 + T_n + \text{RT}_2^2$ is Π_1^1 -conservative over $\text{RCA}_0 + T_n$.*

5 Open questions

Many open questions remain around the first-order part of Ramsey’s theorem for pairs and its consequences. The first and most important question being the following:

Question 5.1. *Is $\text{RCA}_0 + \text{RT}_2^2$ Π_1^1 -conservative over $\text{RCA}_0 + \text{BS}_2^0$?*

For every $\alpha < \epsilon_0$, $\text{RCA}_0 + \text{BS}_2^0 + \text{WF}(\alpha) + \text{WKL} + \text{COH}$ is Π_1^1 conservative over $\text{RCA}_0 + \text{BS}_2^0 + \text{WF}(\alpha)$. The situation seems more complex for ADS and EM, which both seem to require $\text{WF}(\omega^{\alpha \times \omega})$ to preserve $\text{WF}(\alpha)$.

Question 5.2. *For which $\alpha < \epsilon_0$ is $\text{RCA}_0 + \text{WF}(\alpha) + \text{RT}_2^2$ Π_1^1 -conservative over $\text{RCA}_0 + \text{BS}_2^0 + \text{WF}(\alpha)$?*

The proofs of Proposition 3.1 and Proposition 3.17 required the ground model to satisfy $\text{WF}(\omega^\alpha)$ to preserve $\text{WF}(\alpha)$. It is natural to wonder whether this assumption is necessary. One way to formalize this idea would be to prove that $\text{RCA}_0 + \text{RT}_2^2 \rightarrow \forall n (\text{WF}(\omega_n) \rightarrow \text{WF}(\omega_{n+1}))$, where $\omega_0 = 1$ and $\omega_{n+1} = \omega^{\omega_n}$. This is not the case, as $\text{RCA}_0 + \text{RT}_2^2$ is $\forall \Pi_3^0$ -conservative over RCA_0 , hence does not prove $\text{WF}(\omega^\omega)$. One could also wonder whether $\text{RCA}_0 + \text{RT}_2^2$ proves some weaker well-foundedness closure properties which do not already follow from $\text{RCA}_0 + \text{BS}_2^0$. Here again, the answer is negative. Le Hou  rou and Levy Patey constructed a model $\mathcal{M} = (M, S)$ of $\text{RCA}_0 + \text{RT}_2^2$ in which $\{b \in M : \mathcal{M} \models \text{WF}(\omega^b)\} = \sup\{a \cdot k : k \in \omega\}$ for some $a \in M \setminus \omega$. In other words, this cut has no better closure properties than additivity [personal communication]. This result is tight, as $\text{RCA}_0 \vdash \forall a (\text{WF}(\omega^a) \rightarrow \text{WF}(\omega^{2a}))$. Note that one can also show that $\text{RCA}_0 + \text{RT}_2^2$ does not prove $\forall n (\text{WF}(\omega^n) \rightarrow \text{WF}(\omega^{2^n}))$ using proof length. Indeed, by Ko  dziejczyk, Wong and Yokoyama [19], $\text{RCA}_0 + \text{RT}_2^2$ does not have speedup over RCA_0 for $\forall \Pi_3^0$ consequences, but in an incoming paper, Ko  dziejczyk and Yokoyama [personal communication] proved that $\text{RCA}_0 + \forall n (\text{WF}(\omega^n) \rightarrow \text{WF}(\omega^{2^n}))$ has super-exponential speedup over RCA_0 for $\forall \Pi_3^0$ consequences.

Acknowledgements

The authors are thankful to Leszek Kołodziejczyk for his careful reading and his improvement suggestions.

References

- [1] David R. Bélanger. Conservation theorems for the cohesiveness principle. 2022. Available at <https://arxiv.org/abs/2212.13011>.
- [2] Andrey Bovykin and Andreas Weiermann. The strength of infinitary Ramseyan principles can be accessed by their densities. *Ann. Pure Appl. Logic*, 168(9):1700–1709, 2017.
- [3] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey’s theorem for pairs. *The Journal of Symbolic Logic*, 66(1):1–55, 2001.
- [4] C. T. Chong and K. J. Mourad. The degree of a Σ_n cut. *Ann. Pure Appl. Logic*, 48(3):227–235, 1990.
- [5] C. T. Chong, Theodore A. Slaman, and Yue Yang. Π_1^1 -conservation of combinatorial principles weaker than Ramsey’s theorem for pairs. *Adv. Math.*, 230(3):1060–1077, 2012.
- [6] C. T. Chong, Theodore A. Slaman, and Yue Yang. The inductive strength of Ramsey’s Theorem for Pairs. *Adv. Math.*, 308:121–141, 2017.
- [7] Damir D. Dzharfarov and Carl Mummert. *Reverse mathematics—problems, reductions, and proofs*. Theory and Applications of Computability. Springer, Cham, [2022] ©2022.
- [8] Benedict Eastaugh. *Slicing the truth: on the computability theoretic and reverse mathematical analysis of combinatorial principles* [book review of MR3244278]. *Studia Logica*, 105(4):873–879, 2017.
- [9] P. Erdős and L. Moser. On the representation of directed graphs as unions of orderings. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 9:125–132, 1964.
- [10] António M. Fernandes, Fernando Ferreira, and Gilda Ferreira. Analysis in weak systems. In *Logic and computation*, volume 33 of *Tributes*, pages 231–261. Coll. Publ., [London], 2017.
- [11] Marta Fiori-Carones, Leszek Aleksander Kołodziejczyk, Tin Lok Wong, and Keita Yokoyama. An isomorphism theorem for models of Weak König’s Lemma without primitive recursion. *Journal of the European Mathematical Society*, 2021. To appear.

- [12] Petr Hájek. Interpretability and fragments of arithmetic. In *Arithmetic, proof theory, and computational complexity (Prague, 1991)*, volume 23 of *Oxford Logic Guides*, pages 185–196. Oxford Univ. Press, New York, 1993.
- [13] Petr Hájek and Pavel Pudlák. *Metamathematics of first-order arithmetic*, volume 3. Cambridge University Press, 2017.
- [14] Denis R. Hirschfeldt, Carl G. Jockusch, Jr., Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman. The strength of some combinatorial principles related to Ramsey’s theorem for pairs. In *Computational prospects of infinity. Part II. Presented talks*, volume 15 of *Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.*, pages 143–161. World Sci. Publ., Hackensack, NJ, 2008.
- [15] Denis R. Hirschfeldt and Richard A. Shore. Combinatorial principles weaker than Ramsey’s theorem for pairs. *Journal of Symbolic Logic*, 72:171 – 206, 2007.
- [16] Jeffry Lynn Hirst. *COMBINATORICS IN SUBSYSTEMS OF SECOND ORDER ARITHMETIC*. ProQuest LLC, Ann Arbor, MI, 1987. Thesis (Ph.D.)—The Pennsylvania State University.
- [17] Carl Jockusch and Frank Stephan. A cohesive set which is not high. *Math. Logic Quart.*, 39(4):515–530, 1993.
- [18] Carl G. Jockusch, Jr. Ramsey’s theorem and recursion theory. *J. Symbolic Logic*, 37:268–280, 1972.
- [19] Leszek Aleksander Kołodziejczyk, Tin Lok Wong, and Keita Yokoyama. Ramsey’s theorem for pairs, collection, and proof size. *Journal of Mathematical Logic*, page 2350007, 2023.
- [20] Leszek Aleksander Kołodziejczyk and Keita Yokoyama. In search of the first-order part of Ramsey’s theorem for pairs. In *Connecting with computability*, volume 12813 of *Lecture Notes in Comput. Sci.*, pages 297–307. Springer, Cham, [2021] ©2021.
- [21] Alexander P. Kreuzer. Primitive recursion and the chain antichain principle. *Notre Dame J. Form. Log.*, 53(2):245–265, 2012.
- [22] Alexander P. Kreuzer and Keita Yokoyama. On principles between Σ_1 - and Σ_2 -induction, and monotone enumerations. *J. Math. Log.*, 16(1):1650004, 21, 2016.
- [23] Manuel Lerman, Reed Solomon, and Henry Towsner. Separating principles below Ramsey’s theorem for pairs. *J. Math. Log.*, 13(2):1350007, 44, 2013.
- [24] Jiayi Liu. RT22 does not imply WKL0. *The Journal of Symbolic Logic*, 77(2):609–620, 2012.

- [25] Joseph Roy Mileti. *Partition theorems and computability theory*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)—University of Illinois at Urbana-Champaign.
- [26] Benoit Monin and Ludovic Patey. SRT_2^2 does not imply RT_2^2 in ω -models. *Adv. Math.*, 389:Paper No. 107903, 32, 2021.
- [27] Ludovic Patey. Partial orders and immunity in reverse mathematics. *Computability*, 7(4):323–339, 2018.
- [28] Ludovic Patey and Keita Yokoyama. The proof-theoretic strength of Ramsey’s theorem for pairs and two colors. *Adv. Math.*, 330:1034–1070, 2018.
- [29] David Seetapun and Theodore A. Slaman. On the strength of Ramsey’s theorem. volume 36, pages 570–582. 1995. Special Issue: Models of arithmetic.
- [30] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Perspectives in Logic. Cambridge University Press, 2 edition, 2009.
- [31] Stephen G. Simpson and Rick L. Smith. Factorization of polynomials and Σ_1^0 induction. volume 31, pages 289–306. 1986. Special issue: second Southeast Asian logic conference (Bangkok, 1984).
- [32] Theodore A. Slaman. Σ_n -bounding and Δ_n -induction. *Proc. Amer. Math. Soc.*, 132(8):2449–2456, 2004.
- [33] Keita Yokoyama. On conservativity for theories in second order arithmetic. In *Proceedings of the 10th Asian Logic Conference*, pages 375–386. World Scientific, 2010.