

THE REVERSE MATHEMATICS OF CARLSON'S THEOREM FOR LOCATED WORDS

TRISTAN BOMPARD, LU LIU, AND LUDOVIC PATEY

ABSTRACT. In this article, we give two proofs of Carlson's theorem for located words in ACA_0^+ . The first proof is purely combinatorial, in the style of Towsner's proof of Hindman's theorem. The second uses topological dynamics to show that an iterated version of Hindman's theorem for bounded sums implies Carlson's theorem for located words.

1. INTRODUCTION

We study the metamathematics of a partition theorem for words due to Carlson [3] from the viewpoint of reverse mathematics. Also our motivation is foundational, the metamathematical study of theorems in combinatorics usually consists in seeking for combinatorially simpler proofs of existing theorems. This is in particular the case of this article, where we give two new proofs of Carlson's theorem: a direct purely combinatorial one in the style of Towsner [25], and another from the Finite Union theorem using the tools of topological dynamics. This article might therefore be of interest to both combinatoricians who can ignore the metamathematical considerations, and logicians who wonder about the optimal axioms to prove Carlson's theorem.

1.1. Reverse mathematics

This is a foundational program started in 1974 by Harvey Friedman, whose goal is to study the optimal axioms for proving ordinary theorems. It uses the framework of subsystems of second-order arithmetics, with a base theory, RCA_0 , capturing *computable mathematics*. The early study of reverse mathematics has shown the emergence of four systems of axioms, namely, WKL_0 , ACA_0 , ATR_0 and $\Pi_1^1\text{-CA}_0$, listed in increasing order in terms of logical strength, such that most theorems are either provable in RCA_0 (hence computably true), or equivalent modulo RCA_0 to one of these four systems. This observation is known as the *Big Five phenomenon*. See Simpson [23] for a reference book on the early reverse mathematics.

The study of combinatorial theorems, especially coming from Ramsey's theory, has relativized the Big Five phenomenon. Ramsey's theorem for pairs is the most famous example of theorem which fails this observation in a strong sense: its logical strength is strictly in between RCA_0 and ACA_0 (see Specker [24], Jockusch [17] and Seetapun and Slaman [22]), and incomparable with WKL_0 (see Jockusch [17] and Liu [20]). Combinatorial theorems are notoriously hard to study in reverse mathematics, and often require to find new elementary proofs of existing theorems. This is in particular the case of Hindman's theorem, whose logical strength is still an active study of research. See Hirschfeldt [16] or Dzhafarov and Mummert [8] for an introduction to the reverse mathematics of combinatorial theorems.

1.2. Hindman's theorem

Hindman's theorem [13] is a partition theorem about the integers. Given a set $X \subseteq \mathbb{N}$, we let $\text{FS}(X)$ be the set of non-empty finite sums of distinct elements from X , that is,

$$\text{FS}(X) = \{\Sigma F : F \subseteq X \wedge 0 < |F| < \infty\}$$

A *finite coloring* of a set X is a function of the form $X \rightarrow C$, where C is a finite set, identified as a set of colors. Given a finite coloring $f : X \rightarrow C$, we say that a subset $Y \subseteq X$ is *f-homogeneous* if every element in Y is given the same color by f .

Theorem 1.1 (Hindman). *For every finite coloring $f : \mathbb{N} \rightarrow C$, there is an infinite set $Y \subseteq \mathbb{N}$ such that $\text{FS}(Y)$ is f-homogeneous.*

We shall refer to the previous theorem as Hindman's theorem (HT). There exist multiple proofs of Hindman's theorem. The original proof from Hindman [13], a short proof from Baumgartner [1], an ultrafilter proof from Galvin and Glazer (see Hindman and Strauss [15]), a proof using topological dynamics by Furstenberg and Weiss [10], and a simple proof by Towsner [25]. The two first proofs were analyzed in reverse mathematics by Blass, Hirst, and Simpson [2]. They showed that the original proof from Hindman holds in ACA_0^+ , while Baumgartner's proof can be formalized in the much stronger system $\Pi_2^1\text{-TI}_0$. The analysis of the proofs from Galvin and Glazer and from Furstenberg and Weiss are more tricky, since they use third-order objects : ultrafilters in the former case, and Ellis enveloping semigroups in the latter case. Montalbán and Shore [21] studied the ultrafilter proof using the tools of conservativity, and proved that the existence of an idempotent ultrafilter is a conservative extension of ACA_0 augmented with an iterated version of Hindman's theorem. Kreuzer [19] studied the existence of idempotent ultrafilters from the viewpoint of higher-order reverse mathematics, and proved that iterated Hindman's theorem is equivalent to the Auslander-Ellis theorem in topological dynamics. Lastly, the proof from Towsner holds in ACA_0^+ . The exact reverse mathematical strength of Hindman's theorem remains one of the biggest open questions in reverse mathematics.

Hindman's theorem is a typical example where the quest for optimal axioms motivates the search for new elementary proofs of existing theorems. This motivated for example the purely combinatorial proof by Towsner [25], which is arguably elementary as it is obtained by combining and iterating only very simple constructions. On the other hand, reverse mathematics is sensitive to the operation of iterating constructions. Iterations are very elementary from a mathematical perspective, but this yields sometimes computationally complex objects which then require strong existence axioms.

Hindman's theorem is equivalent to a partition theorem about finite sets, known as the Finite Union theorem. Let $\mathcal{P}_f(\mathbb{N})$ be the set of all non-empty finite subsets of \mathbb{N} . A subset $X \subseteq \mathcal{P}_f(\mathbb{N})$ is a *block sequence* if for every $E, F \in X$, either $\max E < \min F$, or $\max F < \min E$. Given a set $X \subseteq \mathcal{P}_f(\mathbb{N})$, we write $\text{FU}(X)$ for the set of all non-empty finite unions of elements from X , that is,

$$\text{FU}(X) = \{\cup F : F \subseteq X \wedge 0 < |F| < \infty\}$$

Theorem 1.2 (Finite Union). *For every finite coloring $f : \mathcal{P}_f(\mathbb{N}) \rightarrow C$, there is an infinite block sequence $Y \subseteq \mathcal{P}_f(\mathbb{N})$ such that $\text{FU}(Y)$ is f -homogeneous.*

The Finite Union theorem (FUT) is known to be equivalent to Hindman's theorem over RCA_0 , through the one-to-one correspondence which to $n \in \mathbb{N}$ associates the finite set $F \in \mathcal{P}_f(\mathbb{N})$ such that $\sum_{i \in F} 2^i = n$. The Finite Union theorem is more convenient than Hindman's theorem from many viewpoints, as the set $\text{FU}(Y)$ is closed under finite unions, while the set $\text{FS}(Y)$ is not closed under finite sums.

1.3. Carlson's theorem for words

Carlson's theorem for words [3] is a unifying theorem generalizing both Hindman's theorem and the Hales-Jewett theorem. It comes from a long line of partition theorems about variable words, among which the Hales-Jewett theorem [12], the Graham-Rothschild theorem [11] and the Carlson-Simpson lemma [4].

Definition 1.3. A *variable word* over an alphabet A is a finite sequence w over $A \cup \{\star\}$, where \star is a variable which occurs at least once in w . Given a variable word w and a letter $a \in A$, we write $w[a]$ for the string of length $|w|$ where all the occurrences of \star are replaced by a .

We write $W(A)$ and $V(A)$ for the sets of all words and variable words over A , respectively. The most basic – and arguably the most important – theorem about variable words is the Hales-Jewett theorem. It plays the same role on the semigroup of words as the pigeonhole principle does in the semigroup of integers.

Theorem 1.4 (Hales-Jewett). *Fix a finite alphabet A . For every finite coloring $f : W(A) \rightarrow C$, there is a variable word w such that $\{w[a] : a \in A\}$ is f -homogeneous.*

The Hales-Jewett theorem was generalized to arbitrary finite dimensions by Graham and Rothschild, and to infinite dimensions by Carlson and Simpson. Carlson's theorem for words generalizes both the Carlson-Simpson lemma and Hindman's theorem as follows.

Definition 1.5. Given a finite or infinite sequence of variable words $(w_n)_{n < \ell}$ with $\ell \in \mathbb{N} \cup \{\mathbb{N}\}$, a word $u \in W(A \cup \{\star\})$ is *extracted*¹ from $(w_n)_{n < \ell}$ if there exists a finite sequence of indices $j_0 < \dots < j_{n-1}$ and a finite sequence of letters $a_0, \dots, a_{n-1} \in A \cup \{\star\}$ such that

$$u = w_{j_0}[a_0]w_{j_1}[a_1] \dots w_{j_{n-1}}[a_{n-1}]$$

We write $\langle (w_n)_{n < \ell} \rangle_A$ and $\langle (w_n)_{n < \ell} \rangle_{A\star}$ for the set of all words and variable words extracted from $(w_n)_{n < \ell}$, respectively.

Theorem 1.6 (Carlson for words). *For every finite coloring $f : W(A) \rightarrow C$, there is an infinite sequence $(w_n)_{n \in \mathbb{N}}$ of variable words such that $\langle (w_n)_{n \in \mathbb{N}} \rangle_A$ is f -homogeneous.*

Carlson's theorem for words implies Hindman's theorem by considering colorings which depend only on the length of the words. Indeed, for every $f : \mathbb{N} \rightarrow C$, letting $g : W(A) \rightarrow C$ be defined by $g(u) = f(|u|)$, then for any infinite g -homogeneous sequence of variable words $(w_n)_{n \in \mathbb{N}}$, letting $Y = \{|w_n| = n \in \mathbb{N}\}$, the set $\text{FS}(Y)$ is f -homogeneous.

Remark 1.7. Carlson [3] actually proved a stronger theorem about finite colorings of variable words: For every finite coloring $f : V(A) \rightarrow C$, there is an infinite sequence $(w_n)_{n \in \mathbb{N}}$ of variable words such that $\langle (w_n)_{n \in \mathbb{N}} \rangle_{A\star}$ is f -homogeneous. This statement, that we shall refer as Carlson's theorem for variable words, implies its version for words. The only known proofs of Carlson's theorem for variable words involve ultrafilters [3] or the Ellis envelopping semigroup [9].

While Carlson's theorem for variable words is only known to admit an ultrafilter proof and a topological dynamical proof, Karagiannis [18] gave a Baumgartner-style proof of Carlson's theorem for words. Despite their combinatorial simplicity, Baumgartner-style proofs involve strong set existence axioms from a metamathematical viewpoint. In this article, we give an alternative proof of Carlson's theorem for words in the style of Towsner [25], which yields the same known upper bound as Hindman's theorem, namely, ACA_0^+ .

Carlson's theorem for words admits a formulation in terms of located words, which plays the same role as the Finite Union theorem for Hindman's theorem.

Definition 1.8. Fix a finite alphabet.

- (1) A *located word* over A is a function from a finite nonempty subset of \mathbb{N} into A . Let FIN_A be the collection of all located words over A .
- (2) A *located variable word* over A is a finite partial function from \mathbb{N} into $A \cup \{\star\}$, that takes the value \star at least once. Let $\text{FIN}_{A\star}$ be the collection of all located variable words.

Note that we have $\text{FIN}_{A \cup \{\star\}} = \text{FIN}_A \sqcup \text{FIN}_{A\star}$. Given a located variable word $p \in \text{FIN}_{A\star}$ and a letter $a \in A$, we write $p[a]$ for the located words with $\text{dom}(p[a]) = \text{dom } p$ and $p[a](x) = a$ if $x = \star$, and $p[a](x) = p(x)$ otherwise. We also let $p[\star] = p$. We equip the collections FIN_A and $\text{FIN}_{A\star}$ with a partial ordering defined as

$$p < q \text{ if } \max \text{dom } p < \min \text{dom } q$$

Definition 1.9. A *block sequence* is a totally ordered set $X \subseteq \text{FIN}_{A\star}$. Given a block sequence X , we let

- (1) $[X]_A = \{p_0[a_0] \cup \dots \cup p_k[a_k] \in \text{FIN}_A : p_0, \dots, p_k \in \text{FIN}_{A\star}, a_0, \dots, a_k \in A\}$
- (2) $[X]_{A\star} = \{p_0[a_0] \cup \dots \cup p_k[a_k] \in \text{FIN}_{A\star} : p_0, \dots, p_k \in \text{FIN}_{A\star}, a_0, \dots, a_k \in A \cup \{\star\}\}$

In other words, $[X]_{A\star} = [X]_{A \cup \{\star\}} \cap \text{FIN}_{A\star}$.

Theorem 1.10 (Carlson for located words). *For every finite coloring $f : \text{FIN}_A \rightarrow C$, there is an infinite block sequence $X \subseteq \text{FIN}_{A\star}$ such that $[X]_A$ is f -homogeneous.*

¹The terminology comes from [7].

Carlson's theorem for located words implies its version for words by collapsing the domain of the partial functions to obtain an initial segment of \mathbb{N} . This implication will be formally proven in Proposition 2.15. Like the Finite Union theorem, Carlson's theorem for located words is more convenient to manipulate since $[X]_A$ is closed under finite unions.

1.4. Organization of the paper

In Section 2, we discuss various versions of Carlson's theorem for words and justify the use of located words by the ability to iterate the theorem. In Section 3, we give a Towsner-style proof of Carlson's theorem for located words. This proof can be formalized in ACA_0^+ . In Section 4, we develop some basics of topological dynamics for located words, as a preparation for Section 5. In Section 5 we state two versions of the Auslander-Ellis theorem for located words, and use them to give an alternative proof of Carlson's theorem for located words. We also prove that AET for located words follows from an iterated version of the Finite Union theorem. Since the latter theorem holds in ACA_0^+ , this yields a second proof of Carlson's theorem for located words in ACA_0^+ . Last, in Section 6, we state a few remaining open questions.

2. VERSIONS OF CARLSON'S THEOREM FOR WORDS

One desirable feature of a partition theorem is the ability to iterate it, to obtain a simultaneous solution to multiple instances. This can be achieved whenever the combinatorial space representing the solution is isomorphic to the original space. We then obtain a stronger statement, saying that any partition of a specific combinatorial space admits a combinatorial subspace within one of the parts. In this section, we study the corresponding strengthenings for the Hindman's theorem, Carlson's theorem for words, and justify why the Finite Union theorem and Carlson's theorem for located words is more convenient in terms of iterations. In what follows, we consider only partial semigroups where the product is defined only on distinct elements.

2.1. Strong Finite Union theorem and Hindman's theorem

In this section, we shall see that both the Finite Union theorem and Hindman's theorem are equivalent to their strong version over RCA_0 . However, there exists a natural semigroup isomorphism between the Finite Union theorem, its strong version and the strong version of Hindman's theorem, while the equivalence with Hindman's theorem requires some extra work.

The Finite Union theorem admits a natural iterable version thanks to the closure property of its combinatorial space.

Theorem 2.1 (Strong Finite Union). *For every infinite block sequence $X \subseteq \mathcal{P}_f(\mathbb{N})$ and every finite coloring $f : \text{FU}(X) \rightarrow C$, there is an infinite block sequence $Y \subseteq \text{FU}(X)$ such that $\text{FU}(Y)$ is f -homogeneous.*

The strong Finite Union theorem is an immediate consequence of the Finite Union theorem since for every infinite block sequence $X = \{F_0 < F_1 < \dots\} \subseteq \mathcal{P}_f(\mathbb{N})$, there is a natural isomorphism ι from $(\mathcal{P}_f(\mathbb{N}), \cup)$ to $(\text{FU}(X), \cup)$ defined by $\iota(E) = \bigcup_{n \in E} F_n$.

Lemma 2.2 (RCA_0). *The Finite Union theorem implies its strong version.*

Proof. Given $X = \{F_0 < F_1 < \dots\}$ and $f : \text{FU}(X) \rightarrow C$, let $g : \mathcal{P}_f(\mathbb{N}) \rightarrow C$ be defined by $g(E) = f(\iota(E))$. By the Finite Union theorem, there is an infinite block sequence $Z \subseteq \mathcal{P}_f(\mathbb{N})$ such that $\text{FU}(Z)$ is g -homogeneous. Let $Y = \{\iota(E) : E \in Z\}$. The set $Y \subseteq \text{FU}(X)$ is an infinite block sequence and $\text{FU}(Y) = \{\iota(E) : E \in \text{FU}(Z)\}$, so $\text{FU}(Y)$ is f -homogeneous. \square

It follows that the Finite Union theorem is equivalent to its strong version over RCA_0 . The strong version of Hindman's theorem requires more work, as given an infinite set $X \subseteq \mathbb{N}$ and an infinite set $Y \subseteq \text{FS}(X)$, $\text{FS}(Y)$ is not necessarily a subset of $\text{FS}(X)$.

Theorem 2.3 (Strong Hindman). *For every infinite set $X \subseteq \mathbb{N}$ and every finite coloring $f : \text{FS}(X) \rightarrow C$, there is an infinite set Y such that $\text{FS}(Y) \subseteq \text{FS}(X)$ and $\text{FS}(Y)$ is f -homogeneous.*

Unlike the Finite Union theorem, there is no natural isomorphism from $(\mathbb{N}, +)$ to $(\text{FS}(X), +)$, unless it satisfies some extra conditions.

Definition 2.4 (Carlucci et al. [5]). Given $n = 2^{n_0} + \dots + 2^{n_{\ell-1}}$ with $n_0 < \dots < n_{\ell-1}$, we let $\lambda(n) = n_0$, $\mu(n) = n_{\ell-1}$. An infinite set $A = \{a_0 < a_1 < \dots\}$ is *2-apart* if for every n , $\mu(a_n) < \lambda(a_{n+1})$.

The notion of 2-apartness for sets of integers is equivalent to the notion of block sequence for sets of finite sets. Suppose $X \subseteq \mathbb{N}$ is infinite and 2-apart. Then there is an isomorphism ι from $(\mathbb{N}, +)$ to $(\text{FS}(X), +)$ defined by $\iota(\sum_{i \in E} 2^i) = \sum_{i \in E} a_i$, where $X = \{a_0 < a_1 < \dots\}$. The notion of 2-apartness is not preserved by subsets of finite sums in general: there exist sets $X, Y \subseteq \mathbb{N}$ such that X is 2-apart, $\text{FS}(Y) \subseteq \text{FS}(X)$ but Y is not 2-apart. The following stronger relation preserves 2-apartness.

Definition 2.5. Given two infinite sets $X, Y \subseteq \mathbb{N}$, we write $Y \leq_{\text{FS}} X$ if there is an infinite block sequence $H \subseteq \mathcal{P}_f(X)$ such that $Y = \{\sum F : F \in H\}$.

Note that if $Y \leq_{\text{FS}} X$, then $\text{FS}(Y) \subseteq \text{FS}(X)$. Moreover, if X is 2-apart, then so is Y . The following lemma was proven by Hindman [13] in his original paper.

Lemma 2.6 (Hindman, RCA_0). *For every infinite set $X \subseteq \mathbb{N}$, there is an infinite 2-apart set $Y \leq_{\text{FS}} X$.*

Proof. We first prove by induction over $k \in \mathbb{N}$ that for every infinite set $X \subseteq \mathbb{N}$, there is some $n \in \text{FS}(X)$ such that $\lambda(n) \geq k$. The case $k = 0$ is trivial by taking $n = \min X$. Suppose by induction hypothesis it holds for k , but not for $k+1$. Then for every finite set $F \subseteq X$, there is some $n \in \text{FS}(X \setminus F)$ such that $\lambda(n) = k$. Let F_0, F_1 be two non-empty subsets of X such that $\max F_0 < \min F_1$ and $\lambda(\sum F_i) = k$. Then $\lambda(\sum F_0 \cup F_1) > k$, contradicting our hypothesis. This proves our claim.

Using the claim, we construct an infinite sequence F_0, F_1, \dots of finite non-empty subsets of X such that $\max F_n < \min F_{n+1}$ and $\lambda(\sum F_{n+1}) > \lambda(\sum F_n)$. First, $F_0 = \{\min X\}$. Assume F_0, \dots, F_n are defined. By our claim, since $X \setminus \{0, 1, \dots, \max F_n\}$ is infinite, there is a finite set $F_{n+1} \subseteq X \setminus \{0, 1, \dots, \max F_n\}$ such that $\lambda(\sum F_{n+1}) > \lambda(\sum F_n)$. By construction, the set $Y = \{\sum F_n : n \in \mathbb{N}\}$ is 2-apart and $B \leq_{\text{FS}} A$. \square

We are now ready to prove strong Hindman's theorem from Hindman's theorem.

Lemma 2.7 (RCA_0). *Hindman's theorem implies its strong version.*

Proof. Given X and $f : \text{FS}(X) \rightarrow C$, by Lemma 2.6 there is an infinite 2-apart set $X_0 \leq_{\text{FS}} X$. Let ι be the canonical isomorphism from $(\mathbb{N}, +)$ to $(\text{FS}(X_0), +)$. Let $g : \text{FS}(\mathbb{N}) \rightarrow C$ be defined by $g(n) = f(\iota(n))$. By Hindman's theorem, there is an infinite set $Z \subseteq \mathbb{N}$ such that $\text{FS}(Z)$ is g -homogeneous. Let $Y = \{\iota(n) : n \in Z\}$. The set Y satisfies $\text{FS}(Y) \subseteq \text{FS}(X)$, and $\text{FS}(Y) = \{\iota(n) : n \in \text{FS}(Z)\}$, so $\text{FS}(Y)$ is f -homogeneous. \square

Thanks to the equivalence between Hindman's theorem and the Finite Union theorem, we have the following equivalence.

Proposition 2.8 (Folklore). *The following statements are equivalent over RCA_0 :*

- (1) *Hindman's theorem*
- (2) *Strong Hindman's theorem*
- (3) *Finite Union theorem*
- (4) *Strong Finite Union theorem*

Proof. We have already proven the equivalences (1) \leftrightarrow (2) and (3) \leftrightarrow (4). The proof of (3) \rightarrow (1) is immediate, using the isomorphism ι from $(\mathcal{P}_f(\mathbb{N}), \cup)$ to $(\mathbb{N}, +)$ defined by $\iota(E) = \sum_{i \in E} 2^i$. Last, (2) \rightarrow (3) is immediate, using the isomorphism ι from $(\text{FS}(X), +)$ to $(\mathcal{P}_f(\mathbb{N}), \cup)$ defined by $\iota(\sum_{i \in E} 2^i) = E$. \square

Remark 2.9. The previous considerations show that there exists a direct correspondance between the Finite Union theorem, its strong version, and the strong version of Hindman's theorem. On the other hand, the proof of any of these theorems from Hindman's theorem requires

a preliminary lemma, namely, Lemma 2.6. In particular, the Finite Union theorem is naturally equivalent to its strong version, while the equivalence between Hindman's theorem and its strong version is arguably less natural.

2.2. Strong Carlson's theorem for located words and words

Carlson's theorem for located words and for words both admit a strong version. As before, Carlson's theorem for located words is naturally equivalent to its strong version and to the strong version of Carlson's theorem for words. However, it remains open whether Carlson's theorem for words implies its strong version over RCA_0 .

Theorem 2.10 (Strong Carlson for located words). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_{A^\star} \rightarrow C$ be a coloring. There is an infinite block sequence $Y \subseteq [X]_{A^\star}$ such that $[Y]_{A^\star}$ is f -homogeneous.*

The proof of strong Carlson's theorem for located words from its weak version follows from the isomorphism ι from $(\text{FIN}_{A \cup \{\star\}}, \cup)$ to $([X]_{A \cup \{\star\}}, \cup)$ defined by $\text{dom } \iota(q) = \bigsqcup_{n \in \text{dom } q} \text{dom } p_n$ and for $m \in \text{dom } q$ and $n \in \text{dom } p_n$, $\iota(q)(n) = p_n(q(m))$, where $X = \{p_0 < p_1 < \dots\}$ is an infinite block sequence.

Lemma 2.11 (RCA_0). *Carlson's theorem for located words implies its strong version.*

Proof. Given $X = \{p_0 < p_1 < \dots\}$ and $f : [X]_A \rightarrow C$, let $g : \text{FIN}_A \rightarrow C$ be defined by $g(p) = f(\iota(p))$. By Carlson's theorem for located words, there is an infinite block sequence $Z \subseteq \text{FIN}_{A^\star}$ such that $[Z]_A$ is g -homogeneous. Let $Y = \{\iota(p) : p \in Z\}$. The set $Y \subseteq [X]_{A^\star}$ is an infinite block sequence and $[Y]_A = \{\iota(p) : p \in [Z]_A\}$, so $[Y]_A$ is f -homogeneous. \square

Theorem 2.12 (Strong Carlson for words). *Let $(w_n)_{n \in \mathbb{N}}$ be an infinite sequence of variable words over A and let $f : \langle (w_n)_{n \in \mathbb{N}} \rangle_A \rightarrow C$ be a finite coloring. There is an infinite sequence of variable words $(u_n)_{n \in \mathbb{N}}$ such that $\langle (u_n)_{n \in \mathbb{N}} \rangle_A \subseteq \langle (w_n)_{n \in \mathbb{N}} \rangle_A$ and $\langle (u_n)_{n \in \mathbb{N}} \rangle_A$ is f -homogeneous.*

Strong Carlson's theorem for words follows from Carlson's theorem for located words, using the surjective morphism ι from $(\text{FIN}_{A \cup \{\star\}}, \cup)$ to $(\langle (w_n)_{n \in \mathbb{N}} \rangle_{A \cup \{\star\}}, \cdot)$ defined by collapsing the domains as follows: $\iota(p) = w_{n_0}[p(n_0)] \cdots w_{n_{k-1}}[p(n_{k-1})]$ where $\text{dom } p = \{n_0 < \dots < n_{k-1}\}$.

Lemma 2.13 (RCA_0). *Carlson's theorem for located words implies strong Carlson's theorem for words.*

Proof. Fix $(w_n)_{n \in \mathbb{N}}$ and f . Let $g : \text{FIN}_A \rightarrow C$ be defined by $g(p) = f(\iota(p))$. By Carlson's theorem for located words, there is an infinite block sequence $X = \{p_0 < p_1 < \dots\} \subseteq \text{FIN}_{A^\star}$ such that $[X]_A$ is g -homogeneous. Let $(u_n)_{n \in \mathbb{N}}$ be defined by $u_n = \iota(p_n)$. Then $\langle (u_n)_{n \in \mathbb{N}} \rangle_A = \{\iota(p) : p \in [X]_A\}$, hence $\langle (u_n)_{n \in \mathbb{N}} \rangle_A$ is f -homogeneous. \square

As for Hindman's theorem, the strong version of Carlson's theorem for words implies Carlson's theorem for located words in a natural way. Indeed, let $(w_n)_{n \in \mathbb{N}}$ be the sequence of variable words over A defined by $w_n = \star \cdots \star$ with $|w_n| = 2^n$. Any element in $\langle (w_n)_{n \in \mathbb{N}} \rangle_{A \cup \{\star\}}$ can be uniquely written in the form $w_{n_0}[a_0] \cdots w_{n_{k-1}}[a_{k-1}]$, where $n_0 < \dots < n_{k-1} \in \mathbb{N}$ and $a_0, \dots, a_{k-1} \in A \cup \{\star\}$. Thus, there is a natural isomorphism ι from $(\langle (w_n)_{n \in \mathbb{N}} \rangle_{A \cup \{\star\}}, \cdot)$ to $(\text{FIN}_{A \cup \{\star\}}, \cup)$ defined by $\iota(w_{n_0}[a_0] \cdots w_{n_{k-1}}[a_{k-1}]) = p$ such that $\text{dom}(p) = \{n_0 < \dots < n_{k-1}\}$ and $p(n_i) = a_i$.

Lemma 2.14 (RCA_0). *Strong Carlson's theorem for words implies Carlson's theorem for located words.*

Proof. Let $(w_n)_{n \in \mathbb{N}}$ be the sequence of variable words over A defined as above and let ι be the corresponding isomorphism. Given a finite coloring $f : \text{FIN}_A \rightarrow C$, let $g : \langle (w_n)_{n \in \mathbb{N}} \rangle_A \rightarrow C$ be defined by $g(w) = f(\iota(w))$. By strong Carlson's theorem for words, there is an infinite sequence of variable words $(u_n)_{n \in \mathbb{N}}$ such that $\langle (u_n)_{n \in \mathbb{N}} \rangle_A \subseteq \langle (w_n)_{n \in \mathbb{N}} \rangle_A$ and $\langle (u_n)_{n \in \mathbb{N}} \rangle_A$ is g -homogeneous. Let $Y = \{\iota(u_n) : n \in \mathbb{N}\}$. Then Y is an infinite block sequence and $[Y]_A = \{\iota(w) : w \in \langle (u_n)_{n \in \mathbb{N}} \rangle_A\}$, hence $[Y]_A$ is f -homogeneous. \square

Put altogether, we obtain the following equivalences over RCA_0 .

Proposition 2.15. *The following statements are equivalent over RCA_0 :*

- (1) *Carlson's theorem for located words*
- (2) *Strong Carlson's theorem for located words*
- (3) *Strong Carlson's theorem for words*

Proof. We have proven (1) \rightarrow (3), (1) \rightarrow (2) and (3) \rightarrow (1). Last, (2) \rightarrow (1) since strong Carlson's theorem for located words is a generalization of Carlson's theorem for words. \square

As mentioned at the start of the section, we leave the following question open.

Question 2.16. Does CT for words imply CT for located words over RCA_0 ?

3. TOWNSNER-STYLE PROOF OF CARLSON'S THEOREM FOR LOCATED WORDS

As mentioned in the introduction, Hindman's theorem admits multiple proofs: an ultrafilter one from Galvin and Glazer (see Hindman and Strauss [15]), a simple one from Baumgartner [1] and a proof by Towsner [25]. The latter one yields the best known upper bound in terms of reverse mathematics of Hindman's theorem.

Carlson's theorem for words original proof involves ultrafilters, and Karagiannis [18] later gave a Baumgartner-style proof of it. In this section, we give a Towsner-style proof of Carlson's theorem for located words. Its reverse mathematical analysis shows that Carlson's theorem for located words holds in ACA_0^+ . We shall refine this analysis in Section 5 by showing that Carlson's theorem for located words follows from an iterated version of the Finite Union theorem, which is also known to hold in ACA_0^+ .

We will be using the following theorem as a blackbox.

Theorem 3.1 (Hales-Jewett, RCA_0). *Let A be a finite alphabet and $f : \text{FIN}_A \rightarrow C$ be a finite coloring. Then there is a located variable word $p \in \text{FIN}_{A^\star}$ such that $[p]_A$ is f -homogeneous.*

The previous theorem does not require $\text{B}\Sigma_2^0$ since it admits a finite combinatorial version which requires less induction.

Definition 3.2. Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_A \rightarrow C$ be a coloring. A block sequence $Y \subseteq [X]_{A^\star}$ is

- (1) *weakly f -thin for color $i \in C$* if for every $p \in [Y]_{A^\star}$, there is some $a \in A$ such that $f(p[a]) \neq i$
- (2) *f -thin for color $i \in C$* if for every $p \in [Y]_A$, $f(p) \neq i$.

Lemma 3.3 (RCA_0). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_A \rightarrow C$ be a coloring such that X is weakly f -thin for some color $i \in C$. Then there is an infinite block sequence $Y \subseteq [X]_{A^\star}$ which is f -thin for color i .*

Proof. We build a sequence of located variable words $q_0 < q_1 < \dots$ by induction. For every infinite block sequence X , let $\iota_X : \text{FIN}_{A \cup \{\star\}} \rightarrow [X]_{A \cup \{\star\}}$ be the canonical isomorphism.

Let $g : \text{FIN}_A \rightarrow C$ be defined by $g(p) = f(\iota_X(p))$. By the Hales-Jewett theorem (Theorem 3.1), there is a located variable word p over A such that $[p]_A$ is g -homogeneous. Let $q_0 = \iota_X(p)$. Then $[q_0]_A$ is f -homogeneous. Since $q_0 \in X$, then there is some $a \in A$ such that $f(q_0[a]) \neq i$, hence $\{q_0\}$ is f -thin for color i .

Assume $q_0 < \dots < q_{n-1}$ are located variable words over A such that $F = \{q_0, \dots, q_{n-1}\}$ is f -thin for color i . Let $g : \text{FIN}_A \rightarrow C^{|[F]_A|+1}$ be defined by

$$g(p) = \langle f(q \cup \iota_{X-F}(p)) : q \in [F]_A \cup \{\emptyset\} \rangle$$

By the Hales-Jewett theorem (Theorem 3.1), there is a located variable word p over A such that $[p]_A$ is g -homogeneous. Let $q_n = \iota_{X-F}(p)$. Fix some $q \in [F]_A \cup \{\emptyset\}$. Then $[q \cup q_n]_A$ is f -homogeneous. Since $q \cup q_n \in X$, then there is some $a \in A$ such that $f(q \cup q_n[a]) \neq i$, thus $[q \cup q_n]_A$ is f -thin for color i . Since it is the case for every $q \in [F]_A \cup \{\emptyset\}$, then $F \cup \{q_n\}$ is f -thin for color i . \square

The following definitions are direct adaptations of Towsner's notions of half matches and full matches to variable words (see [25]).

Definition 3.4. Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_A \rightarrow C$ be a coloring. Let $F \subseteq [X]_{A^\star}$ be a finite set and $Y \subseteq [X - F]_{A^\star}$ be an infinite block sequence. We say that

- (1) F *half-matches* Y for color $i \in C$ if for every $q \in [Y]_A$ such that $f(q) = i$, there is some $p \in F$ such that for every $a \in A$, $f(p[a] \cup q) = i$.
- (2) F *half-matches* Y if it half-matches Y for every color $i \in C$.
- (3) F *full-matches* Y if for every $q \in [Y]_A$, there is some $p \in F$ such that for every $a \in A$, $f(p[a]) = f(p[a] \cup q) = f(q)$.

Lemma 3.5 (RCA₀). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence, let $f : [X]_A \rightarrow C$ be a coloring, and $i \in C$ be a color. There is a finite set $F \subseteq [X]_{A^\star}$ and an infinite block sequence $Y \subseteq [X - F]_{A^\star}$ such that F half-matches Y for color i .*

Proof. Suppose first that for every finite block sequence $F \subseteq [X]_{A^\star}$, there is a located variable word $q \in [X - F]_{A^\star}$ and some $a \in A$ such that for every $p \in [F]_{A^\star}$, there is some $b \in A$ such that $f(p[b] \cup q[a]) \neq i$. Then build an infinite block sequence $\{p_0 < p_1 < \dots\} \subseteq [X]_{A^\star}$ such that for every n , there is some $a_n \in A$ such that for every $p \in [p_0, \dots, p_{n-1}]_{A^\star}$, there is some $b \in A$ such that $f(p[b] \cup p_n[a_n]) \neq i$. Let $Y = \{q_0 < q_1 < \dots\}$ be the block sequence defined by $q_n = p_{2n} \cup p_{2n+1}[a_{2n+1}]$. Note that $Y \subseteq [X]_{A^\star}$.

We claim that Y is weakly f -thin for color i . Let $p \in [Y]_{A^\star}$. Then $p = \bigcup_{n \in \text{dom } q} q_n[q(n)]$ for some located variable word $q \in \text{FIN}_{A^\star}$. Let $m = \max \text{dom } q$. We have $q_m[q(m)] = p_{2m}[q(m)] \cup p_{2m+1}[a_{2m+1}]$. Let $r = \bigcup_{n \in \text{dom } q \setminus \{m\}} q_n[q(n)]$. Then $p = r \cup p_{2m}[q(m)] \cup p_{2m+1}[a_{2m+1}]$, with $r \in [p_0, \dots, p_{2m-1}]_{A^\star}$, so by definition of $\{p_0 < p_1 < \dots\}$, there is some $b \in A$ such that $f(r[b] \cup p_{2m}[q(m)] \cup p_{2m+1}[a_{2m+1}]) \neq i$. In other words, there is some $b \in A$ such that $f(p[b]) \neq i$. Hence Y is weakly f -thin for color i . By Lemma 3.3, there is an infinite block sequence $Z \subseteq [Y]_{A^\star}$ which is f -thin for color i . In particular, $\min Z$ half-matches $Z - \{\min Z\}$ for color i .

Suppose now that there is a finite block sequence $F \subseteq [X]_{A^\star}$ such that for every located variable word $q \in [X - F]_{A^\star}$ and every $a \in A$, there is some $p \in [F]_{A^\star}$ such that for every $b \in A$, $f(p[b] \cup q[a]) = i$. Then by assumption, F half-matches $X - F$ for color i . \square

Lemma 3.6 (RCA₀ + IΣ₂⁰). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_A \rightarrow C$ be a coloring. There is a finite set $F \subseteq [X]_{A^\star}$ and an infinite block sequence $Y \subseteq [X - F]_{A^\star}$ such that F half-matches Y .*

Proof. Let $C = \{i_0, i_1, \dots, i_{|C|-1}\}$. We build a finite sequence of pairs

$$(F_0, Y_0), \dots, (F_{|C|}, Y_{|C|})$$

inductively as follows: Initially, $F_0 = \emptyset$ and $Y_0 = X$. Assuming F_s and Y_s have been defined, by Lemma 3.5 there is a finite set $F_{s+1} \subseteq [X_s]_{A^\star}$ and an infinite block sequence $Y_{s+1} \subseteq [X_s - F_{s+1}]_{A^\star}$ such that F_{s+1} half-matches Y_{s+1} for color i_s . Let $F = \bigcup_{s \leq |C|} F_s$ and $Y = Y_{|C|}$.

We claim that F half-matches Y . Let $q \in [Y]_A$. Let s be such that $f(q) = i_s$. Then since F_{s+1} half-matches Y_{s+1} for color i_s and $q \in [Y_{s+1}]_A$, there is some $p \in F_{s+1} \subseteq F$ such that for every $a \in A$, $f(p[a] \cup q) = i_s = f(q)$. \square

Lemma 3.7 (ACA₀). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_A \rightarrow C$ be a coloring. One of the following holds:*

- (1) *There is an infinite block sequence $Y \subseteq [X]_{A^\star}$ which is f -thin for some color $i \in C$.*
- (2) *There is a finite set $F \subseteq [X]_{A^\star}$ and an infinite block sequence $Y \subseteq [X - F]_{A^\star}$ such that F full-matches Y .*

Proof. Construct an infinite sequence of finite sets of located variable words F_1, F_2, \dots , an infinite sequence of infinite block sequences Y_0, Y_1, \dots and an infinite sequence of colorings f_0, f_1, \dots as follows.

Initially, $Y_0 = X$ and $f_0 = f$. Suppose Y_s and f_s have been defined. By Lemma 3.6, there is a finite set $F_{s+1} \subseteq [Y_s]_{A^\star}$ and an infinite block sequence $Y_{s+1} \subseteq [Y_s - F_{s+1}]_{A^\star}$ such that F_{s+1} half-matches Y_{s+1} for the coloring f_s . Let f_{s+1} be defined on $[X_{s+1}]_A$ by letting $f_{s+1}(q) = \langle p, f_s(q) \rangle$, where $p \in F_{s+1}$ is such that for every $a \in A$, $f_s(p[a] \cup q) = f_s(q)$.

Suppose first that there is some s such that for every $q \in [Y_{s+1}]_A$, there is some $p_1 \in F_1, \dots, p_s \in F_s$ and $p \in [p_1, \dots, p_s]_{A^\star}$ such that for every $a \in A$, $f(p[a]) = f(p[a] \cup q) = f(q)$. Then, letting

$$F = \bigcup_{p_1 \in F_1, \dots, p_s \in F_s} [p_1, \dots, p_s]_{A^\star}$$

the set F full-matches Y_{s+1} and we are done.

Suppose now that for every $s \in \mathbb{N}$, there is some $q_s \in [Y_{s+1}]_A$ such that for every $p_1 \in F_1, \dots, p_s \in F_s$ and $p \in [p_1, \dots, p_s]_{A^\star}$, there is some $a \in A$ such that if $f(p[a] \cup q_s) = f(q_s)$, then $f(p[a]) \neq f(q_s)$. By the pigeonhole principle, there is a color $i \in C$ and an infinite sequence of integers $s_0 < s_1 < \dots$ such that for every $r \in \mathbb{N}$, $f(q_{s_r}) = i$. By choice of the colorings f_0, f_1, \dots and the definitions of half-matches, for each such r , we can find a sequence $p_1 \in F_1, \dots, p_{s_r} \in F_{s_r}$ such that for every $p \in [p_1, \dots, p_{s_r}]_{A^\star}$ and every $a \in A$, $f(p[a] \cup q_{s_r}) = f(q_{s_r}) = i$. Then, by our supposition, for every $p \in [p_1, \dots, p_{s_r}]_{A^\star}$, there is some $a \in A$ such that $f(p[a]) \neq i$.

By weak König's lemma (which holds in ACA_0), there is an infinite block sequence $Y \subseteq [X]_{A^\star}$ which is weakly f -thin for color i . By Lemma 3.3, there is an infinite block sequence $Z \subseteq [Y]_{A^\star}$ which is f -thin for color i . \square

Lemma 3.8 (ACA_0). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $f : [X]_A \rightarrow C$ be a coloring. There is a finite set $F \subseteq [X]_{A^\star}$ and an infinite block sequence $Y \subseteq [X - F]_{A^\star}$ such that F full-matches Y .*

Proof. By induction on $|C|$. Applying Lemma 3.7 with $|C| = 1$, the second case must hold and we are done. Suppose now that $|C| > 0$ and that the property holds for $|C| - 1$. Applying Lemma 3.7, either the second case holds, in which case we are done, or the first case holds, and we apply the induction hypothesis. \square

Theorem 3.9 (ACA_0^+). *Let $X \subseteq \text{FIN}_{A^\star}$ be an infinite block sequence and let $(f_n)_{n \in \mathbb{N}}$ be a sequence of colorings $[X]_A \rightarrow C$. There is an infinite block sequence $Y \subseteq [X]_{A^\star}$ such that for every n , there is some finite set $F \subseteq Y$ such that $[Y - F]_A$ is f_n -homogeneous.*

Proof. As in Lemma 3.7, construct an infinite sequence of finite sets of located variable words F_1, F_2, \dots , an infinite sequence of infinite block sequences Y_0, Y_1, \dots and an infinite sequence of colorings g_0, g_1, \dots as follows.

Initially, $Y_0 = X$ and $g_0 = f_0$. Suppose Y_s and g_s have been defined. By Lemma 3.8, there is a finite set $F_{s+1} \subseteq [Y_s]_{A^\star}$ and an infinite block sequence $Y_{s+1} \subseteq [Y_s - F_{s+1}]_{A^\star}$ such that F_{s+1} full-matches Y_{s+1} for the coloring g_s . Let g_{s+1} be defined on $[X_{s+1}]_A$ by letting $g_{s+1}(q) = \langle p, g_s(q), f_{s+1}(q) \rangle$, where $p \in F_{s+1}$ is such that for every $a \in A$, $g_s(p[a]) = g_s(p[a] \cup q) = g_s(q)$.

For every s , there is a sequence $p_1 \in F_1, \dots, p_s \in F_s$ such that for each $t \leq s$, $[p_t, \dots, p_s]_A$ is f_t -homogeneous. By weak König's lemma, there is an infinite block sequence $Y \subseteq [X]_{A^\star}$ such that for every n , there is some finite set $F \subseteq Y$ such that $[Y - F]_A$ is f_n -homogeneous. \square

4. TOPOLOGICAL DYNAMICS IN $C^{\text{FIN}A}$

Topological dynamics studies recurrence phenomena in the context of compact topological spaces. Furstenberg and Weiss [10] have shown that it was a very powerful tool to study combinatorial theorems. They gave in particular a proof of Hindman's theorem based on the Auslander-Ellis theorem, a theorem of topological dynamics informally stating that for every finite coloring $f : \mathbb{N} \rightarrow C$, there exists another coloring $g : \mathbb{N} \rightarrow C$ similar to f , but which enjoys a strong recurrence property. The proof of the Auslander-Ellis theorem involved the notion of Ellis' enveloping semigroup, which is a third-order object related to the notion of ultrafilter.

Blass, Hirst and Simpson [2] studied the work of Furstenberg and Weiss in the context of reverse mathematics, and got rid of the use of Ellis' enveloping semigroup by using iterated

Hindman's theorem in the proof of the Auslander-Ellis theorem. Kreuzer [19] proved the equivalence between iterated Hindman's theorem, the Auslander-Ellis theorem and the existence of an idempotent ultrafilter in higher-order reverse mathematics.

The original work of Furstenberg and Weiss [10] was on finite colorings of the semigroup $(\mathbb{N}, +)$. Furstenberg and Katznelson [9] extended their work to the semigroup $(\mathcal{W}(A), \cdot)$ of words over a finite alphabet A , and proved Carlson's theorem for words and variable words using topological dynamics, using Ellis' enveloping semigroup.

In this section, we build on the work of Furstenberg and Katznelson and develop some basics of topological dynamics for located words. The tools will be used in Section 5 to state and prove multiple versions of the Auslander-Ellis theorem from the iterated Finite Union theorem.

Consider FIN_A as a partial semigroup with the \cup operation, defined only whenever $p < q$. Fix a finite set C of colors. C^{FIN_A} is the space of all mappings from FIN_A to C . Given $m \in \mathbb{N}$, $n \in \mathbb{N} \cup \{\infty\}$, we write $\text{FIN}_A(m, n)$ for the set of all located words $p \in \text{FIN}_A$ such that $m \leq \text{dom}(p) < n$. Given $\ell \in \mathbb{N}$ and $p \in \text{FIN}_A(\ell, \infty)$, let $S_p^\ell(f) : \text{FIN}_A(0, \ell) \rightarrow C$ be defined by $S_p^\ell(f)(q) = f(q \cup p)$. Whenever $p = \emptyset$, we simply write $S^\ell(f)$ for $S_\emptyset^\ell(f)$. We say that a finite coloring $h : \text{FIN}_A(0, \ell) \rightarrow C$ is a *factor* of $f : \text{FIN}_A \rightarrow C$ if there is some $p \in \text{FIN}_A$ such that $S_p^\ell(f) = h$.

4.1. Recurrence

Informally, a coloring is recurrent if any initial segment of it occurs arbitrarily far. The located words being non-linearly ordered, there exist multiple notions of recurrence for colorings of located words.

Definition 4.1. A coloring $f : \text{FIN}_A \rightarrow C$ is

- (1) *weakly recurrent* if for every $\ell \in \mathbb{N}$, there is some located word $p \in \text{FIN}_A(\ell, \infty)$ such that $S_p^\ell(f) = S^\ell(f)$.
- (2) *recurrent* if for every $\ell \in \mathbb{N}$, there is some located variable word $p \in \text{FIN}_{A^\star}(\ell, \infty)$ such that for each $a \in A$, $S_{p[a]}^\ell(f) = S^\ell(f)$.
- (3) *uniformly recurrent* if for every $\ell \in \mathbb{N}$, there is some $m > \ell$ such that for every $p \in \text{FIN}_A(m, \infty)$, there is some $q \in \text{FIN}_A(\ell, m)$ for which $S^\ell(f) = S_{q \cup p}^\ell(f)$.

The terminology suggests that uniform recurrence is a stronger notion than recurrence, although it is not obvious from the definitions. The following lemma shows that it is the case, thanks to the Hales-Jewett theorem.

Lemma 4.2 (RCA₀). *If f is uniformly recurrent, then it is recurrent.*

Proof. Fix $\ell \in \mathbb{N}$. Let $m > \ell$ be such that for every $p \in \text{FIN}_A(m, \infty)$, there is some $q \in \text{FIN}_A(\ell, m)$ for which $S^\ell(f) = S_{q \cup p}^\ell(f)$.

Let $\pi_m : \text{FIN}_A \rightarrow \text{FIN}_A(m, \infty)$ be the canonical bijection. Let $g : \text{FIN}_A \rightarrow \text{FIN}_A(\ell, m)$ be defined for every $p \in \text{FIN}_A(m, \infty)$ by letting $g(p)$ be the least $q \in \text{FIN}_A(\ell, m)$ (in any fixed order) such that $S_{q \cup \pi_m(p)}^\ell(f) = S^\ell(f)$. By the Hales-Jewett theorem (Theorem 3.1), there is a located variable word $v \in \text{FIN}_{A^\star}$ such that $[v]_A$ is g -homogeneous, for some color $q \in \text{FIN}_A(\ell, m)$. Let $w = q \cup \pi_m(v)$. In particular, for every $a \in A$, $g(v[a]) = q$, so $S_{w[a]}^\ell(f) = S_{q \cup \pi_m(v[a])}^\ell(f) = S^\ell(f)$. \square

The following lemma shows that recurrent colorings are simple instances of Carlson's theorem for located words, in that they admit solutions computable in the instances.

Lemma 4.3 (RCA₀). *For every recurrent coloring $f : \text{FIN}_A \rightarrow C$, there is an infinite block sequence $X \subseteq \text{FIN}_{A^\star}$ such that $[X]_A$ is f -homogeneous for color $f(\emptyset)$.*

Proof. We build a sequence $p_0 < p_1 < \dots$ of located variable words inductively. Since f is recurrent, there is some located variable word $p_0 \in \text{FIN}_{A^\star}$ such that for each $a \in A$, $S_{p_0[a]}^0(f) = S^0(f)$. In particular, $f(p_0[a]) = f(\emptyset)$ for every $a \in A$.

Assume $p_0 < \dots < p_n$ are located variable words such that $[p_0, \dots, p_n]_A$ is f -homogeneous for color $f(\emptyset)$. Let $\ell = 1 \max \text{dom } p_n$. Since f is recurrent, there is some located variable

word $p_{n+1} \in \text{FIN}_{A^\star}(\ell, \emptyset)$ such that for each $a \in A$, $S_{p_{n+1}[a]}^\ell(f) = S^\ell(f)$. Let $p \in [p_0, \dots, p_n]_A \cup \{\emptyset\}$. Note that $\max \text{dom } p < \ell$ and that $f(p) = f(\emptyset)$. Then for every $a \in A$,

$$f(p \cup p_{n+1}[a]) = S_{p_{n+1}[a]}^\ell(f)(p) = g(p) = f(\emptyset)$$

Thus, $[p_0, \dots, p_n, p_{n+1}]_A$ is f -homogeneous for color $f(\emptyset)$. \square

4.2. Minimal subshifts

In general, a dynamical system induces a notion of subshift, which is a class of colorings closed under application of the homeomorphism. In the case of colorings of located words, the situation is slightly more complex because of the partiality of the semigroup. We can however define some notion of subshift, and recover the standard properties saying that any minimal subshift contains only uniformly recurrent colorings.

Definition 4.4. A non-empty closed class $\mathcal{C} \subseteq C^{\text{FIN}_A}$ is a *subshift* if for every $f \in \mathcal{C}$, every $\ell \in \mathbb{N}$ and $p \in \text{FIN}_A(\ell, \infty)$, there is some $g \in \mathcal{C}$ such that $S^\ell(g) = S_p^\ell(f)$.

A subshift $\mathcal{C} \subseteq C^{\text{FIN}_A}$ is *minimal* if there is no subshift $\mathcal{D} \subsetneq \mathcal{C}$.

Lemma 4.5 (WKL₀). *If $\mathcal{C} \subseteq C^{\text{FIN}_A}$ is a minimal subshift and $f \in \mathcal{C}$, then f is uniformly recurrent.*

Proof. Suppose f is not uniformly recurrent. Then there is some $\ell \in \mathbb{N}$ such that for every bound $m > \ell$, there is some located word $p \in \text{FIN}_A(m, \infty)$ such that for every $q \in \text{FIN}_A(\ell, m)$, $S_{q \cup p}^\ell(f) \neq S^\ell(f)$. Let \mathcal{D} be the Π_1^0 class of all $g \in \mathcal{C}$ which does not contain $S^\ell(f)$ as a factor. Note that \mathcal{D} is a subshift. As $f \notin \mathcal{D}$, we have that $\mathcal{D} \subsetneq \mathcal{C}$. By minimality assumption, then $\mathcal{D} = \emptyset$. By WKL, there is a bound $m > \ell$ such that for every $g \in \mathcal{C}$, there is some $q \in \text{FIN}_A(\ell, m)$ such that $S_q^\ell(g) = S^\ell(f)$.

Fix some $p \in \text{FIN}_A(m, \infty)$. By definition of a subshift, there is some $g \in \mathcal{C}$ such that $S^m(g) = S_p^m(f)$. In particular, for every $q \in \text{FIN}_A(\ell, m)$, $S_q^\ell(g) = S_{q \cup p}^\ell(f)$. Since $g \in \mathcal{C}$, there is some $q \in \text{FIN}_A(\ell, m)$ such that $S_q^\ell(g) = S^\ell(f)$. But $S_q^\ell(g) = S_{q \cup p}^\ell(f)$, so $S_{q \cup p}^\ell(f) = S^\ell(f)$. This contradicts our initial choice of ℓ . \square

From a purely mathematical viewpoint, the existence of minimal subshifts follow from Zorn's lemma. Day [6] however proved that their existence follow from ACA₀ in the context of colorings of \mathbb{N} . The proof of the existence of minimal subshifts in our case follows exactly the same argument.

Lemma 4.6 (ACA₀, Day [6]). *Every subshift contains a minimal subshift.*

Proof. Let $\mathcal{C} \subseteq C^{\text{FIN}_A}$ be a subshift. Fix an enumeration of all finite colorings h_0, h_1, \dots of the form $\text{FIN}_A(0, \ell) \rightarrow C$ for some $\ell \in \mathbb{N}$. Define a sequence $\mathcal{C}_0 \supseteq \mathcal{C}_1 \supseteq \dots$ of subshifts as follows: $\mathcal{C}_0 = \mathcal{C}$. Given \mathcal{C}_i , let \mathcal{D} be the class of $f \in \mathcal{C}_i$ which does not contain h_i as a factor. If $\mathcal{D} \neq \emptyset$, then $\mathcal{C}_{i+1} = \mathcal{D}$, otherwise $\mathcal{C}_{i+1} = \mathcal{C}_i$. Then $\bigcap_i \mathcal{C}_i$ is a minimal subshift. \square

4.3. Orbit closures

The notion of orbit closure of a coloring is central in topological dynamics. There is no clear definition of orbit in the case of colorings of located words, but the notion of orbit closure can be directly defined.

Definition 4.7. The *orbit closure* of $f \in C^{\text{FIN}_A}$ is the collection $\overline{\text{Orb}(f)}$ of all $g \in C^{\text{FIN}_A}$ such that for every $\ell \in \mathbb{N}$, and every $q \in \text{FIN}_A(\ell, \infty)$, there is some $p \in \text{FIN}_A(\ell, \infty)$ such that $S_q^\ell(g) = S_p^\ell(f)$.

In particular, $\overline{\text{Orb}(f)}$ is a $\Pi_1^0(f')$ class and is the class of path of the following tree:

$$T_f = \bigcup_m \left\{ h \in C^{\text{FIN}_A(0, m)} : \begin{array}{l} \forall \ell < m \forall q \in \text{FIN}_A(\ell, m) \\ \exists p \in \text{FIN}_A(\ell, \infty) S_p^\ell(f) = S_q^\ell(h) \end{array} \right\}$$

Given a finite coloring $h : \text{FIN}_A(0, \ell) \rightarrow C$, we write $[h]$ for the class of all $f \in C^{\text{FIN}_A}$ such that $S^\ell(f) = h$.

Lemma 4.8 (RCA₀). *For every f , $\overline{\text{Orb}(f)}$ is a subshift.*

Proof. $\overline{\text{Orb}(f)}$ is clearly closed as $\overline{\text{Orb}(f)} = [T_f]$.

Fix some $g \in \overline{\text{Orb}(f)}$, $\ell \in \mathbb{N}$ and $p \in \text{FIN}_A(\ell, \infty)$. Let us show that there is some $h \in \overline{\text{Orb}(f)}$ such that $S^\ell(h) = S_p^\ell(g)$. Let h be defined by $h(q) = g(q \cup p)$ for $q \in \text{FIN}_A(0, \ell)$, and $h(q) = g(q)$ otherwise. By definition, $S^\ell(h) = S_p^\ell(g)$. We claim that $h \in \overline{\text{Orb}(f)}$. Fix some $\ell_1 \in \mathbb{N}$, some $q \in \text{FIN}_A(\ell_1, \infty)$. If $q \in \text{FIN}_A(0, \ell)$, then $S_q^{\ell_1}(h) = S_{q \cup p}^{\ell_1}(g)$. Since $g \in \overline{\text{Orb}(f)}$, there is some $p_1 \in \text{FIN}_A(\ell_1, \infty)$ such that $S_{q \cup p}^{\ell_1}(g) = S_{p_1}^{\ell_1}(f)$. In particular $S_q^{\ell_1}(h) = S_{p_1}^{\ell_1}(f)$. If $q \notin \text{FIN}_A(0, \ell)$, then $S_q^{\ell_1}(h) = S_q^{\ell_1}(g)$. Again, since $g \in \overline{\text{Orb}(f)}$, there is some $p_1 \in \text{FIN}_A(\ell_1, \infty)$ such that $S_q^{\ell_1}(g) = S_{p_1}^{\ell_1}(f)$. In both cases, $S_q^{\ell_1}(h) = S_{p_1}^{\ell_1}(f)$ for some $p_1 \in \text{FIN}_A(\ell_1, \infty)$. \square

Lemma 4.9 (ACA₀, Kreuzer [19]). *If f is uniformly recurrent, then $\overline{\text{Orb}(f)}$ is minimal.*

Proof. Suppose that $\overline{\text{Orb}(f)}$ is not minimal. Then there is some $g \in \overline{\text{Orb}(f)}$ such that $\overline{\text{Orb}(g)} \subsetneq \overline{\text{Orb}(f)}$. In particular, $f \notin \overline{\text{Orb}(g)}$, so there some $\ell \in \mathbb{N}$ such that $\overline{\text{Orb}(g)} \cap [S^\ell(f)] = \emptyset$.

Since f is uniformly recurrent, there is a bound $m > \ell$ such that for every $p \in \text{FIN}_A(m, \infty)$, there is some $q \in \text{FIN}_A(\ell, m)$ such that $S_{q \cup p}^\ell(f) = S^\ell(f)$. Since $g \in \overline{\text{Orb}(f)}$, there is some $p \in \text{FIN}_A(m, \infty)$ such that $S^m(g) = S_p^m(f)$. In particular, for every $q \in \text{FIN}_A(\ell, m)$, $S_q^\ell(g) = S_{q \cup p}^\ell(f)$. Let $q \in \text{FIN}_A(\ell, m)$ be such that $S_{q \cup p}^\ell(f) = S^\ell(f)$. Then $S_q^\ell(g) = S_{q \cup p}^\ell(f) = S^\ell(f)$. But by definition of a subshift, there is some $h \in \overline{\text{Orb}(Y)}$ such that $S^\ell(h) = S_q^\ell(g)$, contradicting the fact that $\overline{\text{Orb}(g)} \cap [S^\ell(f)] = \emptyset$. \square

Lemma 4.10 (ACA₀, Day [6]). *If $\overline{\text{Orb}(f)}$ is minimal, then f is uniformly recurrent.*

Proof. $\overline{\text{Orb}(f)}$ is a minimal subshift containing f . By Lemma 4.5, f is uniformly recurrent. \square

Lemma 4.11 (ACA₀). *For every $f \in C^{\text{FIN}_A}$, there is a coloring $g \in \overline{\text{Orb}(f)}$ such that $\overline{\text{Orb}(g)}$ is minimal.*

Proof. By Lemma 4.6, $\overline{\text{Orb}(f)}$ contains a minimal subshift $\mathcal{D} \subseteq \overline{\text{Orb}(f)}$. Let $g \in \mathcal{D}$. In particular, $\overline{\text{Orb}(g)} \subseteq \mathcal{D}$ is a subshift. By minimality of \mathcal{D} , $\overline{\text{Orb}(g)} = \mathcal{D}$. \square

5. AUSLANDER-ELLIS THEOREM FOR C^{FIN_A}

We have seen that recurrent colorings are simple instances of Carlson's theorem for located words. Although colorings are not recurrent in general, the Auslander-Ellis theorem says that every coloring is close to a recurrent coloring. Here again, due to the non-linearity of the ordering of words, there exists multiple notions of proximality.

Definition 5.1. Fix two colorings $f, g : \text{FIN}_A \rightarrow C$.

- (1) f and g are *weakly proximal* if for every $\ell \in \mathbb{N}$, there is a located word $p \in \text{FIN}_A(\ell, \infty)$ such that $S_p^\ell(g) = S_p^\ell(f)$.
- (2) f and g are *proximal* if for every $\ell \in \mathbb{N}$, there is a located variable word $p \in \text{FIN}_{A^\star}(\ell, \infty)$ such that for each $a \in A$, $S_{p[a]}^\ell(g) = S_{p[a]}^\ell(f)$.
- (3) g is *strongly proximal* to f if for every $\ell \in \mathbb{N}$, there is a located variable word $p \in \text{FIN}_{A^\star}(\ell, \infty)$ such that for each $a \in A$, $S^\ell(g) = S_{p[a]}^\ell(g) = S_{p[a]}^\ell(f)$.

Note that if f is recurrent, then it is strongly proximal to itself. There exists multiple candidate statements for adapting the Auslander-Ellis theorem to C^{FIN_A} since there exists three notions of proximality. Two of them are of interest:

Theorem 5.2 (Auslander-Ellis for C^{FIN_A}). *For every coloring $f : \text{FIN}_A \rightarrow C$, there is a uniformly recurrent coloring $g : \text{FIN}_A \rightarrow C$ weakly proximal to f .*

Theorem 5.3 (Weak Auslander-Ellis for C^{FIN_A}). *For every coloring $f : \text{FIN}_A \rightarrow C$, there is a coloring $g : \text{FIN}_A \rightarrow C$ strongly proximal to f .*

In what follows, let $HJ(k, \ell)$ be an integer large enough so that for every $h : \text{FIN}_A(HJ(k, \ell)) \rightarrow \ell$, there is a located variable word $p \in \text{FIN}_{A^\star}(HJ(k, \ell))$ such that $[p]_A$ is h -homogeneous. The following lemma shows that the Auslander-Ellis theorem for C^{FIN_A} implies its weak version.

Lemma 5.4 (RCA₀). *If g is uniformly recurrent and weakly proximal to f , then it is strongly proximal to f .*

Proof. Fix $\ell \in \mathbb{N}$. Since g is uniformly recurrent, there is some $m > \ell$ be such that for every $p \in \text{FIN}_A(m, \infty)$, there is some $q \in \text{FIN}_A(\ell, m)$ such that $S_{q \cup p}^\ell(g) = S^\ell(g)$. Let $N = HJ(k, |\text{FIN}_A(\ell, m)|)$. Since g is weakly proximal to f , there is some $p \in \text{FIN}_A(m + N, \infty)$ such that $S_p^{m+N}(g) = S_p^{m+N}(f)$. Let $h : \text{FIN}_A(m, m + N) \rightarrow \text{FIN}_A(\ell, m)$ be defined by $h(v) = q$ such that $S_{q \cup v \cup p}^\ell(g) = S^\ell(g)$. By the finite Hales-Jewett theorem, there is a located variable word $u \in \text{FIN}_A(m, m + N)$ and some $q \in \text{FIN}_A(\ell, m)$ such that for every $a \in A$, $h(u[a]) = q$. In other words, for every $a \in A$, $S_{q \cup u[a] \cup p}^\ell(g) = S^\ell(g)$. Note that $S_{q \cup u[a] \cup p}^\ell(g) = S_{q \cup u[a] \cup p}^\ell(f)$. Thus, letting $w = q \cup u[a] \cup p$, for every $a \in A$, $S^\ell(g) = S_{w[a]}^\ell(g) = S_{w[a]}^\ell(f)$. \square

Corollary 5.5 (RCA₀). *The Auslander-Ellis theorem for C^{FIN_A} implies its weak version.*

Proof. Immediate by Lemma 5.4. \square

5.1. Proof of Carlson's theorem for located words

The same way Furstenberg and Weiss [10] proved that the original Auslander-Ellis theorem implies Hindman's theorem, we will prove that the weak version of the Auslander-Ellis theorem for C^{FIN_A} implies Carlson's theorem for located words.

Lemma 5.6 (RCA₀). *Suppose $f, g : \text{FIN}_A \rightarrow C$ are colorings such that g is strongly proximal to f . Then there is an infinite block sequence $X \subseteq \text{FIN}_{A^\star}$ such that $[X]_A$ is both f -homogeneous and g -homogeneous for color $g(\emptyset)$.*

Proof. We build a sequence $p_0 < p_1 < \dots$ of located variable words inductively.

Since g is strongly proximal to f , there is some located variable word $p_0 \in \text{FIN}_{A^\star}$ such that for each $a \in A$, $S_{p_0[a]}^0(f) = S_{p_0[a]}^0(g) = S^0(g)$. In particular, $f(p_0[a]) = g(p_0[a]) = g(\emptyset)$ for every $a \in A$.

Assume $p_0 < \dots < p_n$ are located variable words such that $[p_0, \dots, p_n]_A$ is both f -homogeneous and g -homogeneous for color $g(\emptyset)$. Let $\ell = 1 \max \text{dom } p_n$. Since g is strongly proximal to f , there is some located variable word $p_{n+1} \in \text{FIN}_{A^\star}(\ell, \emptyset)$ such that for each $a \in A$,

$$S_{p_{n+1}[a]}^\ell(f) = S_{p_{n+1}[a]}^\ell(g) = S^\ell(g)$$

Let $p \in [p_0, \dots, p_n]_A \cup \{\emptyset\}$. Note that $\max \text{dom } p < \ell$ and that $g(p) = g(\emptyset)$. Then for every $a \in A$,

$$f(p \cup p_{n+1}[a]) = S_{p_{n+1}[a]}(f)(p) = g(p) = g(\emptyset)$$

$$g(p \cup p_{n+1}[a]) = S_{p_{n+1}[a]}(g)(p) = g(p) = g(\emptyset)$$

Thus, $[p_0, \dots, p_n, p_{n+1}]_A$ is both f -homogeneous and g -homogeneous for color $g(\emptyset)$. \square

Corollary 5.7 (RCA₀). *The weak Auslander-Ellis theorem for C^{FIN_A} implies Carlson's theorem for located words.*

Proof. Immediate by Lemma 5.6. \square

5.2. Proof of the Auslander-Ellis theorem for C^{FIN_A}

The purpose of this section is to prove the Auslander-Ellis theorem for C^{FIN_A} from an iterated version of the Finite Union theorem. Together with the previous section, we will obtain a proof of Carlson's theorem for located words from the iterated Finite Union theorem. The proof follows the structure of Section 5 of Blass, Hirst and Simpson [2].

We can actually refine our statement about the Finite Union theorem and consider only bounded unions. Given some $r \in \mathbb{N}$, let

$$\text{FU}^{\leq r}(X) = \{\cup F : F \subseteq X \wedge 0 < |F| \leq n\}$$

Theorem 5.8 (Iterated Finite Union (bounded version)). *Let $X \subseteq \mathcal{P}_f(\mathbb{N})$ be an infinite block sequence. For every sequence $(f_n)_{n \in \mathbb{N}}$ of colorings $\text{FU}(X) \rightarrow C$, there is an infinite block sequence $Y \subseteq \text{FU}(X)$ such that for every n , there is some finite set $F \subseteq Y$ such that $\text{FU}^{\leq r}(Y - F)$ is f_n -homogeneous.*

We will write $\text{IFUT}^{\leq r}$ as a shorthand for the iterated Finite Union theorem for unions of at most r elements. We need a notion of weak block sequence playing the same role as the notion of block sequence, but for located words instead of located variable words.

Definition 5.9. A *weak block sequence* is a totally ordered set $X \subseteq \text{FIN}_A$. Given a weak block sequence X and some $r \in \mathbb{N}$, we let

$$[X]_A^{\leq r} = \{p_0 \cup \dots \cup p_{k-1} \in \text{FIN}_A : k \leq r \wedge p_0, \dots, p_{k-1} \in \text{FIN}_A\}$$

The following notion is an adaptation of the IP-limit to combinatorial spaces of located words.

Definition 5.10. Given $f, g : \text{FIN}_A \rightarrow C$ and an infinite weak block sequence $X \subseteq \text{FIN}_A$, we write $\text{FU}^{\leq 2}\text{-lim}_X(f) = g$ if for every $\ell \in \mathbb{N}$, there is a finite set $F \subseteq X$ such that for every $p \in [X - F]_A^{\leq 2}$, $S_p^\ell(f) = S^\ell(g)$.

Lemma 5.11 (RCA₀). *If $g = \text{FU}^{\leq 2}\text{-lim}_X(f)$ for an infinite weak block sequence $X \subseteq \text{FIN}_A$ then g is weakly proximal to f .*

Proof. Suppose $g = \text{FU}^{\leq 2}\text{-lim}_X(f)$. Given $\ell \in \mathbb{N}$, let $F_0 \subseteq X$ be such that for every $p \in [X - F_0]_A^{\leq 2}$, $S_p^\ell(f) = S^\ell(g)$. Fix $p \in X - F_0$ and let $m = 1 + \max p$. Let $F_1 \subseteq X$ be such that for every $q \in [X - F_1]_A^{\leq 2}$, $S_q^m(f) = S^m(g)$. Fix $q \in X - F_1$. In particular, $S_{p \cup q}^\ell(f) = S_p^\ell(g)$. Since $p, p \cup q \in [X - F_0]_A^{\leq 2}$, then $S_p^\ell(f) = S^\ell(g)$ and $S_{p \cup q}^\ell(f) = S^\ell(g)$. It follows that

$$S_p^\ell(f) = S^\ell(g) = S_{p \cup q}^\ell(f) = S_p^\ell(g)$$

In other words, g is weakly proximal to f . □

Lemma 5.12 (RCA₀ + IFUT^{≤2}). *For every coloring $f : \text{FIN}_A \rightarrow C$ and every infinite weak block sequence $X \subseteq \text{FIN}_A$, there is an infinite weak block sequence $Y \subseteq [X]_A$ such that $\text{FU}^{\leq 2}\text{-lim}_Y(f)$ exists.*

Proof. Let $B = \{\text{dom } p : p \in X\}$. There exists a canonical bijection $\pi : \text{FU}(B) \rightarrow [X]_A$.

For every $\ell \in \mathbb{N}$ and finite coloring $h : \text{FIN}_A(0, \ell) \rightarrow C$, let $C_h = \{F \in \text{FU}(B) : S_{\pi(F)}^\ell(f) = h\}$. Thus $\langle C_h : h \in \bigcup_\ell C^{\text{FIN}_A(0, \ell)} \rangle$ is a countable sequence of 2-colorings of $\text{FU}(B)$. By IFUT^{≤2}, there is an infinite set $D \subseteq \text{FU}(B)$ such that for every $h : \text{FIN}_A(0, \ell) \rightarrow C$, there is a finite set $G_h \subseteq D$ such that $\text{FU}^{\leq 2}(D - G_h)$ is homogeneous for C_h . Moreover, the sequence $\langle G_h : h \in \bigcup_\ell C^{\text{FIN}_A(0, \ell)} \rangle$ exists by ACA₀, which follows from RCA₀ + IFUT^{≤2}.

We claim that for every $\ell \in \mathbb{N}$, there is at most one $h : \text{FIN}_A(0, \ell) \rightarrow C$ such that $\text{FU}^{\leq 2}(D - G_h) \subseteq C_h$. Indeed, if there are $h_0, h_1 : \text{FIN}_A(0, \ell) \rightarrow C$ such that $\text{FU}^{\leq 2}(D - G_{h_i}) \subseteq C_{h_i}$ for each $i < 2$, then pick some $F \in D - (G_{h_0} \cup G_{h_1})$. Then $F \in C_{h_0} \cap C_{h_1}F$, so by definition of C_{h_i} , $S_{\pi(F)}^\ell(f) = h_i$, so $h_0 = h_1$.

For every $\ell \in \mathbb{N}$, let $h_\ell : \text{FIN}_A(0, \ell) \rightarrow C$ witness the claim. Note that $h_\ell \subseteq h_{\ell+1}$. The sequence $\langle h_\ell : \ell \in \mathbb{N} \rangle$ can be obtained computably from the sequence $\langle G_h : h \in \bigcup_\ell C^{\text{FIN}_A(0, \ell)} \rangle$. Let $g : \text{FIN}_A \rightarrow C$ be such that $S^\ell(g) = h_\ell$ for every $\ell \in \mathbb{N}$. Let $Y = \{\pi(F) : F \in D\}$. Note that $Y \subseteq [X]_A$ is a weak block sequence. By construction, $\text{FU}^{\leq 2}\text{-lim}_Y(f) = g$. □

Lemma 5.13 (RCA₀ + IFUT^{≤2}). *For every coloring $f : \text{FIN}_A \rightarrow C$ and every $g \in \overline{\text{Orb}(f)}$, there is a coloring $h \in \overline{\text{Orb}(g)}$ such that $h = \text{FU}^{\leq 2}\text{-lim}_X(f)$ for some infinite weak block sequence $X \subseteq \text{FIN}_A$.*

Proof. Fix f and g . Assume $f \neq g$, otherwise the result follows from Lemma 5.12 directly.

For every $\ell \in \mathbb{N}$, let $\mathcal{V}_\ell = C^{\text{FIN}_A(0, \ell)}$, and let $\mathcal{V} = \bigcup_\ell \mathcal{V}_\ell$. Let $T \subseteq \mathcal{V}$ be a code for the subshift $\overline{\text{Orb}(g)}$ such that T has no leaves. For each $\ell \in \mathbb{N}$, let $T_\ell = T \cap \mathcal{V}_\ell$. Let \mathcal{U}_ℓ be the open class

induced by the leaves of T_ℓ . Last, for every $p \in \text{FIN}_A(\ell, \infty)$, let $\mathcal{W}_{\ell,p} = \{h \in C^{\text{FIN}_A} : S_p^\ell(h) \in T_\ell\}$. Note that for all $\ell \in \mathbb{N}$ and $p \in \text{FIN}_A(\ell, \infty)$, $\overline{\text{Orb}(g)} \subseteq \mathcal{W}_{\ell,p}$ and $\mathcal{W}_{\ell,p}$ is open.

Define the weak block sequence $p_0 < p_1 < \dots$ inductively as follows. Let $p_0 \in \text{FIN}_A$ be arbitrary. Given p_ℓ defined, with $m_\ell = 1 + \max \text{dom } p_\ell$, since $\bigcap_{p \in \text{FIN}_A(0, m_\ell)} \mathcal{W}_{\ell,p}$ is an open cover of $\overline{\text{Orb}(g)}$, there is some bound $b \in \mathbb{N}$ such that $\mathcal{U}_b \subseteq \bigcap_{p \in \text{FIN}_A(0, m_\ell)} \mathcal{W}_{\ell,p}$. By definition of T_b , since $g \in \overline{\text{Orb}(g)}$, there is some $h \in T_b$ be such that $S^b(g) = h$. In particular, $[h] \subseteq \mathcal{U}_b$. Since $g \in \overline{\text{Orb}(f)}$, then there is some located word $p_{\ell+1} \in \text{FIN}_A(b, \infty)$ such that $S_{p_{\ell+1}}^b(f) = h$. Moreover, since $f \neq g$, the located word $p_{\ell+1}$ can be taken to be non-empty. In particular, $[S_{p_{\ell+1}}^b(f)] \subseteq \mathcal{U}_b \subseteq \bigcap_{p \in \text{FIN}_A(0, m_\ell)} \mathcal{W}_{\ell,p}$. Thus for each $\ell \in \mathbb{N}$, $S_p^\ell(f) \in T_\ell$ for all sufficiently large $p \in [p_0, p_1, \dots]_A$.

By Lemma 5.12, there is an infinite weak block sequence $X \subseteq [p_0, p_1, \dots]_A$ such that $h = \text{FU}^{\leq 2} - \lim_X(f)$ exists. Then $h \in \bigcap_\ell \mathcal{U}_\ell = \overline{\text{Orb}(Y)}$. \square

We are now ready to prove the Auslander-Ellis theorem for C^{FIN_A} from the iterated Finite Union theorem.

Proof of Theorem 5.2 over $\text{RCA}_0 + \text{IFUT}^{\leq 2}$. Let $f : \text{FIN}_A \rightarrow C$ be a coloring. By Lemma 4.6, let $g \in \overline{\text{Orb}(f)}$ be such that $\overline{\text{Orb}(g)}$ is minimal. By Lemma 5.13, there is a coloring $h \in \overline{\text{Orb}(g)}$ and an infinite weak sequence $X \subseteq \text{FIN}_A$ such that $h = \text{FU}^{\leq 2} - \lim_X(f)$. By Lemma 5.11, h is weakly proximal to f . Moreover, $\overline{\text{Orb}(h)} = \overline{\text{Orb}(h)}$ by minimality, hence h is uniformly recurrent by Lemma 4.5. \square

This gives us in particular a proof of Carlson's theorem from the iterated Finite Union theorem for unions of at most 2 elements.

Proof of Theorem 1.10 over $\text{RCA}_0 + \text{IFUT}^{\leq 2}$. Let $f : \text{FIN}_A \rightarrow C$ be a coloring. By Theorem 5.2, there is a coloring g which is uniformly recurrent, and weakly proximal to f . By Lemma 5.4, g is strongly proximal to f . By Lemma 5.6, there is an infinite block sequence $X \subseteq \text{FIN}_{A^*}$ such that $[X]_A$ is f -homogeneous. \square

Corollary 5.14. $\text{RCA}_0 \vdash \text{IFUT}^{\leq 2} \rightarrow \text{FUT}$

Proof. Immediate, since $\text{RCA}_0 + \text{IFUT}^{\leq 2}$ proves Carlson's theorem for located words (Theorem 1.10), which itself implies the Finite Union theorem. \square

It is a major open question in combinatorics whether Hindman's theorem for bounded sums implies Hindman's theorem (see Hindman, Leader and Strauss [14, Question 12]). Corollary 5.14 can be seen as a partial negative answer.

6. OPEN QUESTIONS

There are many remaining open questions in the reverse mathematics of Ramsey's theory. We mention a few of them. The first one, maybe the most important, is the following.

Question 6.1. Does (iterated) Hindman's theorem hold in ACA_0 ?

We have seen that the iterated Finite Union theorem implies Carlson's theorem for located words. However, the proof does not seem to be adaptable to replace IFUT with its non-iterated version.

Question 6.2. Does Hindman's theorem imply Carlson's theorem for (located) words?

Last, as mentioned in the introduction, Carlson proved a stronger statement about variable words, which has no known elementary proof.

Question 6.3. What are the reverse mathematics of Carlson's theorem for variable (located) words?

ACKNOWLEDGEMENT

Patey is partially supported by grant ANR "ACTC" #ANR-19-CE48-0012-01.

REFERENCES

- [1] James E. Baumgartner. A short proof of Hindman’s theorem. *Journal of Combinatorial Theory, Series A*, 17(3):384–386, 1974.
- [2] Andreas R. Blass, Jeffrey L. Hirst, and Stephen G. Simpson. Logical analysis of some theorems of combinatorics and topological dynamics. *Logic and Combinatorics, S. Simpson, ed., Contemporary Math*, 69:125–156, 1987.
- [3] Timothy J. Carlson. Some unifying principles in Ramsey theory. *Discrete Math.*, 68(2-3):117–169, 1988.
- [4] Timothy J. Carlson and Stephen G. Simpson. A dual form of Ramsey’s theorem. *Adv. in Math.*, 53(3):265–290, 1984.
- [5] Lorenzo Carlucci, Leszek Aleksander Kołodziejczyk, Francesco Lepore, and Konrad Zdanowski. New bounds on the strength of some restrictions of Hindman’s theorem. *Computability*, 9(2):139–153, 2020.
- [6] Adam R. Day. On the strength of two recurrence theorems. *J. Symb. Log.*, 81(4):1357–1374, 2016.
- [7] Pandelis Dodos and Vassilis Kanellopoulos. *Ramsey theory for product spaces*, volume 212 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2016.
- [8] Damir Dzhafarov and Carl Mummert. *Reverse Mathematics: Problems, Reductions, and Proofs*. Springer, 2022.
- [9] H. Furstenberg and Y. Katznelson. Idempotents in compact semigroups and Ramsey theory. *Israel J. Math.*, 68(3):257–270, 1989.
- [10] H. Furstenberg and B. Weiss. Topological dynamics and combinatorial number theory. *J. Analyse Math.*, 34:61–85 (1979), 1978.
- [11] R. L. Graham and B. L. Rothschild. Ramsey’s theorem for n -parameter sets. *Trans. Amer. Math. Soc.*, 159:257–292, 1971.
- [12] A. W. Hales and R. I. Jewett. Regularity and positional games. *Trans. Amer. Math. Soc.*, 106:222–229, 1963.
- [13] Neil Hindman. Finite sums from sequences within cells of a partition of N . *J. Combinatorial Theory Ser. A*, 17:1–11, 1974.
- [14] Neil Hindman, Imre Leader, and Dona Strauss. Open problems in partition regularity. volume 12, pages 571–583. 2003. Special issue on Ramsey theory.
- [15] Neil Hindman and Dona Strauss. *Algebra in the Stone-Čech compactification*. De Gruyter Textbook. Walter de Gruyter & Co., Berlin, 2012. Theory and applications, Second revised and extended edition [of MR1642231].
- [16] Denis R. Hirschfeldt. *Slicing the truth*, volume 28 of *Lecture Notes Series. Institute for Mathematical Sciences. National University of Singapore*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015. On the computable and reverse mathematics of combinatorial principles, Edited and with a foreword by Chitao Chong, Qi Feng, Theodore A. Slaman, W. Hugh Woodin and Yue Yang.
- [17] Carl G. Jockusch. Ramsey’s theorem and recursion theory. *Journal of Symbolic Logic*, 37(2):268–280, 1972.
- [18] Nikolaos Karagiannis. A combinatorial proof of an infinite version of the Hales-Jewett theorem. *J. Comb.*, 4(2):273–291, 2013.
- [19] Alexander P. Kreuzer. Minimal idempotent ultrafilters and the auslander-ellis theorem, 2013.
- [20] Lu Liu. RT_2^2 does not imply WKL_0 . *Journal of Symbolic Logic*, 77(2):609–620, 2012.
- [21] Antonio Montalbán and Richard A. Shore. Conservativity of ultrafilters over subsystems of second order arithmetic. *J. Symb. Log.*, 83(2):740–765, 2018.
- [22] David Seetapun and Theodore A. Slaman. On the strength of Ramsey’s theorem. *Notre Dame Journal of Formal Logic*, 36(4):570–582, 1995.
- [23] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Cambridge University Press, 2009.
- [24] Ernst Specker. Ramsey’s theorem does not hold in recursive set theory. In *Studies in Logic and the Foundations of Mathematics*, volume 61, pages 439–442. Elsevier, 1971.
- [25] Henry Towsner. A simple proof and some difficult examples for Hindman’s theorem. *Notre Dame J. Form. Log.*, 53(1):53–65, 2012.

E-mail address: tristanbompard2.7@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, HNP-LAMA, CENTRAL SOUTH UNIVERSITY, CITY CHANGSHA, HUNAN PROVINCE, CHINA. 410083

E-mail address: g.jiayi.liu@gmail.com

CNRS, ÉQUIPE DE LOGIQUE, UNIVERSITÉ DE PARIS, PARIS, FRANCE

E-mail address: ludovic.patey@computability.fr

URL: <http://ludovicpatey.com>