

Π_4^0 conservation of Ramsey's theorem for pairs

Quentin Le Hou  rou Ludovic Levy Patey
Keita Yokoyama

June 2023

Abstract

In this article, we prove that Ramsey's theorem for pairs and two colors is a $\forall\Pi_4^0$ conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0$, where a $\forall\Pi_4^0$ formula consists of a universal quantifier over sets followed by a Π_4^0 formula. The proof is an improvement of a result by Patey and Yokoyama [22] and a step towards the resolution of the longstanding question of the first-order part of Ramsey's theorem for pairs. For this, we introduce a new general technique for proving Π_4^0 -conservation theorems.

MSC classes Primary: 03B30, 03F30 Secondary: 05D10

1 Introduction

Among the theorems studied in Reverse Mathematics, Ramsey's theorem for pairs plays a significant role, as it escapes the structural phenomenon of the so-called Big Five. The study of its ω -models yielded many longstanding open problems, and each of them motivated the discovery of new techniques in Computability Theory [24, 3, 18, 19]. See Hirschfeldt [10] for a gentle introduction to the Reverse Mathematics of Ramsey's theorem.

Given a set $X \subseteq \mathbb{N}$, we write $[X]^n$ for the set of unordered n -tuples over X . A set $H \subseteq \mathbb{N}$ is *homogeneous* for a coloring $f : [\mathbb{N}]^n \rightarrow k$ if f is constant over $[H]^n$.

Statement (Ramsey's theorem for n -tuples and k colors). RT_k^n : *For every coloring $f : [\mathbb{N}]^n \rightarrow k$, there is an infinite f -homogeneous set.*

From a proof-theoretic perspective, Ramsey's theorem for pairs also raised many challenging open questions, among which the characterization of its first-order part. The *first-order part* of a theorem T of second-order arithmetic is the set of its first-order consequences, that is, the sentences in the language \mathcal{L}_{PA} which are provable by T . RT_2^2 is known to imply the collection principle for Σ_2^0 formulas ($\text{B}\Sigma_2^0$) over RCA_0 (see Hirst [11]) and to be Π_1^1 conservative over $\text{RCA}_0 + \text{I}\Sigma_2^0$ (see Cholak, Jockusch and Slaman [3]). On the other hand, Chong, Slaman and Yang [5] proved that RT_2^2 does not imply $\text{I}\Sigma_2^0$ over RCA_0 . The following question is arguably the most important open question of the reverse mathematics of Ramsey's theorem:

Question 1.1. *Is $\text{RT}_2^2 \Pi_1^1$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$?*

The reader can refer to [17] for a history of the quest for the first-order part of Ramsey's theorem for pairs. Thanks to an isomorphism theorem for weak König's lemma (WKL_0), Fiori-Carones, Kołodziejczyk, Wong and Yokoyama [6] proved that in order to prove Π_1^1 conservation of RT_2^2 over $\text{RCA}_0 + \text{B}\Sigma_2^0$, it is sufficient to study only a fixed level in the arithmetic hierarchy.

Theorem 1.2 (Fiori-Carones et al. [6]). *RT_2^2 is Π_1^1 conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$ iff it is $\forall\Pi_5^0$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.*

Here, a formula is $\forall\Pi_n^0$ if it is of the form $\forall X \varphi(X)$ where φ is a Π_n^0 formula. Patey and Yokoyama [22] proved the following theorem:

Theorem 1.3 (Patey and Yokoyama [22]). *RT_2^2 is $\forall\Pi_3^0$ conservative over RCA_0 .*

Note that the $\forall\Pi_3^0$ consequences of RCA_0 and $\text{RCA}_0 + \text{B}\Sigma_2^0$ coincide, by a parameterized version of the Parsons, Paris and Friedman conservation theorem (see [12] or [2]). In this article, we make one further step towards the characterization of the first-order part of Ramsey's theorem for pairs, by proving the following theorem:

Theorem 1.4 (Main theorem). *$\text{WKL}_0 + \text{RT}_2^2$ is $\forall\Pi_4^0$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.*

The proof of the main theorem follows the structure of Patey and Yokoyama [22], using an indicator for $\forall\Pi_4^0$ conservation defined by Yokoyama [27, Section 4]. It is based on a variant of the notion of α -largeness, which provides a new general technique for proving $\forall\Pi_4^0$ -conservation theorems.

1.1 Quantitative largeness and Π_3^0 sentences

A family of finite sets of natural numbers $\mathcal{L} \subseteq [\mathbb{N}]^{<\mathbb{N}}$ is said to be a *largeness notion* if any infinite set has a finite subset in \mathcal{L} and \mathcal{L} is closed under supersets. Ketonen and Solovay [13] defined a quantitative notion of largeness, called ω^n -largeness (see Definition 2.1), to measure the size of sets necessary to satisfy Σ_1^0 formulas which WKL_0 -provably hold over infinite sets. More precisely, the following theorem holds:

Theorem 1.5 (Generalized Parsons theorem for WKL_0 [22]). *Let $\psi(x, y, F)$ be a Σ_0 formula with exactly the displayed free variables. Assume that*

$$\text{WKL}_0 \vdash \forall x \forall X (X \text{ is infinite} \rightarrow \exists F \subseteq_{\text{fin}} X \exists y \psi(x, y, F))$$

Then there exists some $n \in \omega$ such that $|\Sigma_1^0|$ proves $\forall x \forall Z \subseteq_{\text{fin}} (x, \infty)$

$$Z \text{ is } \omega^n\text{-large} \rightarrow \exists F \subseteq Z \exists y < \max Z \psi(x, y, F)$$

The generalized Parsons theorem for WKL_0 plays a key role in conservation results, as it provides quantitative finitary counterparts to infinitary theorems. See Patey and Yokoyama [22] for a proof of Theorem 1.5.

For the purpose of $\forall\Pi_4^0$ conservation over $\text{WKL}_0 + \text{B}\Sigma_2^0$, we shall define a new quantitative notion of largeness, called ω^n -largeness(T) (see Definition 2.5), where T is any fixed Π_3^0 sentence, and prove the following generalized Parsons theorem:

Theorem 1.6 (Generalized Parsons theorem for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$). *Let $\psi(x, y, F)$ be a Σ_0 formula with exactly the displayed free variables. Assume that*

$$\text{WKL}_0 + \text{B}\Sigma_2^0 + T \vdash \forall x \forall X (X \text{ is infinite} \rightarrow \exists F \subseteq_{\text{fin}} X \exists y \psi(x, y, F))$$

Then there exists some $n \in \omega$ such that $\text{I}\Sigma_1^0$ proves $\forall x \forall Z \subseteq_{\text{fin}} (x, \infty)$

$$Z \text{ is } \omega^n\text{-large}(T) \text{ and exp-sparse} \rightarrow \exists F \subseteq Z \exists y < \max Z \psi(x, y, F)$$

In Section 3, we shall develop a framework for $\forall\Pi_4^0$ conservation over $\text{B}\Sigma_2^0$, and will in particular relate through Theorem 3.12 the notion of $\forall\Pi_4^0$ conservation to provability over $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$, for a Π_3^0 sentence T .

1.2 Prerequisites

Along the paper, we shall resort to some well-known conservation theorems without explicitly stating them.

Statement (Weak König's lemma). **WKL**: *Every infinite binary tree admits an infinite path.*

To follow the standard reverse mathematical practice, we shall write WKL_0 for the theory $\text{RCA}_0 + \text{WKL}$. Weak König's lemma plays a central role in Reverse Mathematics, as it captures compactness arguments.

Theorem 1.7 (Hájek [8]). $\text{WKL}_0 + \text{B}\Sigma_2^0$ is a Π_1^1 -conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0$.

Ramsey's theorem for pairs admits a decomposition into two combinatorially simpler statements, namely, the Ascending Descending Sequence principle and the Erdős-Moser theorem.

Statement (Ascending Descending Sequence). **ADS**: *Every infinite linear order admits an infinite ascending or descending sequence.*

Given a coloring $f : [\mathbb{N}]^2 \rightarrow 2$, a set $H \subseteq \mathbb{N}$ is f -transitive if for every $\{x, y, z\} \in [H]^3$ with $x < y < z$ and every $i < 2$, if $f(x, y) = f(y, z) = i$, then $f(x, z) = i$.

Statement (Erdős-Moser theorem). **EM**: *For every infinite coloring $f : [\mathbb{N}]^2 \rightarrow 2$, there is an infinite f -transitive subset.*

Ramsey's theorem for pairs and two colors is equivalent to $\text{ADS} \wedge \text{EM}$ (see Bovykin and Weiermann [1]). Chong, Slaman and Yang [4, Corollary 4.4] proved the following theorem, among other Π_1^1 conservations of combinatorial statements:

Theorem 1.8 (Chong, Slaman and Yang [4]). *ADS is a Π_1^1 conservative extension of $\text{RCA}_0 + \text{BS}_2^0$.*

The following theorem was proven by Patey and Yokoyama [22] and is a generalization of an amalgamation theorem for Π_1^1 conservation by Yokoyama [25].

Theorem 1.9 (Amalgamation [22]). *Fix $n \geq 1$. Let T be a theory extending IS_1^0 which consists of sentences of the form $\forall X \exists Y \theta(X, Y)$ where θ is Π_{n+2}^0 , and let Γ_1 and Γ_2 be sentences of the same form as T . If $T + \Gamma_i$ is a $\forall \Pi_{n+2}^0$ -conservative extension of T for $i = 1, 2$, then, $T + \Gamma_1 + \Gamma_2$ is a $\forall \Pi_{n+2}^0$ -conservative extension of T .*

In particular, putting Theorem 1.8 and Theorem 1.9 together, it is sufficient to prove that EM is a $\forall \Pi_4^0$ conservative extension of $\text{RCA}_0 + \text{BS}_2^0$ to prove our main theorem.

Every Σ_1^0 formula $\varphi(G)$ can be expressed in normal form as $\exists k \psi(G \upharpoonright_k)$, where $\psi(\sigma)$ is a monotonous Σ_1^0 formula, that is, if $\psi(\sigma)$ and $\sigma \preceq \tau$, then $\psi(\tau)$ holds. All over this paper, we shall use the following standard compactness argument:

Lemma 1.10 (Folklore). *$\text{WKL}_0 + \text{BS}_2^0$ proves that for every Σ_1^0 formula $\varphi(G) \equiv \exists k \psi(G \upharpoonright_k)$ in normal form such that $\forall A \varphi(A)$ holds, there exists some $N_1 \in \mathbb{N}$ such that for every $\sigma \in 2^{N_1}$, $\psi(\sigma)$ holds. Moreover, for every Δ_2^0 predicate A , $\varphi(A)$ holds.*

Proof. Suppose there is no such N_1 . Let T be the tree of all $\sigma \in 2^{<\mathbb{N}}$ such that $\neg \psi(\sigma)$. By monotonicity of ψ , T is a tree, and by assumption, it is infinite. By WKL_0 , there is some $A \in [T]$. Then for every k , $\neg \psi(A \upharpoonright_k)$, so $\neg \varphi(A)$, contradicting our assumption. Let N_1 witness the statement of the lemma and let A be a Δ_2^0 predicate. By BS_2^0 , A is regular, so $A \upharpoonright_{N_1}$ is coded. By choice of N_1 , $\psi(A \upharpoonright_{N_1})$ holds, so $\exists k \psi(A \upharpoonright_k)$ holds, hence $\varphi(A)$ holds. \square

1.3 Structure of this paper

In Section 2, we define two quantitative notions of largeness. The first one, ω^n -largeness, was defined by Ketonen and Solovay [13] and admits a generalized Parsons theorem for provability over WKL_0 . The second one, ω^n -largeness(T), is new, and plays the same role as ω^n -largeness, but for provability over $\text{WKL}_0 + \text{BS}_2^0 + T$, where T is any fixed Π_3^0 sentence.

Then, in Section 3, we develop a framework for $\forall \Pi_4^0$ conservation of RT-like theorems. For this, given a RT-like theorem Γ , we first define in Section 3.1 a notion of n -density(Γ), which informally asserts the existence of sufficiently large finite sets, such that n consecutive applications of Γ yield a large set.

Following techniques initially defined by Kirby and Paris [14], we prove that RT_2^2 is a $\forall\Pi_4^0$ conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0 +$ “there exists n -dense sets for every $n \in \omega$ ” (see Theorem 3.9). $\forall\Pi_4^0$ conservation over $\text{RCA}_0 + \text{B}\Sigma_2^0$ is then reduced to whether $\text{RCA}_0 + \text{B}\Sigma_2^0$ proves the existence of these n -dense(Γ) sets, for every $n \in \omega$. It is however difficult to directly prove the existence of n -dense(Γ) sets, and we therefore resort in Section 3.2 to the quantitative notion of largeness, ω^n -largeness(T), to prove this existence step by step, by handling one application of Γ at a time.

In Section 4, we introduce the grouping principle, originally defined by Patey and Yokoyama [22], and prove that it is a Π_1^1 conservative extension of $\text{RCA}_0 + \text{B}\Sigma_2^0$. This principle was shown very useful to construct ω^{n+1} -large sets from sequences of ω^n -large sets after an application of RT_2^2 in [22]. Using the generalized Parsons theorem for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$, we derive a finitary version of the grouping principle in Section 4.1. This will be the actual version which will be used in the last section. In Section 5, we apply this framework to prove that EM , ADS and RT_2^2 are $\forall\Pi_4^0$ conservative extensions of $\text{RCA}_0 + \text{B}\Sigma_2^0$.

In Section 6, we give two proofs that $\text{RCA}_0 + \text{B}\Sigma_2^0$ is a conservative extension of RCA_0 for a restricted class of $\forall\Pi_4^0$ formulas.

In Section 7, we prove the tightness of the upper bound of largeness for the pigeonhole principle. Last, in Section 8, we open the discussion to proof sizes and state some related open questions.

2 Largeness

In this section, we define some quantitative notions of largeness both for WKL_0 and for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$, where T is any Π_3^0 sentence. The first notion of largeness is originally due to Ketonen and Solovay [13], and is already well-understood. The second notion is new, so we conduct a systematic study of it, by proving some combinatorial lemmas, and a generalized Parsons theorem, which will often be used in the remainder of this article.

2.1 Largeness for WKL_0

Ketonen and Solovay [13] defined a quantitative notion of largeness, called α -largeness for every $\alpha < \epsilon_0$, and proved a partition theorem for this notion. We give here an equivalent inductive definition in the restricted case where α is of the form $\omega^n \cdot k$ with $n, k \in \omega$. This definition will serve as a basis for the quantitative notion of largeness for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$.

Definition 2.1 (Largeness for WKL_0). *A set $X \subseteq_{\text{fin}} \mathbb{N}$ is*

- ω^0 -large if $X \neq \emptyset$.
- $\omega^{(n+1)}$ -large if $X \setminus \min X$ is $(\omega^n \cdot \min X)$ -large
- $\omega^n \cdot k$ -large if there are k ω^n -large subsets of X

$$X_0 < X_1 < \dots < X_{k-1}$$

where $A < B$ means that for all $a \in A$ and $b \in B$, $a < b$.

Following the convention of Kołodziejczyk and Yokoyama [16], we always assume that $\min X \geq 3$ to avoid some degenerate behavior when $\min X$ is too small. The equivalence between Definition 2.1 and the original one is a consequence of [9, Theorem II.3.21]. This notion of largeness enjoys many desirable properties: RCA_0 proves that ω^n -largeness is a largeness notion for every $n \in \omega$. Moreover, it satisfies the generalized Parsons theorem for RCA_0 (Theorem 1.5).

Remark 2.2. *The definition above is also a notion of largeness for $\text{WKL}_0 + \text{B}\Sigma_2^0$, and Theorem 1.5 also holds if one replaces provability over WKL_0 by provability over $\text{WKL}_0 + \text{B}\Sigma_2^0$, as $\text{WKL}_0 + \text{B}\Sigma_2^0$ is a $\forall\Pi_3^0$ conservative extension of WKL_0 (see [12] or [2]).*

The main result of Ketonen and Solovay is the following partition theorem:

Theorem 2.3 (Ketonen and Solovay [13]). *Fix $k \in \omega$. RCA_0 proves that for every ω^{k+4} -large set $X \subseteq_{\text{fin}} \mathbb{N}$ and for every coloring $f : [X]^2 \rightarrow k$, there is an ω -large f -homogeneous subset $Y \subseteq X$.*

The theorem of Ketonen and Solovay was later generalized by Kołodziejczyk and Yokoyama [16, Theorem 1.6], who proved that if X is $\omega^{300^{k-1}n}$ -large, then for every coloring $f : [X]^2 \rightarrow k$, there is an ω^n -large f -homogeneous subset.

2.2 Largeness for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$

We now adapt the notion of ω^n -largeness, to obtain a quantitative notion of largeness which will play the same role, but for provability over $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$ where T is a Π_3^0 sentence.

Such a development might seem a bit ad-hoc and unrelated to $\forall\Pi_4^0$ conservation at first sight, but it will make full sense in Section 3.2, where it will be shown to be very useful for proving the existence of density notions. The reason of the use of a Π_3^0 sentence comes from the standard proof that a statement Γ is $\forall\Pi_4^0$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$. Assume that some sentence of the form $\forall A \forall a \exists x \forall y \exists z \zeta(A \upharpoonright_z, a, x, y, z)$ is not provable over $\text{RCA}_0 + \text{B}\Sigma_2^0$, where ζ is a Σ_0^0 -formula. Then there exists some model $\mathcal{M} = (M, S)$ with some set $A \in S$ and some integer a such that $\mathcal{M} \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \forall x \exists y \forall z \neg \zeta(A \upharpoonright_z, a, x, y, z)$. Working in a language enriched with constants A and a , and letting $T \equiv \forall x \exists y \forall z \neg \zeta(A \upharpoonright_z, a, x, y, z)$, we want to create a model of $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ which furthermore satisfies Γ . We therefore naturally end up reasoning over $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ for a Π_3^0 sentence T .

For the remainder of this article, and unless specified, the arithmetic hierarchy is extended to allow the use of the constants A and a inside formulas. Let $\theta(x, y, z)$ be the Σ_0^0 -formula $\neg \zeta(A \upharpoonright_z, a, x, y, z)$ and let $T \equiv \forall x \exists y \forall z \theta(x, y, z)$.

Definition 2.4. *Two finite sets $X < Y$ are T -apart if*

$$\forall x < \max X \exists y < \min Y \forall z < \max Y \theta(x, y, z)$$

Note that T -apartness is a transitive relation. Moreover, if $X < Y$ are T -apart and $X_0 \subseteq X$ and $Y_0 \subseteq Y$, then X_0, Y_0 are T -apart.

Definition 2.5 (Largeness for $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$). *A set $X \subseteq_{\text{fin}} \mathbb{N}$ is*

- ω^0 -large(T) if $X \neq \emptyset$
- $\omega^{(n+1)}$ -large(T) if $X \setminus \min X$ is $(\omega^n \cdot \min X)$ -large(T)
- $\omega^n \cdot k$ -large(T) if there are k pairwise T -apart ω^n -large(T) subsets of X

$$X_0 < X_1 < \dots < X_{k-1}$$

As for ω^n -largeness, we will only consider sets X such that $\min X \geq 3$. In addition, we will also require that $\min X \geq a$ to ensure that the constant a will always be in the new model obtained by building a proper cut (see Proposition 2.16).

Note that if we take $\theta(x, y, z)$ to be the \top formula, then $\omega^n \cdot k$ -largeness(T) is exactly $\omega^n \cdot k$ -largeness. We first prove some basic closure properties for ω^n -largeness(T). A stronger closure property will be proven in Section 2.4.

Lemma 2.6. $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ proves that for every $b \in \mathbb{N}$, if ω^b -largeness(T) is a largeness notion, then

1. For every $c \in \mathbb{N}$, $\omega^b \cdot c$ -largeness(T) is a largeness notion ;
2. ω^{b+1} -largeness(T) is a largeness notion.

Proof. Suppose that ω^b -largeness(T) is a largeness notion.

Fix an infinite set X . We prove by internal $\Sigma_1^0(X)$ induction over $c \geq 1$, that X contains an $\omega^b \cdot c$ -large(T) subset. The case $c = 1$ follows from the fact that ω^b -largeness(T) is a largeness notion. Assume the case k holds. Let $F \subseteq X$ be an $\omega^b \cdot k$ -large(T) set. Since T holds, then by $\text{B}\Sigma_2^0$, there is some $d \in X$ such that

$$\forall x < \max F \exists y < d \forall z \theta(x, y, z)$$

Since $X \setminus [0, d]$ is infinite and ω^b -largeness(T) is a largeness notion, there is an ω^b -large(T) subset $G \subseteq X \setminus [0, d]$. Then $F \cup G$ is an $\omega^b \cdot (c + 1)$ -large(T) subset of X . Since every infinite set X contains a $\omega^b \cdot c$ -large(T) subset, then $\omega^b \cdot c$ -largeness(T) is a largeness notion.

Suppose now that for every $c \geq 1$, $\omega^b \cdot c$ -largeness(T) is a largeness notion. In particular, for every infinite set X , there is an $\omega^b \cdot (\min X)$ -large(T) set $F \subseteq X \setminus \{\min X\}$, hence $\{\min X\} \cup F$ is an ω^{b+1} -large(T) subset of X . Thus, ω^{b+1} -largeness(T) is a largeness notion. \square

Proposition 2.7. For every $n \in \omega$, $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ proves that for every $c \geq 1$, $\omega^n \cdot c$ -largeness(T) is a largeness notion.

Proof. We prove by external induction over n that ω^n -largeness(T) is a largeness notion. Then statement of the proposition then follows from Lemma 2.6. For the base case, since every non-empty finite set X is ω^0 -large(T), then ω^0 -largeness(T) is a largeness notion. The induction case is the second item of Lemma 2.6. \square

A set X is *exp-sparse* if for every $x < y \in X$, $4^x < y$.

Corollary 2.8. *For every $n \in \omega$, $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ proves that for every $c \geq 1$, exp-sparse $\omega^n \cdot c$ -largeness(T) is a largeness notion.*

Proof. Fix $n \in \omega$ and $c \geq 1$. Let X be an infinite set. Then, there is an infinite exp-sparse subset $Y \subseteq X$. By Proposition 2.7, there is an $\omega^n \cdot c$ -large(T) subset $F \subseteq Y$. In particular, F is an exp-sparse $\omega^n \cdot c$ -large(T) subset of X . \square

2.3 Largeness(T) combinatorics

The following lemma is inspired from Kołodziejczyk and Yokoyama [16, Lemma 2.2], but required some variations of the combinatorics to handle the constraints based on T .

Lemma 2.9. *$|\Sigma_1^0$ proves that for all $b \in \mathbb{N}$, every ω^{2b} -large(T) exp-sparse set X and every coloring $f : X \rightarrow \min X$, there is an ω^b -large(T) f -homogeneous subset $Y \subseteq X$.*

Proof. By induction over b .

Case $b = 0$. For every ω^0 -large(T) set X and every $f : X \rightarrow \min X$, every 1-element subset of X is f -homogeneous and ω^0 -large(T).

Case $b > 0$. Fix an ω^{2b} -large(T) exp-sparse set X and $f : X \rightarrow \min X$. Since X is ω^{2b} -large(T), there are $\min X$ pairwise T -apart ω^{2b-1} -large(T) subsets $X_0 < \dots < X_{\min X - 1}$ of $X \setminus \min X$. Suppose X_0 is not f -homogeneous, otherwise we are done. Let $t < \min X$ be maximal such that $f[\bigcup_{i < t} X_i] \supseteq f[X_t]$. Note that $t \geq 1$ exists since otherwise,

$$2 \leq |f[X_0]| < |f[X_0 \cup X_1]| < \dots < |f[X_0 \cup \dots \cup X_{\min X - 1}]|$$

and therefore $|f[X]| \geq \min X + 1$, contradicting the assumption that $f : X \rightarrow \min X$.

Since X_t is ω^{2b-1} -large(T), there are $\min X_t$ pairwise T -apart ω^{2b-2} -large(T) subsets $Y_0 < \dots < Y_{\min X_t - 1}$ of $X_t \setminus \min X_t$.

By induction hypothesis, for every $j < \min X_t$, there is a ω^{b-1} -large(T) subset $Z_j \subseteq Y_j$ which is f -homogeneous for some color $c_j < \min X$. Since X is exp-sparse, $4^{\max X_{t-1}} < \min X_t$ so $(\max X_{t-1})^2 < \min X_t$, thus $\min X \times \max X_{t-1} < \min X_t$. Thus, by the finite pigeonhole principle, there is a subset $J \subseteq \{0, \dots, \min X_t - 1\}$ of size at least $\max X_{t-1}$ and some color $c < \min X$ such that for every $j \in J$, $c_j = c$. By choice of t , $f[\bigcup_{i < t} X_i] \supseteq f[X_t]$, so there is some $x \in \bigcup_{i < t} X_i$ with $f(x) = c$. Thus, the set $\{x\} \cup \bigcup_{j \in J} Z_j$ is an ω^b -large(T) f -homogeneous subset of X . \square

Lemma 2.10. *Let $n, m \in \omega$. $\text{I}\Sigma_1^0$ proves that for every ω^{n+m+1} -large(T) set X , there are ω^n -large(T) pairwise T -apart subsets $X_0 < \dots < X_{k-1}$ of X such that $\{\min X_i : i < k\}$ is ω^m -large (in the regular sense of largeness)*

Proof. By external induction over m , we prove the following statement that directly imply the lemma (assuming that $\min X \geq 3$): for all T -apart pairs $Y_0 < Y_1$ of ω^{n+m} -large(T) sets, there are ω^n -large(T) subsets $X_1 < \dots < X_{k-1}$ of Y_1 such that, letting $X_0 = Y_0$, $\{\min X_i : i < k\}$ is ω^m -large and X_0, \dots, X_{k-1} are pairwise T -apart.

Case $m = 0$. The result is clear, as every non-empty set with an element greater than 3 is ω^0 -large(T).

Case $m > 0$. Let $Y_0 < Y_1$ be ω^{n+m} -large(T) and T -apart, let $Z_0 < \dots < Z_{\min Y_1 - 1}$ be ω^{n+m-1} -large(T) pairwise T -apart subsets of Y_1 . Since Y_0 is ω^{n+m} -large(T), then it is ω -large and therefore $\min Y_1 > \max Y_0 \geq 2 \times (\min Y_0)$.

We can then apply the inductive hypothesis on the pairs

$$(Z_0, Z_1), \dots, (Z_{2(\min Y_0) - 2}, Z_{2(\min Y_0) - 1})$$

to get for every $j < \min Y_0$, families of pairwise T -apart ω^n -large(T) subsets $Z_{2j} = X_{j,0} < \dots < X_{j,k_j}$ of $Z_{2j} \cup Z_{2j+1}$ such that $\{\min X_{j,i} : i < k_j\}$ is ω^{m-1} -large.

Consider the family, $Y_0 < X_{0,0} < X_{0,1} < \dots < X_{0,k_0} < X_{1,0} < X_{1,1} < \dots < X_{\min Y_0 - 1, k_{\min Y_0 - 1}}$. Every block of this family is ω^n -large(T). Since $X_{0,0} \subseteq Y_1$, then Y_0 and $X_{0,0}$ are T -apart. Moreover, for all $j < \min Y_0 - 1$, since $X_{j,k_j} \subseteq Z_{2j+1}$ and $X_{j+1,0} = Z_{2j+2}$, X_{j,k_j} and $X_{j+1,0}$ are T -apart. So by denoting W_i the i -th element of the family and by k the cardinality of the family, $\{\min W_i | i < k\}$ is ω^m -large and the W_i 's are pairwise T -apart. This completes the proof. \square

2.4 Closure of largeness(T)

Given an ordinal $\alpha \leq \epsilon_0$, we let $\text{WF}(\alpha)$ be the $\forall\Pi_3^0$ statement of the well-foundedness of α . It is well-known that, over RCA_0 , $\text{WF}(\alpha)$ is equivalent to the fact that every infinite set admits an α -large subset (see [16, Lemma 3.2]). This motivates the following definition:

Statement. $\text{WF}_T(\omega^b)$: *Every infinite set admits an ω^b -large(T) subset. In other words, ω^b -largeness(T) is a largeness notion.*

For every $n \in \omega$, RCA_0 proves that $\text{WF}(\omega^n)$ holds, but RCA_0 does not prove $\text{WF}(\omega^\omega)$. Thus, given a model $\mathcal{M} = (M, S)$, the set $\text{WF}(\omega^\mathcal{M}) = \{b \in M : \mathcal{M} \models \text{WF}(\omega^b)\}$ is a cut of \mathcal{M} which is proper iff $\mathcal{M} \not\models \text{WF}(\omega^\omega)$. RCA_0 proves that $\text{WF}(\mathcal{M})$ is an additive cut, that is, for every $b \in \mathbb{N}$, $\text{WF}(\omega^b)$ implies $\text{WF}(\omega^{2b})$ (see [16, Lemma 3.2]). We prove the same property for $\text{WF}_T(\omega^\mathcal{M}) = \{b \in M : \mathcal{M} \models \text{WF}_T(\omega^b)\}$.

Lemma 2.11 ($\text{RCA}_0 + \text{B}\Sigma_2^0 + T$). *Let $a, b \in \mathbb{N}$, $k \geq 0$ and $X_0 < X_1 < \dots < X_{k-1}$ be pairwise T -apart ω^a -large(T) sets such that $\{\max X_s : s < k\}$ is ω^{b+1} -large(T). Then $\{\max X_0\} \cup \bigcup_{1 \leq s < k} X_s$ is ω^{a+b} -large(T).*

Proof. By induction over b . If $b = 0$, then since any ω^1 -large(T) set has cardinality at least 2, $k \geq 1$, and by assumption, X_1 is ω^a -large(T), so $\{\max X_0\} \cup \bigcup_{1 \leq s < k} X_s$ is ω^{a+b} -large(T).

Let $b > 0$. Let $Z_0 < Z_1 < \dots < Z_{\ell-1}$ be pairwise T -apart ω^b -large(T) sets such that $\{\ell\} \cup Z_0 \cup \dots \cup Z_{\ell-1}$ is an ω^{b+1} -large(T) subset of $\{\max X_s : s < k\}$. By induction hypothesis, for every $t < \ell$, the set $W_t = \{\min Z_t\} \cup \bigcup\{X_s : \max X_s \in Z_t \setminus \min Z_t\}$ is ω^{a+b-1} -large(T). Note that $\min Z_t = \min W_t$ and $\max W_t = \max Z_t$. Given $i < j < \ell$, since Z_i and Z_j are T -apart, and since $\max Z_i = \max W_i$, $\min Z_j = \min W_j$ and $\max W_j = \max Z_j$, then W_i and W_j are T -apart. Last, since $\max X_0 \leq \ell$, then $\{\max X_0\} \cup W_0 \cup \dots \cup W_{\ell-1}$ is an ω^{a+b} -large(T) subset of $\{\max X_0\} \cup \bigcup_{1 \leq s < k} X_s$. \square

Proposition 2.12. $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ proves that for every $a \in \mathbb{N}$, if ω^a -largeness(T) is a largeness notion, then so is ω^{2a} -largeness(T).

Proof. Suppose ω^a -largeness(T) is a largeness notion. Fix an infinite set X . We build greedily an infinite subsequence $X_0 < X_1 < \dots$ of pairwise T -apart ω^a -large(T) sets. Suppose for the contradiction that there is some $k \in \mathbb{N}$ such that X_{k+1} is not found.

Since T holds, then by $\text{B}\Sigma_2^0$, there is some $b \in X$ such that

$$\forall x < \max X_k \exists y < b \forall z \theta(x, y, z)$$

Since $X \setminus [0, b]$ is infinite and ω^n -largeness(T) is a largeness notion, there is an ω^n -large(T) subset $X_{k+1} \subseteq X \setminus [0, b]$. Then X_0, \dots, X_{k+1} are pairwise T -apart, contradiction.

By Lemma 2.6, ω^{a+1} -largeness(T) is a largeness notion. Since $Y = \{\max Y_s : s \in \mathbb{N}\}$ is an infinite set, then there is an ω^{a+1} -large(T) subset F . By Lemma 2.11, $\bigcup\{X_s : \max X_s \in F\}$ is an ω^{2a} -large(T) subset of X . \square

2.5 A generalized Parsons theorem for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$

We now turn to the proof of a generalized Parsons theorem for this notion of largeness. The proof is an elaboration of the original construction by Kirby and Paris [14]. It is based on the construction of a semi-regular cut which satisfies some extra properties to make it a model of $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$. We start with a few definitions:

Definition 2.13 (Cut). *A cut in a model M of first order arithmetic is a nonempty subset $I \subseteq M$ which is closed by successor, and is an initial segment of M , that is, if $a \in I$ and $b \leq a$, then $b \in I$.*

Definition 2.14 (Semi-regular cut [14]). *A cut $I \subseteq M$ is said to be semi-regular if for every M -finite set $E \subseteq M$ such that $|E| \in I$, $E \cap I$ is bounded in I .*

The following proposition shows that semi-regular cuts are exactly those coding a model of WKL_0 :

Proposition 2.15 (See Scott [23] and Kirby and Paris [14, Proposition 1]). *Let $I \subseteq M$ be a cut and let $\text{Cod}(M/I) = \{S \cap I : S \text{ is } M\text{-finite}\}$. Then I is semi-regular if and only if $(I, \text{Cod}(M/I)) \models \text{WKL}_0$.*

Proposition 2.16. *Given a countable non-standard model $\mathcal{M} = (M, S, a^\mathcal{M}, A^\mathcal{M})$ of IS_1^0 and an M -finite set $Z \subseteq M$ which is ω^{2^c} -large(T) and exp-sparse for some $c \in M \setminus \omega$, there is an initial segment I of M such that $(I, \text{Cod}(M/I), a^\mathcal{M}, A^\mathcal{M} \cap I) \models \text{WKL}_0 + \text{BS}_2^0 + T$ and $I \cap Z$ is infinite in I .*

Proof. Let $(E_i)_{i \in \omega}$ be an enumeration containing all the M -finite sets infinitely many times each and let $(f_i)_{i \in \omega}$ be an enumeration containing infinitely many times all the M -finite functions from $\{0, \dots, \max Z\}$ to a k where $k < \max Z$.

We will build a decreasing sequence of sets $Z = Z_0 \supseteq Z_1 \supseteq \dots$ such that Z_i will be $\omega^{2^{c-i}}$ -large(T) (and still exp-sparse).

At stages of the form $s = 4i$, let Z_{4i} be given. If $\min Z_{4i} \leq |E_i|$, then keep $Z_{4i+1} = Z_{4i}$. If $\min Z_{4i} > |E_i|$, then, letting $Z_{4i}^0 < \dots < Z_{4i}^{\min Z_{4i}-1}$ be $\omega^{2^{c-4i-1}}$ -large(T) such that $\bigsqcup_{j < \min Z_{4i}} Z_{4i}^j \subseteq Z_{4i}$, by the finite pigeonhole principle, there exists a $j < \min Z_{4i}$ such that $E_i \cap Z_{4i}^j = \emptyset$, in this case, take $Z_{4i+1} = Z_{4i}^j$ for such a j .

At stages of the form $s = 4i + 1$, let Z_{4i+1} be given. Let k be the range of f_i , if $\min Z_{4i+1} < k$, then keep $Z_{4i+2} = Z_{4i+1}$. If $\min Z_{4i+1} > k$ then f_i induce a coloring $\tilde{f}_i : Z_{4i+1} \rightarrow \min Z_{4i+1}$ and let Z_{4i+2} be an $\omega^{2^{c-4i-2}}$ -large(T) \tilde{f}_i -homogeneous subset of Z_{4i+1} (by Lemma 2.9).

At stages of the form $s = 4i + 2$, let Z_{4i+2} be given and put $Z_{4i+3} = Z_{4i+2} \setminus \{\min Z_{4i+2}\}$.

At stages of the form $s = 4i + 3$, let Z_{4i+3} be given. Z_{4i+3} contains at least two $\omega^{2^{c-4i-4}}$ -large(T) and T -apart subsets, $Y_0 < Y_1$. Take $Z_{4i+4} = Y_1$.

Finally, let $I = \sup\{\min Z_i \mid i \in \omega\}$. First, note that by convention, for every ω^{2^c} -large(T) set Z , $a^\mathcal{M} \leq \min Z$, hence $a^\mathcal{M} \in I$.

The stages of the form $s = 4i$ ensure that I is a semi-regular cut and therefore $(I, \text{Cod}(M/I)) \models \text{WKL}_0$.

The stages of the form $s = 4i + 2$ ensure that $Z_i \cap I$ is infinite in I for every i (and in particular $Z \cap I$ is infinite in I).

The stages of the form $s = 4i + 1$ ensure that $(I, \text{Cod}(M/I)) \models \text{RT}^1$ (and therefore BS_2^0): let $f : I \rightarrow k \in \text{Cod}(M/I)$, there exists an index $i \in \omega$ such that $f = f_i \restriction I$ (and therefore $f = f_i \restriction I$) and since $k \in I$, we can take the i to be large enough for $\min Z_{4i+1}$ to be bigger than k (Since every such function appears infinitely many times in the enumeration). By construction Z_{4i+2} is f_i -homogeneous, so $Z_{4i+2} \cap I$ is an element of $\text{Cod}(M/I)$ that is f -homogeneous and infinite in I .

The stages of the form $s = 4i + 3$ ensure that $(I, \text{Cod}(M/I), a^\mathcal{M}, A^\mathcal{M} \cap I) \models T$ as for every $k \in I$, there exists an index i such that $k < \min Z_{4i+3}$ and therefore $\forall x < k \exists y < \min Z_{4i+4} \forall z < \max Z_{4i+4} \theta(x, y, z)$ holds. Replacing $A^\mathcal{M}$ by $A^\mathcal{M} \cap I$ inside $\theta(x, y, z)$ does not change its truth value for $x, y, z \in I$, hence $(I, \text{Cod}(M/I), a^\mathcal{M}, A^\mathcal{M} \cap I) \models \forall x < k \exists y \forall z \theta(x, y, z)$. \square

Theorem 1.6 (Generalized Parsons theorem for $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$). *Let $\psi(x, y, F)$ be a Σ_0 formula with exactly the displayed free variables. Assume that*

$$\text{WKL}_0 + \text{B}\Sigma_2^0 + T \vdash \forall x \forall X (X \text{ is infinite} \rightarrow \exists F \subseteq_{\text{fin}} X \exists y \psi(x, y, F))$$

Then there exists some $n \in \omega$ such that $\text{I}\Sigma_1^0$ proves $\forall x \forall Z \subseteq_{\text{fin}} (x, \infty)$

$$Z \text{ is } \omega^n\text{-large}(T) \text{ and exp-sparse} \rightarrow \exists F \subseteq Z \exists y < \max Z \psi(x, y, F)$$

Proof. By contradiction, assume that for all $n \in \omega$, $\text{I}\Sigma_1^0$ does not prove

$$\forall x \forall Z \subseteq_{\text{fin}} (x, \infty) [Z \text{ is } \omega^n\text{-large}(T) \text{ and exp-sparse} \rightarrow \exists F \subseteq Z \exists y < \max Z \psi(x, y, F)]$$

There is a countable model $\mathcal{M} = (M, S) \models \text{I}\Sigma_1^0$, satisfying for every $n \in \omega$

$$\exists x \exists Z \subseteq_{\text{fin}} (x, \infty) \left[\begin{array}{l} Z \text{ is } \omega^{2^n}\text{-large}(T) \text{ and exp-sparse} \\ \text{and } \forall F \subseteq Z \forall y < \max Z \neg \psi(x, y, F) \end{array} \right]$$

We can assume M to be non-standard, so by overspill, there exists a $b \in M \setminus \omega$ such that

$$M \models \exists x \exists Z \subseteq_{\text{fin}} (x, \infty) \left[\begin{array}{l} Z \text{ is } \omega^{2^b}\text{-large}(T) \text{ and exp-sparse} \\ \text{and } \forall F \subseteq Z \forall y < \max Z \neg \psi(x, y, F) \end{array} \right]$$

Let $c \in M$ and $Z \subseteq_{\text{fin}} (c, \infty)$ ω^{2^b} -large(T) be such that $\mathcal{M} \models \forall F \subseteq Z \forall y < \max Z \neg \psi(c, y, F)$. By Proposition 2.16, there is an initial segment I of M such that $(I, \text{Cod}(M/I), a^{\mathcal{M}}, A^{\mathcal{M}} \cap I) \models \text{WKL}_0 + \text{B}\Sigma_2^0 + T$ and $Z \cap I$ infinite in I . Therefore,

$$(I, \text{Cod}(M/I), a^{\mathcal{M}}, A^{\mathcal{M}} \cap I) \models (Z \cap I \text{ is infinite} \wedge \forall F \subseteq_{\text{fin}} Z \cap I \forall y \neg \psi(c, y, F))$$

This contradicts our assumption that

$$\text{WKL}_0 + \text{B}\Sigma_2^0 + T \vdash \forall x \forall X (X \text{ is infinite} \rightarrow \exists F \subseteq_{\text{fin}} X \exists y \psi(x, y, F))$$

□

3 A framework for $\forall\Pi_4^0$ conservation

We now develop a framework for proving that RT-like statements are $\forall\Pi_4^0$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$. This will be divided into two sections: First, building up on the work of Kirby and Paris [14], we prove in Section 3.1 that every RT-like theorem Γ is $\forall\Pi_4^0$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$ + a first-order theory stating the existence of dense(Γ) sets. Then, in Section 3.2, we use the quantitative notion of largeness defined in Section 2 to prove the existence of dense(Γ) sets, assuming the existence of a combinatorially simpler objects.

Given a function $f : [\mathbb{N}]^n \rightarrow k$ and a finite set $G = \{x_0 < \dots < x_{\ell-1}\}$, let $f_G : [\ell]^n \rightarrow k$ be defined by $f_G(i_0, \dots, i_{n-1}) = f(x_{i_0}, \dots, x_{i_{n-1}})$.

Definition 3.1 (RT-like formula). *Given $n, k \in \omega$, an RT-like formula is a Π_2^1 -formula of the form*

$$(\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y))$$

where $\Psi(f, Y)$ is of the form $(\forall G \subseteq_{\text{fin}} Y)\Psi_0(f_G)$ with Ψ_0 a Δ_0^0 -formula.

RT-like statements were introduced and studied in the standard realm by Patey [21], formulated in terms of forbidden patterns. The idea behind the formula Ψ is that theorems from Ramsey theory are about structure and not the actual value of the domain. Thus, every solution is rewritten as a copy of the integers.

Remark 3.2. *The notion of RT-like statement is a particular case of Ramsey-like- Π_2^1 -formula, introduced by defined by Patey and Yokoyama [22]. It already encompasses Ramsey's theorem for pairs, the Ascending Descending Sequence principle and the Erdős-Moser theorem. This section could have been formulated and proven in the more general setting of Ramsey-like- Π_2^1 -formulas, with the cost of some heavier notations as in [22].*

Since an RT-like statement does not distinguish the solutions from the set of the integers, it proves a slightly stronger formula, stating that the class of solutions is dense in the partial order $([\mathbb{N}]^{\mathbb{N}}, \subseteq)$.

Lemma 3.3. *Let $\Gamma \equiv (\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y))$ be an RT-like formula. Then $\text{RCA}_0 + \Gamma$ proves*

$$(\forall f : [\mathbb{N}]^n \rightarrow k)(\forall X \text{ infinite})(\exists Y \subseteq X)(Y \text{ is infinite} \wedge \Psi(f, Y))$$

Proof. Let $f : [\mathbb{N}]^n \rightarrow k$ be a coloring and $X = \{x_0 < x_1 < \dots\}$ be an infinite set. Let $\hat{f} : [\mathbb{N}]^n \rightarrow k$ be defined by $\hat{f}(i_0, \dots, i_{n-1}) = f(x_{i_0}, \dots, x_{i_{n-1}})$. By Γ , there is an infinite set $\hat{Y} \subseteq \mathbb{N}$ such that $\Psi(\hat{f}, \hat{Y})$. Let $Y = \{x_s : s \in \hat{Y}\}$. We claim that $\Psi(f, Y)$ holds.

Say $\Psi(f, Y) \equiv (\forall G \subseteq_{\text{fin}} Y)\Psi_0(f_G)$ with Ψ_0 a Δ_0^0 -formula. Let $G \subseteq_{\text{fin}} Y$, and let $\hat{G} = \{s : x_s \in G\}$. Say $G = \{x_{s_0}, \dots, x_{s_{\ell-1}}\}$. Then

$$f_G(i_0, \dots, i_{n-1}) = f(x_{s_{i_0}}, \dots, x_{s_{i_{n-1}}}) = \hat{f}(s_{i_0}, \dots, s_{i_{n-1}}) = \hat{f}_{\hat{G}}(i_0, \dots, i_{n-1})$$

hence $f_G = \hat{f}_{\hat{G}}$. Since $\Psi(\hat{f}, \hat{Y})$ and $\hat{G} \subseteq_{\text{fin}} \hat{Y}$, then $\Psi_0(\hat{f}_{\hat{G}})$ holds, then so does $\Psi_0(f_G)$. Since it holds for every $G \subseteq_{\text{fin}} Y$, then $\Psi(f, Y)$ holds. \square

3.1 Density(T) for RT-like statements

There exists a very general way to characterize the first-order part of a second-order theory, using the notion of density, originally defined by Kirby and Paris [14]. This was later adapted by Bovykin and Weiermann [1, Theorem 1] to prove Π_2^0 conservation of combinatorial theorems, and generalized by Patey and Yokoyama [22, Theorem 3.4] for $\forall\Pi_3^0$ conservation.

Definition 3.4. Fix an RT-like statement

$$\Gamma = (\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y)).$$

We define inductively the notion of m -density(T, Γ) of a finite set $Z \subseteq \mathbb{N}$ as follows. A set Z is 0-dense(T, Γ) if it is ω -large(T) and a set Z is $(m + 1)$ -dense(T, Γ) if

- (a) for any $f : [Z]^n \rightarrow k$, there is an m -dense(T, Γ) set $Y \subseteq Z$ such that $\Psi(f, Y)$ holds, and,
- (b) for any partition $Z_0 \sqcup \dots \sqcup Z_{\ell-1} = Z$ such that $\ell \leq Z_0 < \dots < Z_{\ell-1}$, one of the Z_i 's is m -dense(T, Γ).
- (c) for any $f : Z \rightarrow \min Z$, there is an m -dense(T, Γ) set $Y \subseteq Z$ which is f -homogeneous
- (d) there is an m -dense(T, Γ) set $Y \subseteq Z$ such that

$$\forall x < \min Z \exists y < \min Y \forall z < \max Y \theta(x, y, z)$$

As for largeness(T), we require that any m -dense(T, Γ) set X satisfies $\min X \geq \max 3, a$.

Remark 3.5.

- (1) In the definition above, item (a) is an indicator for Γ , item (b) for WKL_0 , item (c) for RT^1 and item (d) for T . Strictly speaking, this notion corresponds to m -density($\Gamma, \text{WKL}_0, \text{RT}^1, T$), but for simplicity of notation, we only made explicit the varying parameters.
- (2) Note that item (b) follows from item (c). However, for clarity and explicitness of the indicators, we kept both items.
- (3) Also note that there is a hidden use of $\text{B}\Sigma_2^0$ in item (d). Indeed, one assume that every $x < \min Z$ will have $\min Y$ as a uniform bound. One could have modified this item to require that for every $x < \min Z$, there is an m -dense(T, Γ) set $Y \subseteq Z$ and some $y < \min Y$ such that $\forall z < \max Z \theta(x, y, z)$.

We now prove the core combinatorial lemma which relates the notion of density to the existence of a cut satisfying the desired properties.

Lemma 3.6. Let Γ be an RT-like statement. Given a countable nonstandard model $\mathcal{M} \models \text{I}\Sigma_1^0$ and an M -finite set $Z \subseteq M$ which is c -dense(T, Γ) for some $c \in M \setminus \omega$, then there exists an initial segment I of M such that $(I, \text{Cod}(M/I), a^{\mathcal{M}}, A^{\mathcal{M}} \cap I) \models \text{WKL}_0 + \text{B}\Sigma_2^0 + \Gamma + T$ and $I \cap Z$ is infinite in I .

Proof. Let Γ be an RT-like statement of the form

$$(\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y))$$

where $k, n \in \omega$ and Ψ in the form of Definition 3.1

Let $(E_i)_{i \in \omega}$ be an enumeration containing all the M -finite sets infinitely many times each, let $(f_i)_{i \in \omega}$ be an enumeration of all the M -finite functions from $\{0, \dots, \max Z\}^n$ to k and let $(g_i)_{i \in \omega}$ be an enumeration containing infinitely many times all the M -finite functions from $\{0, \dots, \max Z\}$ to a l where $l < \max Z$.

We will build a decreasing sequence of sets $Z = Z_0 \supseteq Z_1 \supseteq \dots$ such that Z_i will be $(c - i)$ -dense(T, Γ).

At stages of the form $s = 5i$, let Z_{5i} be given. If $\min Z_{5i} \leq |E_i|$, then keep $Z_{5i+1} = Z_{5i}$. If $\min Z_{5i} > |E_i|$, let $E_i = \{e_0, \dots, e_{l-1}\}$ where $e_0 < \dots < e_{l-1}$ and put $Z_{5i}^0 = Z_{5i} \cap [0, e_0)_{\mathbb{N}}$, $Z_{5i}^j = Z_{5i} \cap [e_{j-1}, e_j)_{\mathbb{N}}$ for $1 \leq j < l-1$ and $Z_{5i}^l = Z_{5i} \cap [e_{l-1}, \infty)_{\mathbb{N}}$. Then $Z_{5i} = Z_{5i}^0 \sqcup \dots \sqcup Z_{5i}^l$, thus one of the Z_{5i}^j for $j \leq l$ is $(c - 5i - 1)$ -dense(T, Γ). Put Z_{5i+1} to be such a Z_{5i}^j .

At stages of the form $s = 5i + 1$, let Z_{5i+1} be given. Let l be the range of g_i , if $\min Z_{5i+1} < l$, then keep $Z_{5i+2} = Z_{5i+1}$. If $\min Z_{5i+1} > l$ then g_i induce a coloring $\tilde{g}_i : Z_{5i+1} \rightarrow \min Z_{5i+1}$ and let Z_{5i+2} be an $(c - 5i - 2)$ -dense(T, Γ) \tilde{g}_i -homogeneous subset of Z_{5i+1} .

At stages of the form $s = 5i + 2$, let Z_{5i+2} be given and put $Z_{5i+3} = Z_{5i+2} \setminus \{\min Z_{5i+2}\}$.

At stages of the form $s = 5i + 3$, let Z_{5i+3} be given and let Z_{5i+4} be a $(c - 5i - 4)$ -dense(T, Γ) subset of Z_{5i+3} such that

$$\forall x < \min Z_{5i+3} \exists y < \min Z_{5i+4} \forall z < \max Z_{5i+4} \theta(x, y, z)$$

At stages if the form $s = 5i + 4$, let Z_{5i+4} and f_i be given. Let $Z_{5i+5} \subseteq Z_{5i+4}$ satisfying $\Psi(f_i, Z_{5i+5})$.

Finally, let $I = \sup\{\min Z_i | i \in \omega\}$.

The stages of the form $s = 5i$ ensure that I is a semi-regular cut and therefore $(I, \text{Cod}(M/I)) \models \text{WKL}_0$.

The stages of the form $s = 5i + 2$ ensure that $Z_i \cap I$ is infinite in I for every i (and in particular $Z \cap I$ is infinite in I).

The stages of the form $s = 5i + 1$ ensure that $(I, \text{Cod}(M/I)) \models \text{RT}^1$ (and therefore BS_2^0): let $g : I \rightarrow k \in \text{Cod}(M/I)$, there exists an index $i \in \omega$ such that $g = g_i \restriction I$ (and therefore $g = g_i \restriction I$) and since $k \in I$, we can take the i to be large enough for $\min Z_{5i+1}$ to be bigger than k (Since every such function appears infinitely many times in the enumeration). By construction Z_{5i+2} is g_i -homogeneous, so $Z_{5i+2} \cap I$ is an element of $\text{Cod}(M/I)$ that is g -homogeneous and infinite in I .

The stages of the form $s = 5i + 3$ ensure that $(I, \text{Cod}(M/I), a^{\mathcal{M}}, A^{\mathcal{M}} \cap I) \models T$ as for every $k \in I$, there exists an index i such that $k < \min Z_{5i+3}$ and therefore

$\forall x < k \exists y \forall z < \max Z_{5i+4} \theta(x, y, z)$, so $(I, \text{Cod}(M/I), a^{\mathcal{M}}, A^{\mathcal{M}} \cap I) \models \forall x < k \exists y \forall z \theta(x, y, z)$ (since $\max Z_{5i+4} > I$), so $(I, \text{Cod}(M/I)) \models T$.

The stages of the form $s = 5i + 4$ ensure that $(I, \text{Cod}(M/I)) \models \Gamma$: Let $f : [I]^n \rightarrow k \in \text{Cod}(M/I)$, there exists an index $i \in \omega$ such that $f = f_i \cap I$ (and therefore $f = f_i \upharpoonright I$). By construction, $M \models \Psi(f_i, Z_{5i+5})$ and thus $(I, \text{Cod}(M/I)) \models \Psi(f, Z_{5i+5} \cap I)$ and since $Z_{5i+5} \cap I$ is infinite in I we have $(I, \text{Cod}(M/I)) \models \Gamma$. \square

We can now define a Paris-Harrington-like principle to which our theorem will be $\forall \Pi_4^0$ conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.

Definition 3.7 (Paris-Harrington principle for density(T)). *Fix a Σ_0^0 -formula $\theta(X \upharpoonright_z, a, x, y, z)$ with exactly the displayed free variables and some $n \in \omega$. Let $n\text{-PH}_\theta(\Gamma)$ be the statement*

“For every a, b and every set A , if $\forall x \exists y \forall z \theta(A \upharpoonright_z, a, x, y, z)$ then there is an $n\text{-dense}(\forall x \exists y \forall z \theta(A \upharpoonright_z, a, x, y, z), \Gamma)$ set X such that $\min X > b$.”

The following proposition shows that this Paris-Harrington-like statement can be actually proven by Γ , with the help of compactness.

Proposition 3.8. *Fix a Σ_0^0 -formula $\theta(X \upharpoonright_z, a, x, y, z)$ with exactly the displayed free variables. Then for every $n \in \omega$, $\text{WKL}_0 + \text{B}\Sigma_2^0 + \Gamma \vdash n\text{-PH}_\theta(\Gamma)$.*

Proof. Write $T(X, t) \equiv \forall x \exists y \forall z \theta(X \upharpoonright_z, t, x, y, z)$ and $\Gamma \equiv (\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y))$.

Fix a, b and A a set and assume $T(A, a)$. By external induction on $n \in \omega$, we will prove the stronger property that “ $n\text{-density}(T(A, a), \Gamma)$ is a largeness notion”.

Case $n = 0$. By Proposition 2.7, $\text{RCA}_0 + \text{B}\Sigma_2^0 + T(A, a)$ proves that $\omega\text{-largeness}(T(A, a))$ is a largeness notion.

Case $n > 0$. Suppose the property to be true at rank $n - 1$ and fix Y an infinite set. By the standard compactness argument (see Lemma 1.10), available within $\text{WKL}_0 + \Gamma$, and by Lemma 3.3, there is a depth d_0 such that for every $f : [[0, \dots, d_0] \cap Y]^n \rightarrow k$ there exists a $Z \subseteq [0, d_0] \cap Y$ such that $\Psi(f, Z)$ and Z is $(n - 1)\text{-dense}(T(A, a), \Gamma)$.

Let $\ell = \min Y$. Consider the tree of all ℓ -partition of Y such that no elements of the partition contains a $(n - 1)\text{-dense}(T(A, a), \Gamma)$ set. By the inductive hypothesis and $\text{B}\Sigma_2^0$, this tree has no infinite branch, so, by WKL_0 , there is a depth d_1 such that for every partition $Z_0 \sqcup \dots \sqcup Z_{\ell-1} = Y \cap [0, \dots, d_1]$, one of the Z_i is $(n - 1)\text{-dense}(T(A, a), \Gamma)$.

By $\text{B}\Sigma_2^0$ and our assumption that $\forall x \exists y \forall z \theta(A \upharpoonright_z, a, x, y, z)$, there exists a bound k such that $\forall x < \min Y \exists y < k \forall z \theta(A \upharpoonright_z, a, x, y, z)$. By the inductive hypothesis, there exists some d_2 such that $Y \cap [k, k + 1, \dots, d_2]$ is $(n - 1)\text{-dense}(T(A, a), \Gamma)$.

Finally, take $d = \max\{d_0, d_1, d_2\}$, the set $Y \cap [0, \dots, d - 1]$ is $n\text{-dense}(T(A, a), \Gamma)$ by definition of d_0, d_1, d_2 . We conclude by induction. \square

We are now ready to prove the general theorem which relates Γ to its corresponding Paris-Harrington-like theory.

Theorem 3.9. $\text{WKL}_0 + \text{B}\Sigma_2^0 + \Gamma$ is $\forall\Pi_4^0$ conservative over $\text{RCA}_0 + \{n\text{-PH}_\theta(\Gamma) : n \in \omega, \theta \in \Sigma_0^0\}$

Proof. Assume $\text{RCA}_0 + \{n\text{-PH}_\theta(\Gamma) : n \in \omega, \theta \in \Sigma_0^0\} \not\models \forall X \forall t \exists x \forall y \exists z \eta(X \upharpoonright_z, t, x, y, z)$ with η a Σ_0^0 formula. By compactness, completeness and the Löwenheim-Skolem theorem, there exists some countable model

$$\mathcal{M} = (M, S) \models \text{RCA}_0 + c\text{-PH}_{\neg\eta}(\Gamma) + \{c > n : n \in \omega\} + \forall x \exists y \forall z \neg \eta(A \upharpoonright_z, a, x, y, z)$$

where $A \in S, a \in M$ and c is a new constant symbol.

$\mathcal{M} \models \forall x \exists y \forall z \neg \eta(A, a, x, y, z) + c\text{-PH}_{\neg\eta}(\Gamma)$ so there exists a c -dense($T(A, a), \Gamma$) set X , where $T(A, a) \equiv \forall x \exists y \forall z \neg \eta(A \upharpoonright_z, a, x, y, z)$.

By Lemma 3.6, there exists an initial segment I of M such that $(I, \text{Cod}(M/I)) \models \text{WKL}_0 + \text{B}\Sigma_2^0 + \Gamma + T(A \cap I, a)$. Therefore $\text{WKL}_0 + \text{B}\Sigma_2^0 + \Gamma \not\models \forall X \forall t \exists x \forall y \exists z \eta(X \upharpoonright_z, t, x, y, z)$.

The converse follows from Proposition 3.8. □

3.2 Largeness(T) for RT-like statements

Thanks to Theorem 3.9, our main theorem is reduced to proving $n\text{-PH}_\theta(\text{RT}_2^2)$ over $\text{RCA}_0 + \text{B}\Sigma_2^0$ for every $n \in \omega$ and every Σ_0^0 formula θ . Unfolding the definitions, given a set A and a , letting $T(A, a) \equiv \forall x \exists y \forall z \theta(A \upharpoonright_z, a, x, y, z)$, the goal is to prove over $\text{RCA}_0 + \text{B}\Sigma_2^0 + T(A, a)$ the existence of n -dense($T(A, a), \text{RT}_2^2$) sets. However, the notion of n -density is not easy to manipulate. Thanks to our generalized Parsons theorem for provability over $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$, one can handle one application of RT_2^2 at a time, but in return the solution has to sufficiently large in the sense of ω^n -largeness(T).

Definition 3.10. Fix $r, s \in \omega$ and an RT-like- Π_2^1 -statement $\Gamma \equiv (\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y))$. A set $Z \subseteq_{\text{fin}} \mathbb{N}$ is said to be $\omega^r \cdot s$ -large(T, Γ) if for any $f : [Z]^n \rightarrow k$, there is an $(\omega^r \cdot s)$ -large(T) set $Y \subseteq Z$ such that $\Psi(f, Y)$ holds.

The following proposition relates density to largeness:

Proposition 3.11. Let Γ be an RT-like statement and T be a Π_3^0 formula. Suppose ω^n -largeness(T, Γ) is a largeness notion provably in $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ for every $n \in \omega$. Then n -density(T, Γ) is a largeness notion provably in $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ for every $n \in \omega$.

Proof. Write $T \equiv \forall x \exists y \forall z \theta(x, y, z)$ and $\Gamma \equiv (\forall f : [\mathbb{N}]^n \rightarrow k)(\exists Y)(Y \text{ is infinite} \wedge \Psi(f, Y))$.

By Theorem 1.6, for every $n \in \omega$, since ω^n -largeness(T, Γ) is a largeness notion provably in $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$, there exists some $\ell_n \in \omega$, such that

$$\text{I}\Sigma_1^0 \vdash \forall X (X \text{ is } \omega^{\ell_n}\text{-large}(T) \text{ and exp-sparse} \rightarrow X \text{ is } \omega^n\text{-large}(T, \Gamma))$$

Consider the following inductively defined sequence: $k_0 = 1$ and $k_{n+1} = \max\{2k_n, \ell_{k_n}, k_n + 1\}$. We claim that

$$\text{IS}_1^0 \vdash \forall X (X \text{ is } \omega^{k_n}\text{-large}(T) \text{ and exp-sparse} \rightarrow X \text{ is } n\text{-dense}(T, \Gamma))$$

By Proposition 2.8, $\text{RCA}_0 + \text{BS}_2^0 + T$ proves that exp-sparse ω^k -largeness(T) is a largeness notion for every $k \in \omega$, so if the claim is valid then n -density(T, Γ) is a largeness notion provably in $\text{RCA}_0 + \text{BS}_2^0 + T$ for every $n \in \omega$.

We prove the claim by external induction on $n \in \omega$:

Case $n = 0$, every ω^1 -large(T) set is 0-dense(T, Γ)

Case $n > 0$, assume the property to be true at rank $n - 1$. Let X be ω^{k_n} -large(T) and exp-sparse. We need to check that X is n -dense(T, Γ). Since by Remark 3.5, item (b) follows from (c), then we prove (a), (c) and (d) of Definition 3.4 :

- (a) Since X is $\omega^{\ell_{k_n-1}}$ -large(T) and exp-sparse, X is $\omega^{k_{n-1}}$ -large(T, Γ), so for any $f : [X]^n \rightarrow k$, there is an $(n - 1)$ -dense(T, Γ) subset $Y \subseteq X$ such that $\Psi(f, Y)$ holds.
- (c) Since X is $\omega^{2k_{n-1}}$ -large(T) and exp-sparse, by Lemma 2.9, for every coloring $f : X \rightarrow \min X$, there is an $\omega^{k_{n-1}}$ -large(T) subset $Y \subseteq X$ which is f -homogeneous. By induction hypothesis, Y is $(n - 1)$ -dense(T, Γ).
- (d) Since X is $\omega^{k_{n-1}+1}$ -large(T), there exists $X_0 < \dots < X_{\min X - 1}$ $(n - 1)$ -dense(T, Γ) subsets of X that are pairwise T -apart. So $\forall x < \min X \exists y < \min X_1 \forall z < \max X_1 \theta(x, y, z)$ (since $\min X \leq \max X_0$ and X_0 and X_1 are T -apart).

Therefore, X is indeed n -dense(T, Γ). We conclude by induction. \square

The following theorem is the one which will actually be used in our applications in Section 5.

Theorem 3.12. *Let Γ be an RT-like. If ω^n -largeness(T, Γ) is a largeness notion provably in $\text{RCA}_0 + \text{BS}_2^0 + T$ for every $n \in \omega$ and every Π_3^0 formula T , then $\text{WKL}_0 + \text{BS}_2^0 + \Gamma$ is a $\forall\Pi_4^0$ -conservative extension of $\text{RCA}_0 + \text{BS}_2^0$.*

Proof. Fix some $n \in \omega$ and $\theta(A \upharpoonright_z, a, x, y, z)$ a Σ_0^0 formula. Let us show that $\text{RCA}_0 + \text{BS}_2^0 \vdash n\text{-PH}_\theta(\Gamma)$. Fix a model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{BS}_2^0$, some $a, b \in M$ and some set $A \in S$ such that $\mathcal{M} \models \forall x \exists y \forall z \theta(A \upharpoonright_z, a, x, y, z)$ holds. Let $T(A, a) \equiv \forall x \exists y \forall z \theta(A \upharpoonright_z, a, x, y, z)$. By Proposition 3.11, n -density($T(A, a), \Gamma$) is a largeness notion provably in $\text{RCA}_0 + \text{BS}_2^0 + T(A, a)$ for every $n \in \omega$, thus n -density($T(A, a), \Gamma$) is a largeness notion in \mathcal{M} . Since (b, ∞) is an M -infinite set, then there is an n -dense($T(A, a), \Gamma$) set $X \in S$ such that $\min X > b$. So $\mathcal{M} \models n\text{-PH}_\theta(\Gamma)$. Thus, by Theorem 3.9, $\text{WKL}_0 + \Gamma$ is a $\forall\Pi_4^0$ -conservative extension of $\text{RCA}_0 + \text{BS}_2^0$. \square

4 Π_1^1 conservation of the grouping principle

The grouping principle was defined by Patey and Yokoyama [22] as a combinatorial principle needed for an inductive construction of ω^n -large solutions to the Erdos-Moser theorem. They proved that the grouping principle is Π_3^0 conservative over RCA_0 and that it implies $\text{B}\Sigma_2^0$. In this section, we prove that this principle is actually Π_1^1 conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.

There exist two notions of grouping: a finitary one (FGP_k^n) and an infinitary one (GP_k^n), the former following from the latter. We first define the infinitary version.

Definition 4.1 (\mathcal{L} -grouping). *Let \mathcal{L} be a largeness notion and $f : [\mathbb{N}]^n \rightarrow 2$ a coloring. A (finite or infinite) sequence of \mathcal{L} -large sets $F_0 < F_1 < \dots$ of length $k \in \mathbb{N} \cup \{\infty\}$ is an \mathcal{L} -grouping for f if for every $H \in [k]^n$, f is monochromatic on $\prod_{i \in H} F_i$.*

In particular, for $n = 1$, $F_0 < F_1 < \dots$ is an \mathcal{L} -grouping if for every i , F_i is f -homogeneous.

Statement (Grouping principle). GP_k^n : *For every largeness notion \mathcal{L} and every coloring $f : [\mathbb{N}]^n \rightarrow k$, there is an infinite \mathcal{L} -grouping.*

The remainder of this section is dedicated to the proof of the following theorem.

Theorem 4.2. $\text{WKL}_0 + \text{GP}_2^2$ is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.

Following the reverse mathematical practice, GP_2^2 can be decomposed into the cohesiveness principle (COH) and a Δ_2^0 version of GP_2^1 ($\Delta_2^0\text{-GP}_2^1$).

Statement. $\Delta_2^0\text{-GP}_2^1$: *For every largeness notion \mathcal{L} and every Δ_2^0 set A , there is an infinite \mathcal{L} -grouping for A .*

An infinite set $C \subseteq \mathbb{N}$ is *cohesive* for an infinite sequence of sets R_0, R_1, \dots if for every $x \in \mathbb{N}$, $C \subseteq^* R_x$ or $C \subseteq^* \overline{R}_x$, where \subseteq^* denotes inclusion up to finitely many elements.

Statement (Cohesiveness). COH : *Every infinite sequence of sets admits an infinite cohesive set.*

The following decomposition can be considered as folklore:

Lemma 4.3 (Folklore). $\text{RCA}_0 + \text{B}\Sigma_2^0 \vdash \text{COH} + \Delta_2^0\text{-GP}_2^1 \rightarrow \text{GP}_2^2$.

Proof. Fix some coloring $f : [\mathbb{N}]^2 \rightarrow 2$ and largeness notion \mathcal{L} . By COH , letting $R_x = \{y : f(x, y) = 1\}$, there is an infinite set C which is cohesive for R_0, R_1, \dots . In particular, $\lim_{y \in C} f(x, y)$ exists for every $x \in M$. Say $C = \{c_0 < c_1 < \dots\}$. Let $A(x) = \lim_y f(c_x, c_y)$. By $\Delta_2^0\text{-GP}_2^1$, there is an infinite \mathcal{L} -grouping $F_0 < F_1 < \dots$ for A . For every $s \in M$, let $G_s = \{c_x : x \in F_s\}$. Then for every $s \in M$, either $\forall x \in F_s \lim_{y \in C} f(x, y) = 0$, or $\forall x \in F_s \lim_{y \in C} f(x, y) = 1$. Then, using $\text{B}\Sigma_2^0$, by computably thinning out the sequence, we obtain an M -infinite grouping for f . \square

Given a model $\mathcal{M} = (M, S) \models \text{RCA}_0$ and a set $G \subseteq M$, we write $\mathcal{M}[G]$ for the model whose first-order part is M and whose second-order part consists of the Δ_1^0 -definable sets with parameters in $\mathcal{M} \cup \{G\}$. If $\mathcal{M} \cup \{G\} \models \text{IS}_1^0$, then $\mathcal{M}[G] \models \text{IS}_1^0$. By the amalgamation theorem of Yokoyama [26, Theorem 2.2] and the fact that WKL_0 and COH are both Π_1^1 conservative over $\text{RCA}_0 + \text{BS}_2^0$, Theorem 4.2 is a direct consequence of the following proposition.

Proposition 4.4. *Consider a countable model $\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{BS}_2^0$ topped by a set $Y \in S$. For every largeness notion $\mathcal{L} \in S$ and every $\Delta_2^0(Y)$ -set $A \subseteq M$, there exists a $G \subseteq M$ such that:*

- G is an M -infinite \mathcal{L} -grouping for A or G is a witness that \mathcal{L} is not a largeness notion (i.e. an M -unbounded set with no M -finite subset in \mathcal{L}).
- $Y' \geq (Y \oplus G)'$
- $\mathcal{M}[G] \models \text{RCA}_0 + \text{BS}_2^0$.

Proof. Fix a uniform enumeration of all Y -primitive recursive non-empty tree functionals T_0, T_1, \dots , that is, for every $n \in M$ and every X , T_n^X is an \mathcal{M} -infinite tree. Let

$$\mathcal{C} = \left\{ \bigoplus_a X_a : \forall a \forall n \ X_{\langle a, n \rangle} \in [T_n^{Y \oplus X_0 \oplus \dots \oplus X_a}] \right\}$$

The class \mathcal{C} is $\Pi_1^0(Y)$ and non-empty. There exists a primitive Y -recursive tree whose infinite path are the elements of \mathcal{C} , so by [9, Corollary I.3.10(3)], there is a set $P = \bigoplus_a X_a \in \mathcal{C}$ such that $Y' \geq_T (P \oplus Y)'$ and $\mathcal{M}[P] \models \text{RCA}_0 + \text{BS}_2^0$. Let $\mathcal{N} = (M, \{X_a : a \in M\})$. Note that \mathcal{N} is an ω -extension of \mathcal{M} and that $\mathcal{N} \models \text{WKL}_0 + \text{BS}_2^0$. Assume \mathcal{L} is a largeness notion within \mathcal{N} , otherwise we are done.

Definition 4.5 (Condition). *A condition is a tuple (\vec{F}, \vec{g}) such that \vec{F} is a finite \mathcal{L} -grouping for A , and $\langle g_0, \dots, g_{p-1} \rangle \in \mathcal{N}$ are 2-colorings of M .*

Note that every set $X \in \mathcal{N}$ has a code with respect to P , that is, some $a \in M$ such that $X = X_a$ (where $\bigoplus_a X_a = P$). Thus, since $\langle g_0, \dots, g_{p-1} \rangle \in \mathcal{N}$, a condition (\vec{F}, \vec{g}) can be represented by an integer in M . For simplicity of notation, we identify a condition with its code.

Definition 4.6 (Extension). *A condition $d = (\vec{E}, \vec{h})$ extends a condition $c = (\vec{F}, \vec{g})$ (written $d \leq c$) if:*

- \vec{g} is a subsequence of \vec{h}
- \vec{E} is of the form $\vec{F}, H_0, \dots, H_{s-1}$
- H_0, \dots, H_{s-1} is an \mathcal{L} -grouping for \vec{g} .

Definition 4.7 (Forcing relation). *Let Φ_e be a Turing functional and $c = (\vec{F}, \vec{g})$ a condition:*

- $c \Vdash \Phi_e^{Y \oplus G} \downarrow$ if $\Phi_e^{Y \oplus \vec{F}} \downarrow$
- $c \Vdash \Phi_e^{Y \oplus G} \uparrow$ if $\Phi_e^{Y \oplus (\vec{F}, \vec{E})} \uparrow$ for every finite \mathcal{L} -grouping \vec{E} for \vec{g} such that $\max \vec{F} < \min \vec{E}$.

The relation $c \Vdash \Phi_e^{Y \oplus G} \downarrow$ is $\Delta_0^0(Y)$ and $c \Vdash \Phi_e^{Y \oplus G} \uparrow$ is $\Pi_1^0(Y \oplus \vec{g})$.

Lemma 4.8. Let $c = (\vec{F}, \vec{g})$ be a condition. For every $k \in M$, there exists some \mathcal{M} -finite $\sigma' \in 2^k$ and an extension $d = (\vec{E}, \vec{h}) \leq c$ such that $d \Vdash (G \oplus Y)' \upharpoonright_k = \sigma'$ (Where $d \Vdash (G \oplus Y)' \upharpoonright_k = \sigma'$ is the $\Pi_1^0(Y \oplus \vec{h})$ relation meaning that if $\sigma'(e) = 0$ then $d \Vdash \Phi_e^{G \oplus Y} \uparrow$ and if $\sigma'(e) = 1$ then $d \Vdash \Phi_e^{G \oplus Y} \downarrow$).

Proof. For $s < k$, let $\varphi(s)$ be the formula that holds if for every $(k-s)$ -tuple of colorings $\vec{f} = f_0, \dots, f_{k-s-1}$ there exists a set of indexes $e_0 < \dots < e_{s-1} < k$ and a finite \mathcal{L} -grouping \vec{H} for (\vec{g}, \vec{f}) such that $\max \vec{F} < \min \vec{H}$ and $\Phi_{e_i}^{Y \oplus (\vec{F}, \vec{H})} \downarrow$ for all $i < s$. By Lemma 1.10, $\varphi(s)$ is $\Sigma_1^0(Y \oplus \vec{g})$.

Let $V = \{s < k : \varphi(s)\}$. V is \mathcal{M} -finite since $\mathcal{N} \models \text{IS}_1^0$ and V contains 0. Therefore, one of the following cases holds:

Case 1: $k-1 \in V$. In this case, by considering the coloring induced by A , by Lemma 1.10, there exists a finite \mathcal{L} -grouping \vec{H} for (\vec{g}, A) with $\max \vec{F} < \min \vec{H}$ and such that for every index $e < k$, $\Phi_e^{Y \oplus (\vec{F}, \vec{H})} \downarrow$. In this case, $((\vec{F}, \vec{H}), \vec{g}) \Vdash (G \oplus Y)' \upharpoonright_k = 11 \dots 11$.

Case 2: there is some $s < k-1$ with $s \in V$ but $s+1 \notin V$. Consider the following $\Pi_1^0(Y \oplus \vec{g})$ class: \mathcal{C} is the class of all $(k-s-1)$ -tuple of colorings $\vec{f} = f_0, \dots, f_{k-s-2}$ such that for every set of indexes $e_0 < \dots < e_{s-2} < k$ and every finite \mathcal{L} -grouping \vec{H} for (\vec{g}, \vec{f}) satisfying $\max \vec{F} < \min \vec{H}$, there exists a $i < s-1$ such that $\Phi_{e_i}^{Y \oplus (\vec{F}, \vec{H})} \uparrow$. Since $s+1 \notin V$, the class \mathcal{C} is non-empty. Moreover, since $Y, \vec{g} \in \mathcal{N} \models \text{WKL}_0$, there is some $\vec{f} \in \mathcal{C} \cap \mathcal{N}$.

Then \vec{f}, A is a $(k-s)$ -uple of colorings and since $s \in V$, there exists a set of indexes $e_0 < \dots < e_{s-1} < k$ and a finite \mathcal{L} -grouping \vec{H} for (\vec{g}, \vec{f}, A) such that $\max \vec{F} < \min \vec{H}$ and $\Phi_{e_i}^{Y \oplus (\vec{F}, \vec{H})} \downarrow$ for all $i < s$.

Letting $d = ((\vec{F}, \vec{H}), (\vec{g}, \vec{f}))$ and $\sigma' \in 2^k$ the binary encoding of the set $\{e_0, \dots, e_{s-1}\}$, we have that d extends c and $d \Vdash (G \oplus Y)' \upharpoonright_k = \sigma'$: By choice of \vec{H} and the e_i , it is clear that $d \Vdash \Phi_e^{G \oplus Y} \downarrow$ for every $e < k$ such that $\sigma'(e) = 1$ and by choice of \vec{f} , for every $e < k$ such that $\sigma'(e) = 0$ since $\{e_0, \dots, e_{s-1}, e\}$ is of cardinality $s+1$, no extension of d can force $\Phi_e^{G \oplus Y}$ to stop (and therefore $d \Vdash \Phi_e^{G \oplus Y} \uparrow$).

□

Lemma 4.9. Let $c = (\vec{F}, \vec{g})$ be a condition. For every $k \in M$, there exists an extension $d = (\vec{E}, \vec{g}) \leq c$ such that $\max \vec{E} > k$.

Proof. Consider the $\Pi_1^0(Y \oplus \vec{g})$ -class \mathcal{D} of all colorings h such that there is no \mathcal{L} -grouping \vec{E} with $\max \vec{E} > k$ for \vec{g}, h with \vec{E} is of the form $\vec{F}, H_0, \dots, H_{s-1}$.

If this class was non-empty, then since $Y, \vec{g} \in \mathcal{N} \models \text{WKL}_0$, then there is an element $f \in \mathcal{D} \cap \mathcal{N}$. By RT^1 , there exists an \mathcal{N} -infinite set X homogenous for \vec{g}, f such that $X \cap [k, t]_{\mathbb{N}}$ is not \mathcal{L} -large for any $t > k$ contradicting our assumption on the largeness of \mathcal{L} within \mathcal{N} .

So the class \mathcal{D} is empty and in particular does not contain the coloring A . Therefore, such an extension d exists. \square

Construction. We will build a decreasing sequence (\vec{F}_s, \vec{g}_s) of conditions and then take for G the union of the \vec{F}_s . We will also build an increasing sequence (σ'_s) such that $(G \oplus Y)'$ will be the union of the σ'_s . Initially, we take $\vec{F}_0 = \vec{g}_0 = \emptyset$ and $\sigma'_0 = \epsilon$, and during the construction, we will ensure that we have $|\vec{F}_s|, |\vec{g}_s|, |\sigma'_s| \leq s$ at every stage. Each stage will be either of type \mathcal{R} or of type \mathcal{S} . The stage 0 is of type \mathcal{R} .

Assume that (\vec{F}_s, \vec{g}_s) and σ'_s are already defined. Let $s_0 < s$ be the latest stage at which we switched the stage type. We have three cases.

Case 1: s is of type \mathcal{R} . If there exists some \vec{F} and \vec{g} such that $|\vec{g}|, |\vec{F}| \leq s$ and some $\sigma' \in 2^{s_0}$ such that $(\vec{F}, \vec{g}) \leq (\vec{F}_s, \vec{g}_s)$, and $(\vec{F}, \vec{g}) \Vdash (G \oplus Y)' \upharpoonright_{s_0} = \sigma'$, then let $\vec{F}_{s+1} = \vec{F}$, $\vec{g}_{s+1} = \vec{g}$, $\sigma'_{s+1} = \sigma'$ and let $s+1$ be of type \mathcal{S} . Otherwise, the elements are left unchanged and we go to the next stage.

Case 2: s is of type \mathcal{S} . If there exists some \vec{F} and \vec{g} such that $|\vec{g}|, |\vec{F}| \leq s$, $\max \vec{F} > s_0$ and $(\vec{F}, \vec{g}) \leq (\vec{F}_s, \vec{g}_s)$, then let $\vec{F}_{s+1} = \vec{F}$ and $\vec{g}_{s+1} = \vec{g}$ and let $s+1$ be of type \mathcal{R} . Otherwise, the elements are left unchanged and we go to the next stage.

This completes the construction.

Verification. Since the size of \vec{F}_s, \vec{g}_s and σ'_s are bounded by s , there is a $\Delta_1^0(Y' \oplus P)$ -formula $\phi(s)$ stating that the construction can be pursued up to stage s . Our construction implies that the set $\{s \mid \phi(s)\}$ is a cut, so by $\text{ID}_1^0(Y' \oplus P)$ (which follows from BS_2^0 in $\mathcal{M}[P]$), the construction can be pursued at every stage.

Let $G = \bigcup_{s \in M} \sigma'_s$. By Lemma 4.8 and Lemma 4.9, each type of stage changes \mathcal{M} -infinitely often. Thus, $\{|\sigma'_s| : s \in M\}$ is \mathcal{M} -unbounded, so $(Y \oplus P)' \geq_T (G \oplus Y)'$. Since $Y' \geq_T (Y \oplus P')$, then $Y' \geq_T (G \oplus Y)'$. In particular, $\mathcal{M}[G] \models \text{RCA}_0 + \text{BS}_2^0$ and G is M -infinite and is therefore an M -infinite L -grouping for A . \square

We are now ready to prove the main theorem of this section, namely, that the grouping principle is Π_1^1 conservative over $\text{RCA}_0 + \text{BS}_2^0$. It can be alternatively proved by showing that $\Delta_2^0\text{-GP}_2^1$ and COH both are Π_1^1 -conservative over $\text{RCA}_0 + \text{BS}_2^0$, and using the amalgamation theorem of Yokoyama [26, Theorem 2.2]. However, we give a direct argument for the sake of simplicity:

Proof of Theorem 4.2. Assume $\text{RCA}_0 + \text{BS}_2^0 \not\models \forall X \phi(X)$ for ϕ arithmetical. Then, by completeness and the Löwenheim-Skolem theorem, there exists a model

$\mathcal{M} = (M, S) \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \neg\phi(A)$ for an $A \in S$. We can furthermore assume that \mathcal{M} is topped by A .

By Proposition 4.4 and Chong, Slaman and Yang [4, Theorem 3.2], we can build the following sequence of countable topped model of $\text{RCA}_0 + \text{B}\Sigma_2^0$: $\mathcal{M}_0 = \mathcal{M} \subseteq \mathcal{M}_1 \subseteq \mathcal{M}_2 \subseteq \dots$ satisfying that

- (1) for every i , every largeness notion $\mathcal{L} \in \mathcal{M}_i$ and every set A that is Δ_2^0 in \mathcal{M}_i , there is either a j such that \mathcal{L} is no longer a largeness notion in \mathcal{M}_j , or a set $G \in \mathcal{M}_j$ such that G is an M -infinite \mathcal{L} -grouping for A .
- (2) for every i , every uniform sequence of sets $\vec{R} \in \mathcal{M}_i$, there is some j and an M -infinite \vec{R} -cohesive set $C \in \mathcal{M}_j$.

Let $\mathcal{M}' = \bigcup_{n \in \omega} \mathcal{M}_n$. Clearly, $\mathcal{M}' \models \text{RCA}_0 + \text{B}\Sigma_2^0 + \text{COH} + \Delta_2^0\text{-GP}_2^1 + \neg\phi(A)$. By Lemma 4.3, $\mathcal{M}' \models \text{GP}_2^2$. Therefore, GP_2^2 is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$. \square

Remark 4.10. *The above construction shows that $\text{RCA}_0 + \text{B}\Sigma_2^0$ proves the \ll_2 -basis theorem for $\Delta_2^0\text{-GP}_2^1$ (in the sense of [6]), and thus GP_2^2 also admits \ll_2 -basis theorem within $\text{RCA}_0 + \text{B}\Sigma_2^0$. This implies that the above Π_1^1 -conservation theorem admits poly-time proof transformation. In general, an upcoming result by Ikari and Yokoyama shows that any RT-like principle provable from $\text{WKL}_0 + \text{RT}_2^2$ admits poly-time proof transformation if it is Π_1^1 -conservative over $\text{RCA}_0 + \text{B}\Sigma_2^0$.*

4.1 Finite grouping principle with T -apartness

Thanks to the generalized Parsons theorem, we can turn Π_1^1 conservation of the grouping principle into a quantitative finitary version based on ω^n -largeness(T).

Definition 4.11 (Finite grouping principle for T). *Let $\mathcal{L}_0, \mathcal{L}_1$ be largeness notions and $f : [X]^n \rightarrow k$ a coloring. A finite sequence of \mathcal{L}_0 -large sets $F_0 < F_1 < \dots < F_{k-1}$ is an $(\mathcal{L}_0, \mathcal{L}_1)$ -grouping(T) for f if:*

- for any $H \subseteq_{\text{fin}} \mathbb{N}$, if $H \cap F_i \neq \emptyset$ for every $i < k$, then $H \in \mathcal{L}_1$, and,
- for every $H \in [k]^n$, f is monochromatic on $\prod_{i \in H} F_i$
- for every $i < j < k - 1$, F_i and F_j are T -apart.

Let $(\mathcal{L}_0, \mathcal{L}_1)$ -FGP(T) be the statement that for any infinite set $X \subseteq \mathbb{N}$, there exists some finite set $Y \subseteq X$ such that for any coloring $f : [Y]^n \rightarrow k$, there exists an $(\mathcal{L}_0, \mathcal{L}_1)$ -grouping(T) for f .

Recall that a and A are the first-order and second-order parameters appearing in the formula T .

Proposition 4.12. *Let \mathcal{L}_0 and \mathcal{L}_1 be $\Delta_0^{a,A}$ -definable largeness notions provably in $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$. Then $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$ proves $(\mathcal{L}_0, \mathcal{L}_1)$ -FGP(T).*

Proof. $\forall a \forall A (T \rightarrow (\mathcal{L}_0, \mathcal{L}_1)\text{-FGP}(T))$ is a Π_1^1 statement. So by Π_1^1 conservation of $\text{WKL}_0 + \text{GP}_2^2$ over $\text{RCA}_0 + \text{B}\Sigma_2^0$, it is sufficient to prove the result using $\text{WKL}_0 + \text{B}\Sigma_2^0 + T + \text{GP}_2^2$.

By GP_2^2 , for any infinite set X and for any coloring $f : [X]^2 \rightarrow 2$ there exists an infinite \mathcal{L}_0 -grouping for f . By $\text{B}\Sigma_2^0 + T$, we can furthermore assume the blocks of these groupings to be T -apart.

For each of these grouping $Y_0 < Y_1 < \dots$, we can consider the following finitely branching tree S of finite sequences σ such that $\sigma(i) \in Y_i$ for all $i < |\sigma|$ and such that the finite set $\{\sigma(i) \mid i < |\sigma|\}$ is not \mathcal{L}_1 -large. Since \mathcal{L}_1 is a largeness notion, the tree S has no infinite branch, and therefore by WKL_0 , there is some bound n on the depth of the tree. So $Y_0 < Y_1 < \dots < Y_{n-1}$ is an $(\mathcal{L}_0, \mathcal{L}_1)$ -grouping(T) for the corresponding coloring.

And again by WKL_0 , there exists some finite set $Y \subseteq X$ such that for any coloring $f : [Y]^2 \rightarrow 2$ there exists an $(\mathcal{L}_0, \mathcal{L}_1)$ -grouping(T) for f .

So $\text{WKL}_0 + \text{B}\Sigma_2^0 + T + \text{GP}_2^2 \vdash (\mathcal{L}_0, \mathcal{L}_1)\text{-FGP}(T)$ and therefore $\text{RCA}_0 + \text{B}\Sigma_2^0 + T \vdash (\mathcal{L}_0, \mathcal{L}_1)\text{-FGP}(T)$. \square

We will use the finite grouping principle with T -apartness under the following form. Note that in the following proposition, the integer n might depend not only on k and ℓ , but also on T . However, by adapting the direct combinatorial proof of Kołodziejczyk and Yokoyama [16, Section 2.1] of the finitary grouping principle with explicit bounds, one obtains a bound which does not depend on T (see Remark 4.14).

Proposition 4.13. *For any $k, \ell \in \omega$, there exists $n \in \omega$ such that $\text{I}\Sigma_1^0$ proves that $\forall Z \subseteq_{\text{fin}} \mathbb{N}$, if Z is ω^n -large(T) and exp-sparse then*

$\forall f : [Z]^2 \rightarrow 2$, there exists an $(\omega^k\text{-large}(T), \omega^\ell\text{-large}(T))$ -grouping(T) for f

Proof. Using Theorem 1.6, it is sufficient to show that $\text{WKL}_0 + \text{B}\Sigma_2^0 + T$ proves that $\forall X$ if X is infinite then

$\exists Y \subseteq X \forall f : [Y]^2 \rightarrow 2$, there exists an $(\omega^k\text{-large}(T), \omega^\ell\text{-large}(T))$ -grouping(T) for f

Which we have by Proposition 4.12 (And by Proposition 2.7, ω^k -largeness(T) and ω^ℓ -largeness(T) are largeness notions provably in $\text{RCA}_0 + \text{B}\Sigma_2^0 + T$). \square

Remark 4.14. *The proof of Proposition 4.13 involves a generalized Parsons theorem (Theorem 1.6) which does not provide explicit bounds for the existence of a grouping. Kołodziejczyk and Yokoyama [16, Section 2.1] gave a direct combinatorial proof of the finitary grouping principle, with explicit bounds.*

Largeness(T) shares many combinatorial features with the standard notion of largeness. However, there are some differences, which impact the explicit bounds of the pigeonhole lemma for largeness(T) (Lemma 2.9). Indeed, to obtain an ω^n -large(T) set after an application of RT^1 , one starts with an ω^{2n} -large(T) set, while an ω^{n+1} -large set is sufficient in the case of standard largeness.

Propagating this difference to the proof of Kołodziejczyk and Yokoyama [16, Section 2.1], one obtain (with exp-sparsity) ω^{2n} -largeness(T) for [16, Lemma

2.5], hence ω^{4n+1} -largeness(T) for [16, Lemma 2.6], ω^{16n+5} -largeness(T) for [16, Lemma 2.7] and $\omega^{16^k \times (n+1)}$ -largeness(T) for [16, Theorem 2.4].

For $f : X \rightarrow 2$ a coloring of singletons, the existence of an $(\omega^n, 2)$ -grouping(T) for $g : (x, y) \mapsto f(x)$ yields an ω^n -large(T) f -homogeneous subset of X (by taking the first block of the grouping). We shall see in Section 7 that for all $n \in \omega$ there exists some Π_3^0 -formula T and some ω^{2n-1} -large(T) set X such that X is not ω^n -large(T, RT_2^1), therefore the bound obtained for the existence of an $(\omega^n\text{-large}(T), \omega^k\text{-large}(T))$ -grouping(T) is tight in the sense that it is not possible to obtain one with X $\omega^{n+h(k)}$ -large(T) for some $h : \omega \rightarrow \omega$ as in [16].

5 Applications to RT-like theorems

In this section, we apply the framework developed in Section 3 to prove that the Erdős-Moser theorem, the Ascending Descending Sequence principle and Ramsey's theorem for pairs are $\forall \Pi_4^0$ conservative over $\text{RCA}_0 + \text{BS}_2^0$.

5.1 $\forall \Pi_4^0$ conservation of EM

As explained in Section 1.2, RT_2^2 can be decomposed into two combinatorially simpler statements, namely, ADS and EM. Thanks to the amalgamation theorem (Theorem 1.9), since $\text{RCA}_0 + \text{ADS}$ is a Π_1^1 conservative extension of $\text{RCA}_0 + \text{BS}_2^0$, the heart of the main question lies in the first-order part of EM.

As in Proposition 4.13, the bound k_n depends on n , but also on T by default. However, the alternative combinatorial proof yields explicit bounds which do not depend on T (see Remark 5.2).

Proposition 5.1. *For any $n \in \omega$, there exists some $k_n \in \omega$ such that IS_1^0 proves that $\forall X \subseteq_{\text{fin}} \mathbb{N}$ if X is ω^{k_n} -large(T) and exp-sparse then X is ω^n -large(T, EM).*

Proof. By external induction on n :

Case $n = 0$: Any ω^0 -large(T) set is ω^0 -large(T, EM). Thus, take $k_0 = 0$.

Case $n > 0$, assume the property to be true at rank $n - 1$. By Proposition 4.13, there is some $k_n \in \omega$ such that if X is ω^{k_n} -large(T) and exp-sparse, then

$\forall f : [X]^2 \rightarrow 2$, there exists an $(\omega^{k_n-1}\text{-large}(T), \omega^6\text{-large})$ -grouping(T) for f

Note that ω^6 -largeness is not ω^6 -largeness(T) in the grouping above, but any ω^6 -large(T) set is also ω^6 -large.4

We need to check that X is ω^n -large(T, EM). Let $f : [X]^2 \rightarrow 2$ be an instance of EM. There exists an $(\omega^{k_n-1}\text{-large}(T), \omega^6\text{-large})$ -grouping(T) $X_0 < X_1 < \dots < X_{\ell-1}$ for f .

By the inductive hypothesis, each block X_i is ω^{n-1} -large(T, EM), so for each $i < \ell$ there exists $Y_i \subseteq X_i$ ω^{n-1} -large(T) and transitive for f . By definition of a grouping, f induces a tournament on the pairs of blocks and $\{\min Y_i : i < \ell\}$ is ω^6 -large and therefore ω -large(EM) by Ketonen and Solovay (see Theorem 2.3).

Therefore, there is a subset $I \subseteq \{0, \dots, l-1\}$ such that f is transitive on $\{\min Y_i : i \in I\}$ which is ω -large. The set $Y = \bigcup_{i \in I} Y_i$ is f -transitive and ω^n -large(T). We conclude by induction. \square

Remark 5.2. *One can propagate the explicit bounds of the grouping principle computed in Remark 4.14 to obtain an explicit bound for Proposition 5.1: if X is $\omega^{(16^6+1)^n}$ -large(T), then it is ω^n -large(T, EM).*

Note that we obtain an exponential upper bound for largeness(T, EM) while the upper bound for regular largeness(EM) is polynomial. Since the bounds on the grouping principle are tight, one would need to get rid of the applications of the grouping principle to obtain a better upper bound, which seems unlikely.

Theorem 5.3. $\text{WKL}_0 + \text{EM}$ is $\forall \Pi_4^0$ -conservative over $\text{RCA}_0 + \text{BS}_2^0$.

Proof. By Proposition 5.1, and Proposition 2.8, for all $n \in \omega$ and T a Π_3^0 -formula, $\text{RCA}_0 + \text{BS}_2^0 + T$ proves that ω^n -largeness(T, EM) is a largeness notion. So by Theorem 3.12, EM is a $\forall \Pi_4^0$ -conservative extension of $\text{RCA}_0 + \text{BS}_2^0$. \square

We are now ready to prove the main theorem of this article:

Theorem 1.4. $\text{WKL}_0 + \text{RT}_2^2$ is $\forall \Pi_4^0$ -conservative over $\text{RCA}_0 + \text{BS}_2^0$.

Proof. Since ADS is Π_1^1 -conservative over $\text{RCA}_0 + \text{BS}_2^0$, it is in particular $\forall \Pi_4^0$ -conservative. By the amalgamation theorem (Theorem 1.9) and the fact that $\text{RCA}_0 \vdash \text{RT}_2^2 \leftrightarrow \text{ADS} + \text{EM}$ (see Bovykin and Weiermann [1]), remains to show that $\text{WKL}_0 + \text{EM}$ is also $\forall \Pi_4^0$ -conservative over $\text{RCA}_0 + \text{BS}_2^0$, which is Theorem 5.3. \square

5.2 $\forall \Pi_4^0$ conservation of ADS

In this section, we give a direct inductive proof in $\text{RCA}_0 + \text{BS}_2^0 + T$ that ω^n -largeness(T, ADS) is a largeness notion, with explicit polynomial bounds. The proof is very similar to [16, Theorem 2.11], but requires some minor adaptation to handle T -apartness.

Proposition 5.4. *Fix $n \in \omega$, $|\Sigma_1^0|$ proves that any ω^{4n+4} -large(T) set X is ω^n -large(T, ADS).*

Proof. Let $f : [X]^2 \rightarrow 2$ be a transitive coloring and assume by contradiction that no f -homogeneous ω^n -large(T) subset of X exists.

Given $x, y \in X$ with $x < y$, we say that an interval $[x, y]$ is (i, α) -long(T) if there exists an α -large(T) set $H \subseteq [x, y] \cap X$ such that $x \in H$, $H \cup \{y\}$ is f -homogeneous with colour i and H and $[y, \max X] \cap X$ are T -apart. Define a new colouring $Q : [X]^2 \rightarrow 4n$ as follows:

$$Q(x, y) = \begin{cases} 4k & \text{if } [x, y] \text{ is } (0, \omega^k)\text{-long}(T) \text{ but not } (0, \omega^k + 1)\text{-long}(T), \\ 4k + 1 & \text{if } [x, y] \text{ is } (0, \omega^k + 1)\text{-long}(T) \text{ but not } (0, \omega^{k+1})\text{-long}(T), \\ 4k + 2 & \text{if } [x, y] \text{ is } (1, \omega^k)\text{-long}(T) \text{ but not } (1, \omega^k + 1)\text{-long}(T), \\ 4k + 3 & \text{if } [x, y] \text{ is } (1, \omega^k + 1)\text{-long}(T) \text{ but not } (1, \omega^{k+1})\text{-long}(T), \end{cases}$$

By Theorem 2.3, there exists a Q -homogeneous ω -large subset Y of X . The remainder of the proof is exactly that of [16, Theorem 2.11], *mutatis mutandis*. Y being ω -large (and not ω -large(T)) is sufficient to create T -apart blocks thanks to the definition of an (i, α) -long(T) interval $[x, y]$, which ensure the existence of an α -large(T) subset $H \subseteq [x, y]$ which is T -apart from $[y, \max X]$. \square

Corollary 5.5. *If X is $\omega^{(16^6+1)^{4n+4}}$ -large(T), then it is ω^n -large(T, RT_2^2).*

Proof. Let $f : [X]^2 \rightarrow 2$ be a coloring. By Remark 5.2, there exists an ω^{4n+4} -large(T) f -transitive set $Y \subseteq X$. By Proposition 5.4, there exists an ω^n -large(T) f -homogeneous subset $Z \subseteq Y$. \square

6 A conservation theorem over IS_1^0

Since BS_2^0 is a $\forall\Pi_4^0$ statement which is strictly stronger than IS_1^0 , then BS_2^0 (and *a fortiori* RT_2^2) is not a $\forall\Pi_4^0$ conservative extension of IS_1^0 . However, in this section, we shall give two proofs that it is the case for a particular kind of $\forall\Pi_4^0$ formulas. The first proof uses the techniques developed in this article, while the second is based on the fact that every countable model of IS_n^0 admits a Σ_{n+1}^0 -elementary cofinal extension which is a model of BS_{n+1}^0 .

Definition 6.1. *A formula is weakly $\forall\Pi_4^0$ if it is of the form*

$$\forall A \forall a \exists x \forall y \exists x' < x \forall y' < y \exists z \theta(A \upharpoonright_z, a, x', y', z)$$

where θ is a Σ_0^0 formula.

Intuitively, weakly $\forall\Pi_4^0$ formulas are $\forall\Pi_4^0$ formulas for which the applications of BS_2^0 are “hardcoded” in the syntax of the formula.

Proposition 6.2. *Over $\text{RCA}_0 + \text{BS}_2^0$, every $\forall\Pi_4^0$ formula is equivalent to a weakly $\forall\Pi_4^0$ formula.*

Proof. Let $S = \forall A \forall a \exists x \forall y \exists z \theta(A \upharpoonright_z, a, x, y, z)$ be a $\forall\Pi_4^0$ formula, where θ is Σ_0^0 . Let $S' = \forall A \forall a \exists x \forall y \exists x' < x \forall y' < y \exists z \theta(A \upharpoonright_z, a, x', y', z)$ be its corresponding weakly $\forall\Pi_4^0$ formula. $\text{RCA}_0 \vdash S \rightarrow S'$. We now prove that $\text{RCA}_0 + \text{BS}_2^0 \vdash S' \rightarrow S$. Fix some A and some a . By S' , there is some x such that $\forall y \exists x' < x \forall y' < y \exists z \theta(A \upharpoonright_z, a, x', y', z)$. Let $f : \mathbb{N} \rightarrow x$ be defined by $f(y) = x'$ such that $\forall y' < y \exists z \theta(A \upharpoonright_z, a, x', y', z)$. By RT_2^1 , which is equivalent to BS_2^0 , there is some infinite f -homogeneous set H for some color $x' < x$. We claim that $\forall y' \exists z \theta(A \upharpoonright_z, a, x', y', z)$. Indeed, given y' , there is some $y > y'$ such that $y \in H$. Since $f(y) = x'$, then by definition of f , $\exists z \theta(A \upharpoonright_z, a, x', y', z)$. This completes the proof. \square

For the remainder of this section, fix a Σ_0^0 -formula $\theta(x, y, z)$ (with parameters) and let $T \equiv \forall x \exists y \forall x' < x \exists y' < y \forall z \theta(x', y', z)$.

Proposition 6.3. *For every $n \in \omega$, $\text{RCA}_0 + T$ proves that for every $k \geq 1$, $\omega^n \cdot k$ -largeness(T) is a largeness notion.*

Proof. Exactly the proof of Proposition 2.7, but the new form of T removes the use of $\text{B}\Sigma_2^0$. \square

Proposition 6.4. *If $\text{WKL}_0 + \text{B}\Sigma_2^0 \vdash \neg T$, then $\text{I}\Sigma_1^0 \vdash \neg T$.*

Proof. Assume $\text{WKL}_0 + \text{B}\Sigma_2^0 \vdash \neg T$. Then $\text{WKL}_0 + \text{B}\Sigma_2^0 + T \vdash \perp$. By Theorem 1.6 with $\psi \equiv \perp$, there is some $n \in \omega$ such that $\text{I}\Sigma_1^0$ proves $\forall x \forall Z \subseteq_{\text{fin}} (x, \infty)$

$$Z \text{ is } \omega^n\text{-large}(T) \text{ and exp-sparse} \rightarrow \perp$$

Hence $\text{I}\Sigma_1^0$ proves that exp-sparse ω^n -largeness(T) is not a largeness notion, therefore, by Proposition 6.3, $\text{RCA}_0 + T \vdash \perp$. By Π_1^1 conservation of RCA_0 over $\text{I}\Sigma_1^0$ (see Friedman [7]), $\text{I}\Sigma_1^0 \vdash \neg T$. \square

Proposition 6.5. *$\text{WKL}_0 + \text{B}\Sigma_2^0$ is conservative over $\text{I}\Sigma_1^0$ for weakly $\forall\Pi_4^0$ formulas.*

First proof of Proposition 6.5. Fix a Σ_0^0 -formula $\zeta(A \upharpoonright_z, a, x, y, z)$, and let $T(A, a) \equiv \forall x \exists y \forall x' < x \exists y' < y \forall z \neg \zeta(A \upharpoonright_z, a, x', y', z)$. Suppose $\text{I}\Sigma_1^0 \not\vdash \forall A \forall a \neg T(A, a)$. Then there is a model $\mathcal{M} = (M, S) \models \text{I}\Sigma_1^0 \wedge T(A, a)$ for some $A \in S$ and $a \in M$. Enrich the language with a constant symbol for A and a . In particular, $\text{I}\Sigma_1^0 \not\vdash \neg T(A, a)$, so by Proposition 6.4, $\text{WKL}_0 + \text{B}\Sigma_2^0 \not\vdash \neg T(A, a)$. It follows that there is a model $\mathcal{N} = (N, R) \models \text{WKL}_0 + \text{B}\Sigma_2^0 \wedge T(A, a)$, hence $\mathcal{N} \models \text{WKL}_0 + \text{B}\Sigma_2^0 \wedge \neg \forall A \forall a \neg T(A, a)$, thus $\text{WKL}_0 + \text{B}\Sigma_2^0 \not\vdash \forall A \forall a \neg T(A, a)$. \square

We now give an alternative and more traditional proof of Proposition 6.5 using the notion of cofinal Σ_n^0 -elementary extension.

Definition 6.6. *A model \mathcal{N} is a cofinal extension of $\mathcal{M} \subseteq \mathcal{N}$ (written $\mathcal{M} \subseteq_{\text{cf}} \mathcal{N}$) if for every $x \in \mathcal{N}$, there is some $y \in \mathcal{M}$ such that $x \leq y$. We write $\mathcal{M} \preceq_{n, \text{cf}} \mathcal{N}$ if \mathcal{N} is a cofinal Σ_n^0 -elementary extension of \mathcal{M} .*

The following theorem first appeared in Paris [20], and was independently discovered by Harvey Friedman.

Theorem 6.7 (Paris [20] ; Friedman). *Let $n \in \omega$ and $\mathcal{M} \models \text{I}\Sigma_n^0$. Then there is $\mathcal{N} \succeq_{n+1, \text{cf}} \mathcal{M}$ that satisfies $\text{B}\Sigma_{n+1}^0$.*

We are now ready to give an alternative proof of Proposition 6.5.

Second proof of Proposition 6.5. Fix a Σ_0^0 -formula $\zeta(A \upharpoonright_z, a, x, y, z)$, and let

$$T(A, a) \equiv \forall x \exists y \forall x' < x \exists y' < y \forall z \neg \zeta(A \upharpoonright_z, a, x', y', z)$$

Suppose $\text{I}\Sigma_1^0 \not\vdash \forall A \forall a \neg T(A, a)$. Then there is a model $\mathcal{M} = (M, S) \models \text{I}\Sigma_1^0 \wedge T(A, a)$ for some $A \in S$ and $a \in M$. Enrich the language with a constant

symbol for A and a . By Theorem 6.7, there exists a model $\mathcal{N} = (N, U) \succeq_{2,cf} \mathcal{M}$ of $\mathbf{B}\Sigma_2^0$.

We claim that $\mathcal{N} \models T(A, a)$. Fix some $x \in N$. By cofinality, let $x_1 \in M$ be such that $x \leq x_1$. Since $\mathcal{M} \models T(A, a)$, there is some $y_1 \in M$ such that $\mathcal{M} \models \forall x' < x_1 \exists y' < y_1 \forall z \neg \zeta(A \upharpoonright_z, a, x', y', z)$. Since $\mathcal{M} \models \mathbf{I}\Sigma_1^0$, the previous formula is equivalent to a Π_1^0 formula, so since \mathcal{N} is a Σ_2^0 -elementary extension of \mathcal{M} , $\mathcal{N} \models \forall x' < x_1 \exists y' < y_1 \forall z \neg \zeta(A \upharpoonright_z, a, x', y', z)$. In particular, $\mathcal{N} \models \exists y \forall x' < x_1 \exists y' < y_1 \forall z \neg \zeta(A \upharpoonright_z, a, x', y', z)$, and this for every $x \in N$. Thus $\mathcal{N} \models T(A, a)$. It follows that $\mathbf{B}\Sigma_2^0 \not\models \forall A \forall a \neg T(A, a)$.

Last, By Hájek [8], $\mathbf{WKL}_0 + \mathbf{B}\Sigma_2^0$ is a Π_1^1 conservative extension of $\mathbf{B}\Sigma_2^0$, so $\mathbf{WKL}_0 + \mathbf{B}\Sigma_2^0 \not\models \forall A \forall a \neg T(A, a)$. \square

Corollary 6.8. $\mathbf{WKL}_0 + \mathbf{RT}_2^2$ is weakly $\forall\Pi_4^0$ conservative over $\mathbf{I}\Sigma_1^0$.

Proof. Immediate from the combination of Theorem 1.4 and Proposition 6.5. \square

7 Lower bounds for largeness(T)

The proof structure of the Π_4^0 conservation theorem for Ramsey's theorem for pairs and two colors followed closely the one of Patey and Yokoyama [22]. In particular, the generalized Parsons theorem for $\mathbf{WKL}_0 + \mathbf{B}\Sigma_2^0 + T$ (Theorem 1.6) is used together with the Π_1^1 conservation of the grouping principle, to obtain a finitary version in terms of largeness(T) without explicit bounds (Proposition 4.13). Kołodziejczyk and Yokoyama [16, Theorem 2.4] computed explicit bounds to the finite grouping principle in terms of largeness, by iterating a pigeonhole lemma [16, Lemma 2.2].

The remainder of this section is devoted to the proof that the new bound of Lemma 2.9 for the pigeonhole principle is optimal.

Definition 7.1. An ω^n -decomposition of an ω^n -large set X is a finite sequence of ω^{n-1} -large subsets $X_0 < \dots < X_a$ of $X \setminus \min X$ for some $a \geq \min X - 1$. A finite set X is minimal for ω^n -largeness if it is ω^n -large and for every $x \in X$, $X \setminus \{x\}$ is not ω^n -large.

Note that by regularity of largeness, for every $x, n \in \mathbb{N}$, there is some $y > x$ such that $[x, y]$ is minimal for ω^n -largeness.

Lemma 7.2. Fix $n > 0$. Let X be minimal for ω^n -largeness. Then it admits a unique ω^n -decomposition $X_0 < \dots < X_a$. Moreover, $a = \min X - 1$, $\{\min X\} \cup X_0 \cup \dots \cup X_a = X$ and for every $i < a$, X_i is minimal for ω^{n-1} -largeness.

Proof. Let X be minimal for ω^n -largeness for some $n > 0$. Let $X_0 < \dots < X_a$ be an ω^n -decomposition of X .

First, we claim that $a = \min X - 1$. Indeed, if $a > \min X - 1$, then $X \setminus X_a$ would be ω^n -large, contradicting minimality of X for ω^n -largeness.

Second, we claim that X_i is minimal for ω^{n-1} -largeness. Suppose not. Let $x \in X_i$ be such that $X_i \setminus \{x\}$ is ω^{n-1} -large. Then $X \setminus \{x\}$ would be ω^n -large, again contradicting minimality of X for ω^n -largeness.

Third, let us prove that $\{\min X\} \cup X_0 \cup \dots \cup X_a = X$. Suppose there is some $x \in X \setminus (\{\min X\} \cup X_0 \cup \dots \cup X_a)$. Then once again, $X \setminus \{x\}$ would be ω^n -large.

Last, assume there is another ω^n -decomposition $Y_0 < \dots < Y_a$ of X . Then, by induction over $i \leq a$, we prove that $X_i = Y_i$. Indeed, assuming that $X_j = Y_j$ for every $j < i$, since $X = \{\min X\} \cup X_0 \cup \dots \cup X_a = \{\min X\} \cup Y_0 \cup \dots \cup Y_a$, then either $X_i \subseteq Y_i$, or $Y_i \subseteq X_i$. By minimality of X_i and Y_i for ω^{n-1} -largeness, $X_i = Y_i$. This completes the proof of Lemma 7.2. \square

The previous lemma justifies the following definition:

Definition 7.3. Fix a set X which is minimal for ω^n -largeness. The canonical n -block of X is X itself. For $c < n$, the canonical c -blocks of X are the canonical c -blocks of the sets X_i , where $X_0 < \dots < X_{\min X-1}$ is the unique ω^n -decomposition of X .

Definition 7.4. Let X minimal for ω^n -largeness for some $n \geq 1$. The 0-blockfree subset of X is the set X minus all the elements belonging to a canonical 0-block of X .

Lemma 7.5. Let X minimal for ω^n -largeness for some $n \geq 1$. Its 0-blockfree subset is a minimal ω^{n-1} -large subset of X .

Proof. Proceed by induction on n :

Case $n = 1$: Let X be ω^1 -large, then its 0-blockfree subset is equal to $\{\min X\}$ which is minimal ω^0 -large.

Case $n \geq 2$: Assume the property to be true at rank $n - 1$, let $X_0 < \dots < X_{\min X-1}$ be the canonical ω^n -decomposition of X into ω^{n-1} -large sets. Let X'_i be the 0-blockfree subset of X_i for every $i < \min X$. By the inductive hypothesis, every X'_i is a minimal ω^{n-2} -large subset of X_i , therefore the 0-blockfree subset X' of X is ω^{n-1} -large and $X'_0 < \dots < X'_{\min X-1}$ is the canonical ω^{n-1} -decomposition of X' into ω^{n-2} -large sets. \square

Definition 7.6. Let X be minimal for ω^n -largeness. Consider $\phi_X(x, y, c)$ a Σ_0 formula that is true if and only if x and y are in the same canonical c -block of X for $c \leq n$.

Let $\theta_X(x, y, z)$ be the following Σ_0 formula:

$$(x \in X \wedge z \in X \wedge z \geq y) \rightarrow \exists c \leq n, (\phi_X(y, z, c) \wedge \neg \phi_X(x, y, c) \wedge y > x \wedge y \in X)$$

Let T_X be the formula $\forall x \exists y \forall z \theta_X(x, y, z)$.

All of these formulas are Σ_0 , since there are only finitely many elements of X to consider. Note that n is uniquely determined by X : it is the largest integer less or equal to $\max X$ such that X is ω^n -large. Being ω^n -large is a Σ_0 predicate in X and n , so the formula ϕ_X is Σ_0 uniformly in X .

Lemma 7.7. *Let X be minimal for ω^n -largeness for some $n \geq 1$. For every subsets $A < B$ of X , A and B are T_X -apart if and only if $\theta_X(\max A, \min B, \max B)$ holds.*

Proof. If A and B are T_X -apart, then there exists some $y \leq \min B$ such that $\theta_X(\max A, y, \max B)$. Since $\max A \in X$, $\max B \in X$ and $\max B \geq y$, unfolding the definition of θ_X yield $y \in X$, $y > \max A$ and there exists some $c \leq n$ such that $\phi_X(y, \max B, c)$ and $\neg\phi_X(\max A, y, c)$ holds. Since $\min B \in [y, \max B]$ and $\min B \in X$, $\phi_X(\min B, \max B, c)$ also holds, and since $y \in [\max A, \min B]$, $\neg\phi_X(\max A, \min B, c)$ holds. So $\theta_X(\max A, \min B, \max B)$ holds.

Conversely, if $\theta_X(\max A, \min B, \max B)$ holds: since $\max A \in X$, $\max B \in X$ and $\max B \geq \min B$, there exists some $c \leq n$ such that $\phi_X(\min B, \max B, c)$ and $\neg\phi_X(\max A, \min B, c)$ holds. To show that A and B are T_X -apart, it is sufficient to show that $\theta_X(x, \min B, z)$ holds for every $x \leq \max A$ and $z \leq \max B$. If $x \notin X$ or $z \notin Z$ or $z < \min B$ then $\theta_X(x, \min B, z)$ holds, so assume $x \in X$ and $z \in Z$ and $\min B \leq z$. Since $x \leq \max A$ and $\neg\phi_X(\max A, \min B, c)$ holds, then $\neg\phi_X(x, \min B, c)$ holds and since $z \leq \max B$ and $\phi_X(\min B, \max B, c)$ holds, then $\phi_X(\min B, z, c)$ holds. Therefore, $\forall x \leq \max A \exists y \leq \min B \forall z \leq \max B \theta_X(x, y, z)$ holds. This completes the proof of Lemma 7.7. \square

Lemma 7.8. *Let X be minimal for ω^n -largeness for some $n \geq 1$. Let Y be a canonical c -block of X for some $c \leq n$. Then θ_X and θ_Y are equivalent for elements of Y .*

Proof. Let $x, y \in Y$. Then for every $d \leq c$, $\phi_X(x, y, d)$ holds if and only if $\phi_Y(x, y, d)$ holds as the canonical d -block of Y are exactly the canonical d -block of X contained in Y .

Let $x, y, z \in Y$, if $z < y$ then $\theta_X(x, y, z)$ and $\theta_Y(x, y, z)$ hold. So assume $y \leq z$, $\theta_X(x, y, z)$ holds if and only if there exists some $d \leq n$ such that $\phi_X(y, z, d)$ and $\neg\phi_X(x, y, d)$ and $y > x$. Since $\phi_X(x, y, c)$ holds (as Y is a canonical c -block) then $d < c$. So $\theta_X(x, y, z)$ holds if and only if $\phi_Y(y, z, d)$ and $\neg\phi_Y(x, y, d)$ and $y > x$ which is equivalent to $\theta_Y(x, y, z)$. This completes the proof of Lemma 7.8. \square

Combining Lemma 7.7 and Lemma 7.8 yield the following corollaries:

Corollary 7.9. *Let X be minimal for ω^n -largeness for some $n \geq 1$ and Y a canonical c -block of X for some $c \leq n$.*

1. *Two subsets $A < B$ of Y are T_Y -apart, if and only if they are T_X -apart.*
2. *For $k \leq c$, the ω^k -large(T_Y) subsets of Y are exactly the ω^k -large(T_X) subsets of Y .*

Lemma 7.10. *Let X be minimal for ω^n -largeness for some $n \geq 1$. Let X' be the 0-blockfree subset of X . Then θ_X and $\theta_{X'}$ are equivalent for elements of X' .*

Proof. Let $x, y \in X'$. Then for every $1 \leq c \leq n$, $\phi_X(x, y, c)$ holds if and only if $\phi_{X'}(x, y, c - 1)$ holds, as the canonical $(c - 1)$ -blocks of X' are exactly the canonical c -block of X minus the elements belonging to a canonical 0-block.

Let $x, y, z \in X'$, if $z < y$ then $\theta_X(x, y, z)$ and $\theta_{X'}(x, y, z)$ hold. So assume $y \leq z$, $\theta_X(x, y, z)$ holds if and only if there exists some $c \leq n$ such that $\phi_X(y, z, c)$ and $\neg\phi_X(x, y, c)$ and $y > x$. Since $x, y \in X'$, c cannot be equal to 0. So $\theta_X(x, y, z)$ holds if and only if $y > x$ and $\phi_{X'}(y, z, c - 1)$ and $\neg\phi_{X'}(x, y, c - 1)$ holds, which is equivalent to $\theta_{X'}(x, y, z)$. This completes the proof of Lemma 7.10. \square

Combining Lemma 7.7 and Lemma 7.10 yield the following corollaries:

Corollary 7.11. *Let X be minimal for ω^n -largeness for some $n \geq 1$ and X' be its 0-blockfree subset.*

1. *Two subsets $A < B$ of X' are $T_{X'}$ -apart, if and only if they are T_X -apart.*
2. *For $k \leq n - 1$, the ω^k -large($T_{X'}$) subsets of X' are exactly the ω^k -large(T_X) subsets of X' .*

Lemma 7.12. *Let X be minimal for ω^n -largeness for some $n \geq 1$. Then X is ω^n -large(T_X).*

Proof. For this, it suffices to prove that for every $a < n$, any two canonical a -blocks $Y < Z$ of X are T_X -apart. Let $Y < Z$ be two such blocks. Then $\phi_X(\min Z, \max Z, a)$ holds since $\min Z$ and $\max Z$ belong to the same a -block, and $\phi_X(\max Y, \min Z, a)$ does not hold since $\max Y$ is not in Z . It follows that $\theta_X(\max Y, \min Z, \max Z)$ holds, so by Lemma 7.7, Y and X are T_X -apart. This completes the proof of Lemma 7.12. \square

Lemma 7.13. *Let X be minimal for ω^n -largeness for some $n \geq 1$. Let $X_0 < \dots < X_{\min X - 1}$ be the canonical ω^n -decomposition of X into ω^{n-1} -large sets. If $A < B$ are two T_X -apart subsets of X , then $B \subseteq X_i$ for some i .*

Proof. Since A and B are T_X -apart, then by Lemma 7.7, $\theta_X(\max A, \min B, \max B)$ holds. Since $\max A \in X$, $\max B \in X$ and $\max B \geq \min B$, unfolding the definition of θ_X yield that there exists some $c \leq n$ such that $\phi_X(\min B, \max B, c)$ and $\neg\phi_X(\max A, \min B, c)$ holds. Since $\phi(\max A, \min B, n)$ holds (as $\max A, \min B \in X$), c cannot be equal to n . On the other hand, $\min B$ and $\max B$ are in the same canonical c -block of X , so there are in the same canonical $(n - 1)$ -block. So B is included in X_i for some i . This completes the proof of Lemma 7.13. \square

In the following proposition, recall that if X is minimal for ω^{2n-1} -largeness, then it is ω^{2n-1} -large(T_X) by Lemma 7.12. Thus, the proposition gives us an example of an ω^{2n-1} -large(T_X) set which is not ω^n -large(T_X, RT_2^1).

Proposition 7.14. *Let X be minimal for ω^{2n-1} -largeness for some $n \geq 1$. There exists a coloring $f_X : X \rightarrow 2$ such that there is no f_X -homogeneous ω^n -large(T_X) subset of X .*

Proof. Let f_X be the following 2-coloring of X : for $x \in X$, consider the smallest c such that x is in a c -canonical block of X , then color x with the parity of c .

Proceed by induction on n :

Case $n = 1$: Let X be minimal for ω -largeness (in other words, $|X| = \min X + 1$). Then $f_X(\min X) = 1$ and $f_X(x) = 0$ for every other element of X . So there is no f_X -homogeneous ω -large(T_X) subset of X by minimality of X .

Case n : Assume the property to be true at rank $n - 1$, and let X be minimal for ω^{2n-1} -largeness. Let $X_0 < \dots < X_{\min X - 1}$ be the canonical ω^{2n-1} -decomposition of X into ω^{2n-2} -large sets.

Assume by contradiction that there exists an f_X -homogeneous ω^n -large(T_X) subset $Y \subseteq X$, we can assume Y to be minimal for ω^n -largeness(T_X). Let $Y_0 < \dots < Y_{\min Y - 1}$ be the canonical ω^n -decomposition of Y into ω^{n-1} -large sets.

There are two cases:

- Y is f_X homogeneous for the color 0: Since $f_X(\min X) = 1$, $\min X \notin Y$, so $\min Y > \min X$. By the finite pigeonhole principle, there exists some $i < \min X$, such that X_i contains two elements of the form $\min Y_j$ for some $j < \min Y$. Since $Y_0 < \dots < Y_{\min Y - 1}$, we can assume these two elements to be of the form $\min Y_j, \min Y_{j+1}$, so there is some $j < \min Y - 1$ such that $Y_j \subseteq X_i$ and $\min Y_{j+1} \in X_i$. By Lemma 7.13, since Y_j and Y_{j+1} are T_X -apart, Y_{j+1} must also be included in X_i .

Let $X_{i,0} < \dots < X_{i,\min X_i - 1}$ be the canonical ω^{2n-2} -decomposition of X_i into ω^{2n-3} -large sets. By Corollary 7.9, since Y_j, Y_{j+1} are T_X -apart and subsets of X_i , they are also T_{X_i} -apart. So, by Lemma 7.13 applied to X_i , $Y_{j+1} \subseteq X_{i,i'}$ for some $i' < \min X_i - 1$. By Corollary 7.9, since Y_{j+1} is ω^{n-1} -large(T_X) then it is ω^{n-1} -large($T_{X_{i,i'}}$). But $f_X \upharpoonright X_{i,i'}$ is equal to $f_{X_{i,i'}}$, so Y_{j+1} is an ω^{n-1} -large($T_{X_{i,i'}}$), $f_{X_{i,i'}}$ -homogeneous subset of $X_{i,i'}$, contradicting the induction hypothesis.

- Y is f_X homogeneous for the color 1: Let X' be the 0-blockfree subset of X . Then, by Lemma 7.5, X' is a minimal ω^{2n-2} -large subset of X and its canonical decomposition $X'_0 < \dots < X'_{\min X - 1}$ satisfies that X'_i is equal to X_i minus all the elements belonging to a canonical 0-block. Furthermore, $Y \subseteq X'$ (since all the elements belonging to a canonical 0-block of X are 0-colored).

If $\max Y_0 \in X_0$ then $Y_0 \subseteq X_0$, and otherwise, by the finite pigeonhole principle, there exists some $1 \leq i < \min X$ such that X_i contains two elements of the form $\max Y_j$. Since $Y_0 < \dots < Y_{\min Y - 1}$, in both cases we have $Y_j \subseteq X_i$ for some $j < \min Y$ and $i < \min X$ and therefore $Y_j \subseteq X'_i$. By Corollary 7.11, Y_j is an ω^{n-1} -large($T_{X'_i}$) subset of X'_i . $f_X \upharpoonright X'_i$ is equal to $1 - f_{X'_i}$ (a canonical c -block of X became a canonical $(c - 1)$ -block

of X' when we get rid of the 0-canonical elements), so Y_j is an ω^{n-1} -large($T_{X'_i}$), $f_{X'_i}$ -homogeneous (for the color 0) subset of X'_i , contradicting the induction hypothesis.

We conclude by induction. This completes the proof of Proposition 7.14. \square

8 Open questions

There exists a close connection between explicit bounds computation for largeness, and proof speedup theorems. In particular Kołodziejczyk, Wong and Yokoyama [15] proved that Ramsey's theorem for pairs and two colors has at most polynomial speedup over RCA_0 for $\forall\Pi_3^0$ sentences, using the fact that ω^{300n} -largeness is sufficient to obtain a homogeneous ω^n -large set for every instance of RT_2^2 . The exponential bounds of largeness(T) for Ramsey's theorem for pairs yields the following natural questions:

Question 8.1. *Does RT_2^2 admit exponential proof speedup over $\text{RCA}_0 + \text{BS}_2^0$?*

Question 8.2. *Is there a polynomial p such that for every n , every $\omega^{p(n)}$ -large(T) set is ω^n -large(T, RT_2^2)?*

The lower bound for the pigeonhole principle uses a formula T_X which depends on the considered set X .

Question 8.3. *Is there a Π_3^0 formula T such that for every $n \in \omega$, every ω^n -large(T, RT_2^1) set is ω^{2n-1} -large(T)?*

We note that Question 8.1 is essential for the original question on Π_1^1 -conservation (Question 1.1). By the discussion of [6, Section 5] and an upcoming paper by Ikari and Yokoyama, if $\text{RCA}_0 + \text{RT}_2^2$ is Π_3^0 -conservative over $\text{RCA}_0 + \text{BS}_2^0$, then there exists a poly-time proof transformation between $\text{RCA}_0 + \text{RT}_2^2$ and $\text{RCA}_0 + \text{BS}_2^0$ for Π_1^1 -consequences. Thus, a positive answer to Question 8.1 implies that our Π_4^0 -conservation is the best possible.

Acknowledgement

The authors are thankful to Leszek Kołodziejczyk for insightful comments and discussions.

References

- [1] Andrey Bovykin and Andreas Weiermann. The strength of infinitary ramseyan principles can be accessed by their densities. *Annals of Pure and Applied Logic*, 168(9):1700–1709, 2017.
- [2] Samuel R. Buss, editor. *Handbook of proof theory*, volume 137 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 1998.

- [3] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey’s theorem for pairs. *J. Symbolic Logic*, 66(1):1–55, 2001.
- [4] C. T. Chong, Theodore A. Slaman, and Yue Yang. Π_1^1 -conservation of combinatorial principles weaker than Ramsey’s theorem for pairs. *Adv. Math.*, 230(3):1060–1077, 2012.
- [5] C. T. Chong, Theodore A. Slaman, and Yue Yang. The inductive strength of Ramsey’s Theorem for Pairs. *Adv. Math.*, 308:121–141, 2017.
- [6] Marta Fiori-Carones, Leszek Aleksander Kołodziejczyk, Tin Lok Wong, and Keita Yokoyama. An isomorphism theorem for models of Weak König’s Lemma without primitive recursion. *Journal of the European Mathematical Society*, 2021. To appear.
- [7] Harvey Friedman. Some systems of second order arithmetic and their use. In *Proceedings of the International Congress of Mathematicians (Vancouver, B.C., 1974)*, Vol. 1, pages 235–242. Canad. Math. Congr., Montreal, QC, 1975.
- [8] Petr Hájek. Interpretability and fragments of arithmetic. In P. Clote and J. Krajíček, editors, *Arithmetic, Proof Theory and Computational Complexity*, pages 185–196. Oxford, Clarendon Press, 1993.
- [9] Petr Hájek and Pavel Pudlák. *Metamathematics of first-order arithmetic*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1998. Second printing.
- [10] Denis R. Hirschfeldt. *Slicing the truth*, volume 28 of *Lecture Notes Series. Institute for Mathematical Sciences. National University of Singapore*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015. On the computable and reverse mathematics of combinatorial principles, Edited and with a foreword by Chitat Chong, Qi Feng, Theodore A. Slaman, W. Hugh Woodin and Yue Yang.
- [11] Jeffrey Lynn Hirst. *COMBINATORICS IN SUBSYSTEMS OF SECOND ORDER ARITHMETIC*. ProQuest LLC, Ann Arbor, MI, 1987. Thesis (Ph.D.)—The Pennsylvania State University.
- [12] Richard Kaye. *Models of Peano arithmetic*, volume 15 of *Oxford Logic Guides*. The Clarendon Press, Oxford University Press, New York, 1991. Oxford Science Publications.
- [13] Jussi Ketonen and Robert Solovay. Rapidly growing Ramsey functions. *Ann. of Math. (2)*, 113(2):267–314, 1981.
- [14] L. A. S. Kirby and J. B. Paris. Initial segments of models of Peano’s axioms. In *Set theory and hierarchy theory, V (Proc. Third Conf., Bierutowice, 1976)*, Lecture Notes in Math., Vol. 619, pages 211–226. Springer, Berlin, 1977.

- [15] Leszek Aleksander Kołodziejczyk, Tin Lok Wong, and Keita Yokoyama. Ramsey’s theorem for pairs, collection, and proof size. *Journal of Mathematical Logic*, page 2350007, 2023.
- [16] Leszek Aleksander Kołodziejczyk and Keita Yokoyama. Some upper bounds on ordinal-valued Ramsey numbers for colourings of pairs. *Selecta Math. (N.S.)*, 26(4):Paper No. 56, 18, 2020.
- [17] Leszek Aleksander Kołodziejczyk and Keita Yokoyama. In search of the first-order part of Ramsey’s theorem for pairs. In *Connecting with computability*, volume 12813 of *Lecture Notes in Comput. Sci.*, pages 297–307. Springer, Cham, [2021] ©2021.
- [18] Jiayi Liu. RT22 does not imply WKL0. *The Journal of Symbolic Logic*, 77(2):609–620, 2012.
- [19] Benoit Monin and Ludovic Patey. SRT_2^2 does not imply RT_2^2 in ω -models. *Adv. Math.*, 389:Paper No. 107903, 32, 2021.
- [20] J. B. Paris. A hierarchy of cuts in models of arithmetic. In *Model theory of algebra and arithmetic (Proc. Conf., Karpacz, 1979)*, volume 834 of *Lecture Notes in Math.*, pages 312–337. Springer, Berlin, 1980.
- [21] Ludovic Patey. Ramsey-like theorems and moduli of computation. *J. Symb. Log.*, 87(1):72–108, 2022.
- [22] Ludovic Patey and Keita Yokoyama. The proof-theoretic strength of Ramsey’s theorem for pairs and two colors. *Adv. Math.*, 330:1034–1070, 2018.
- [23] Dana Scott. Algebras of sets binumerable in complete extensions of arithmetic. In *Proc. Sympos. Pure Math., Vol. V*, pages 117–121. Amer. Math. Soc., Providence, RI, 1962.
- [24] David Seetapun and Theodore A. Slaman. On the strength of Ramsey’s theorem. volume 36, pages 570–582. 1995. Special Issue: Models of arithmetic.
- [25] Keita Yokoyama. On Π_1^1 conservativity of Π_2^1 theories in second order arithmetic. In C. T. Chong et al., editor, *Proceedings of the 10th Asian Logic Conference*, pages 375–386. World Scientific, 2009.
- [26] Keita Yokoyama. On conservativity for theories in second order arithmetic. In *Proceedings of the 10th Asian Logic Conference*, pages 375–386. World Scientific, 2010.
- [27] Keita Yokoyama. Notes on the first-order part of Ramsey’s theorem for pairs (proof theory and complexity). 2013.